

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 019 364 B1

(12)

EUROPEAN PATENT SPECIFICATION(45) Date of publication and mention
of the grant of the patent:**09.06.2004 Bulletin 2004/24**(51) Int Cl.7: **C07C 235/00**(86) International application number:
PCT/US1998/017327(21) Application number: **98944476.5**(22) Date of filing: **24.08.1998**(87) International publication number:
WO 1999/011606 (11.03.1999 Gazette 1999/10)**(54) INHIBITORS OF PROTEIN TYROSINE PHOSPHATASE**

PROTEINTYROSINPHOSPHATASE INHIBITOREN

INHIBITEURS DE LA PROTEINE TYROSINE PHOSPHATASE

(84) Designated Contracting States:

**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**

Designated Extension States:

AL LT LV MK RO SI(30) Priority: **28.08.1997 US 57730 P**

(43) Date of publication of application:

19.07.2000 Bulletin 2000/29(73) Proprietor: **BIOVITRUM AB****112 76 Stockholm (SE)**

(72) Inventors:

- **LARSEN, Scott, D.**
Kalamazoo, MI 49009 (US)
- **MAY, Paul, D.**
Richland, MI 49083 (US)
- **BLEASDALE, John**
Portage, MI 49024 (US)
- **LILJEBRIS, Charlotta**
S-752 67 Uppsala (SE)
- **SCHOSTAREZ, Heinrich, Josef**
Portage, MI 49024 (US)
- **BARF, Tjeerd**
S-753 34 Uppsala (SE)

(74) Representative: **Höglund, Lars et al****Biovitrum AB****112 76 Stockholm (SE)**

(56) References cited:

WO-A-96/30332**WO-A-96/33968****WO-A-96/38415****WO-A-98/07699****DD-A- 151 304****DE-A- 2 320 387**

- **PAUL S. CHARIFSON ET AL.: "Peptide ligands of pp60c-src SH2 Domains: A thermodynamic and structural study" BIOCHEMISTRY., vol. 36, 1997, pages 6283-6293, XP002087254 EASTON, PA US**
- **KING F D: "BIOISOSTERES, CONFORMATIONAL RESTRICTION, AND PRO-DRUGS - CASE HISTORY: AN EXAMPLE OF A CONFORMATIONAL RESTRICTION APPROACH" 1994, MEDICINAL CHEMISTRY: PRINCIPLES AND PRACTICE, PAGE(S) 206 - 209, KING F D XP002033086 see page 208**

Remarks:

The file contains technical information submitted after the application was filed and not included in this specification

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description

[0001] The present invention comprises small molecular weight, non-peptidic inhibitors of Protein Tyrosine Phosphatase 1 (PTP1) which are useful for the treatment and/or prevention of Non-Insulin Dependent Diabetes Mellitus (NIDDM).

BACKGROUND OF THE INVENTION

[0002] The mechanism of insulin action depends critically upon the phosphorylation of tyrosine residues in several proteins in the insulin signalling cascade. Enzymes that dephosphorylate these proteins, protein tyrosine phosphatases (PTPs), are important negative regulators of insulin action. Therefore, the use of specific PTP inhibitors may therapeutically enhance insulin action.

[0003] The insulin resistance that is central to noninsulin-dependent diabetes mellitus (NIDDM) appears to involve a defect in an early process in insulin signal transduction rather than a structural defect in the insulin receptor itself. (J.M. Olefsky, W.T. Garvey, R.R. Henry, D. Brillon, S. Matthai and G.R. Freidenberg, G.R. (1988).) Cellular mechanisms of insulin resistance in non-insulin-dependent (Type II) diabetes. (Am. J. Med. 85: Suppl. 5A, 86-105.) A drug that improved insulin sensitivity would have several advantages over traditional therapy of NIDDM using sulfonylureas, which do not alleviate insulin resistance but instead compensate by increasing insulin secretion.

[0004] The binding of insulin to the α -subunits of the insulin receptor permits the β -subunits to catalyze phosphorylation of target proteins on tyrosine residues. There are 22 tyrosine residues in each insulin receptor β -subunit itself and autophosphorylation of at least 6 of these tyrosines, in 3 distinct domains, is known to be involved in insulin action. (C.R. Kahn (1994) Insulin action, diabetogenes, and the cause of type II diabetes. Diabetes 43: 1066-1084.) Autophosphorylation of Tyr⁹⁶⁰ in the juxtamembrane domain is important for receptor internalization and for the interaction of the receptor with downstream signalling molecules such as insulin receptor substrate 1 (IRS-1). (T.J. O'Neill, A. Craparo and T.A. Gustafson (1994) Characterization of an interaction between insulin receptor substrate 1 and the insulin receptor by using the two-hybrid system. Mol. Cell Biol. 14: 6433-6442.) Autophosphorylation of tyrosine residues 1146, 1150 and 1151 in the regulatory domain permits continued tyrosine kinase activity of β -subunits, even after insulin has dissociated from the α -subunits, and activates the kinase toward other protein substrates. (R. Herrera and O.M. Rosen (1986) Autophosphorylation of the insulin receptor *in vitro*: designation of phosphorylation sites and correlation with receptor kinase activation. J. Biol. Chem. 261: 11980-11985.) Deletion of autophosphorylation sites at Tyr¹³¹⁶ and Tyr¹³²² in the C-terminal domain attenuates the metabolic actions of insulin, but augments its mitogenic actions. (H. Maegawa, D. McClain, G. Freidenberg, J. Olefsky, M. Napier, T. Lipari, T. Dull, J. Lee, and A. Ullrich (1988) Properties of a human insulin receptor with a COOH-terminal truncation. II. Truncated receptors have normal kinase activity but are defective in signalling metabolic effects. J. Biol. Chem. 263: 8912-8917.) (Y. Takata, N.J.G. Webster, and J.M. Olefsky (1991) Mutation of the two carboxyl-terminal tyrosines results in an insulin receptor with normal metabolic signalling but enhanced mitogenic signalling properties. J. Biol. Chem. 266: 9135-9139.) Dephosphorylation of these autophosphorylated sites occurs rapidly *in vivo*, suggesting that a protein tyrosine phosphatase (PTPase) is involved in terminating insulin action. A compound that inhibited this PTPase, therefore, should potentiate insulin action. Indeed, vanadate potentiates insulin action, at least in part, by such a mechanism (Y. Schechter (1990). Insulin-mimetic effects of vanadate. Possible implications for future treatment of diabetes. Diabetes 39: 1-5.) The PTPase(s) that act on the insulin receptor, however, has not been identified definitively.

[0005] It has been estimated that the human genome encodes as many as 500 PTP enzymes (T. Hunter (1995) Protein kinases and phosphatases: The Yin and Yang of protein phosphorylation and signalling. Cell 80:225-236), but less than 100 have been identified and have been grouped into 4 sub-families (E.A. Fauman and M.A. Saper (1996) Structure and function of the protein tyrosine phosphatases. Trends Biochem. Sci. 21:413-417.) Members of the tyrosine-specific PTP sub-family are further divided into the receptor PTPases (such as CD45 and LAR) which typically have a large variable extracellular domain, a single transmembrane spanning region, and two intracellular phosphatase catalytic domains and the non-receptor PTPases. This latter group includes PTP that resemble PTP1. (D.A. Pot and J.E. Dixon (1992) A thousand and two protein tyrosine phosphatases. Biochim. Biophys. Acta 1136: 35-43.) There is data to support the proposition that the insulin receptor PTPase may be PTP1-like. For instance, an insulin-dependent association of PTP1 with insulin receptors has been described. (D. Bandyopadhyay, A. Kursari, K.A. Kenner, F. Liu, J. Chernoff, T.A. Gustafson, J. Kusari (1997) Protein-tyrosine phosphatase 1B complexes with the insulin receptor *in vivo* and is tyrosine-phosphorylated in the presence of insulin. J.Biol.Chem. 272: 1639-1645; and L. Seely, et al. (1996) Protein tyrosine phosphatase 1B interacts with the activated insulin receptor. Diabetes 45:1379.) Furthermore, PTP1 dephosphorylates purified insulin receptors sequentially in the order observed *in vivo* (i.e., Tyr¹¹⁵⁰ = Tyr¹¹⁵¹ > Tyr¹¹⁴⁶), (C. Ramachandran, R. Aebersold, N. Tonks and D.A. Pot (1992) Sequential dephosphorylation of a multiply phosphorylated insulin receptor peptide by protein tyrosine phosphatases. Biochemistry 31: 4232-4238) and insulin acutely increases PTP1 mRNA in hepatoma cells. (N. Hashimoto and B.J. Goldstein (1992) Differential regulation of mRNAs

encoding three protein-tyrosine phosphatases by insulin and activation of protein kinase C. Biochem. Biophys. Res. Commun. 188: 1305-1311.) Insulin resistance induced in Rat 1 fibroblasts by high glucose (27 mM) is preceded by an approximate doubling of cytosolic PTP1 activity that is blocked by the insulin-sensitizer, pioglitazone. (H. Maegawa, R. Ide, M. Hasegawa, S. Ugi, K. Egawa, M. Iwanishi, R. Kikkawa, Y. Shigeta, and A. Kashiwagi (1995) Thiazolidinedione derivatives ameliorate high glucose-induced insulin resistance via the normalization of protein tyrosine phosphatase activities. J. Biol. Chem. 270: 7724-7730.) Thus, a specific inhibitor of PTP1 could be used to potentiate insulin action. While there are no known small molecules that specifically inhibit PTP1, it was found that osmotic loading of hepatoma cells with neutralizing antibodies against PTP1b (the human homologue of rat PTP1) resulted in increased autophosphorylation of insulin receptors and phosphorylation of IRS-1 in response to insulin. (F. Ahmad, P.-M. Li, J. Meyerovitch, and B.J. Goldstein (1995) Osmotic loading of neutralizing antibodies demonstrates a role for PTPase 1B in negative regulation of the insulin signalling pathway. Diabetes 44: Suppl. 1 104A.) See also B.J. Goldstein (1993) Regulation of insulin receptor signaling by protein-tyrosine dephosphorylation. Receptor 3: 1-15.)

INFORMATION DISCLOSURE

[0006] International Publication No. WO 96/30332, "O-Malonyltyrosyl Compounds, O-Malonyltyrosyl Compound-Containing Peptides, and Uses thereof," published 3 October 1996, disclose non-phosphorus containing O-malonyltyrosyl compounds, derivatives thereof, uses of the O-malonyltyrosyl compounds in the synthesis of peptides, and O-malonyltyrosyl compound-containing peptides. The O-malonyltyrosyl compounds and O-malonyltyrosyl compound-containing peptides are disclosed as being useful as inhibitors of protein-tyrosine phosphatase; however, no specific non-peptidic compounds or data is disclosed. International Publication No. WO 96/23813, "Peptides and Compounds that Bind to SH2 Domains," published 8 August 1996, discloses tyrosine-containing peptides and compounds which bind to the SH2 domain or domains of various proteins, as well as methods for identifying such peptides and compounds. These peptides and compounds have application as agonists and antagonists of SH2 domain containing proteins, and as diagnostic or therapeutic agents for the diagnosis or treatment of disease conditions.

[0007] International Publication No. WO 96/40113, "Phosphatase Inhibitors," published 19 December 1996, discloses heterocyclic nitrogen containing compounds, such as nitropyridine or nitrothiazole, capable of inhibiting protein tyrosine phosphatase activity. Such molecules are disclosed as being useful to modulate or regulate signal transduction by inhibiting protein tyrosine phosphatase activity and to treat various disease states including diabetes mellitus.

[0008] International Publication No. WO 96/40109, "Methods of Inhibiting Phosphatase Activity and Treatment of Disorders Associated Therewith Using Naphthopyrones and Derivatives Thereof," published 19 December 1996, discloses the use of naphthopyrone compounds to inhibit protein tyrosine phosphatase activity. Such compounds are disclosed as being useful to modulate or regulate signal transduction by inhibiting protein tyrosine phosphatase activity and to treat various disease states including diabetes mellitus.

[0009] Charifson et al. (1997) Biochemistry 36: 6283-6293, discloses peptide ligands of pp60^{c-src} SH2 domains.


[0010] WO 96/33968 discloses water-soluble hydroxamic acid derivatives which inhibit matrix metalloproteinases and/or tumor necrosis factor- α converting enzyme.

[0011] WO 96/38415 discloses 2-amino-3-phenylpropionic acid derivatives useful as hypoglycemic agents and for lowering blood lipid levels.

[0012] The compounds of the present invention have surprising activity in that they are small molecular weight and non-peptidic compounds.

SUMMARY OF THE INVENTION

[0013] A compound of formula I



wherein G¹ is

- a) $-R^2$, or
b) $-NR^8 R^4$;

wherein R^2 is

- a) $-C_1-C_{10}$ alkyl optionally substituted with one or two $-CO_2R^5$ bonded to the same or different carbon atoms or with one $-CO-NH_2$,
b) $-C_3-C_8$ cycloalkyl optionally substituted with one $-CO_2R^5$,
c) $-C_0-C_6$ alkyl-phenyl optionally substituted with one or two $-CO_2R^5$ bonded to the same or different carbon atoms or with $-CH_2CH(CO_2R^5)_2$,
d) $-CH(R^7)NHXR^6$, or
e)



wherein R^3 is

- a) $-C_1-C_{12}$ alkyl, optionally substituted with one to three $-O-$, $-S-$, $-N-$, $-O-C_1-C_4$ alkyl, $-S-C_1-C_4$ alkyl, $-O-G^3$, $-S-G^3$, or $-OH$,
b) $-C_1-C_4$ alkyl $-C_3-C_6$ cycloalkyl,
c) $-C_2-C_{12}$ alkenyl,
d) $-C_3-C_{12}$ alkynyl,
e) $-C_0-C_{10}$ alkyl(G^3)_n wherein alkyl is optionally substituted with one to three $-O-$, $-S-$ or $-N-$, or
f) $-CH(CONH_2)C_1-C_{12}$ alkyl;

wherein R^4 is

- a) $-H$,
b) up to C_{18} alkyl or alkenyl, or
c) $-C_0-C_6$ -alkyl- G^3 ;

wherein R^5 is

- a) $-H$,
b) $-C_1-C_{10}$ alkyl, or
c) $-C_1-C_5$ alkyl-phenyl;

wherein R^6 is

- a) C_1-C_{10} alkyl,
b) C_0-C_6 alkyl- G^3 ,
c) C_1-C_6 alkyl $CONH_2$,
d) C_1-C_6 alkyl $NHCO_2R^5$,
e) C_1-C_6 alkyl- OR^5 ,
f) C_1-C_6 alkyl- $NHSO_2Me$,
g) C_1-C_6 alkyl- $O-G^3$,
h) C_1-C_6 alkyl- $S-G^3$, or
i) $-C_1-C_6$ alkyl- CO_2R^5 ;

wherein R^7 is

- a) $-H$,
b) $-C_1-C_6$ alkyl- G^3
c) $-C_1-C_6$ alkyl- CO_2R^5
d) C_1-C_6 alkyl $CONH_2$,
e) C_1-C_6 alkyl $NHCO_2R^5$,

- f) C₁-C₁₀ alkyl,
- g) up to C₁₀ cycloalkyl,
- h) -C₁-C₆ alkyl- SR⁵, or
- i) -C₁-C₆ alkyl- S(=O)R⁵;

wherein R⁸ is

- a) C₀-C₆ alkyl-G³,
- b) CH(R⁷)CO₂R⁵,
- c) CH(R⁷)CH₂CO₂R⁵, or
- d) CH(R⁷)CONHCH₂CO₂R⁵;

wherein G³ is

- a) phenyl substituted by zero (0) to three (3) R⁹, or
- b) naphthyl substituted by zero (0) to three (3) R⁹;

wherein R⁹ may be any of the following:

- a) C₁-C₈ alkyl substituted by zero (0) to three (3) halo,
- b) C₂-C₈ alkenyl,
- c) OH,
- d) O-C₁-C₅ alkyl,
- e) O-C₀-C₅ alkyl-phenyl
- e) -(CH₂)_n-O-C₁-C₅ alkyl substituted by zero (0) to three (3) hydroxy,
- f) -(CH₂)_n-O-C₂-C₇ alkenyl substituted by zero (0) to three (3) hydroxy,
- g) halo,
- h) NH₂,
- i) amino-C₁-C₅ alkyl,
- j) mono-or di-C₁-C₅ alkylamino,
- k) -C(O)-C₁-C₅ alkyl,
- l) -CHO,
- m) -C(O)-Co -C₅ alkyl-phenyl
- n) -COOR⁵
- o) -CON(R⁵)₂,
- p) -C₃-C₇ cycloalkyl,
- q) -NO₂,
- r) -CN,
- s) -SO₃H,
- t) -SO₂N(R⁵)₂,
- u) -O[(CH₂)₂-O]_n-CH₃,
- v) -[CH₂-O]_n-C₁-C₃ alkyl,
- w) -NR⁵(CO)-NR⁵,
- x) -CF₃,
- y) - NR⁵(CO)C₁-C₅ alkyl,
- z) -N(R⁵)-SO₂-R⁵,
- a1) -O-C(O)-R⁵,
- b1) -S(O)-R⁵,
- c1) -SR⁵, or
- d1) -SO₂-R⁵;

wherein R¹⁰ is

- a) CO₂R⁵, or
- b) 5-tetrazolyl,

wherein R¹¹ is


- a) H, or
b) methyl;

wherein X is -CO- or -SO₂- or -CO₂-;

wherein n is zero, one, two or three;

or a pharmaceutically acceptable salt thereof.


[0014] The present invention particularly provides the compounds of formula III or IV



wherein G¹, R³, and R⁵ are as defined above, and the chiral center depicted in III is in the (S) configuration.

[0015] The present invention in particular relates to compounds of formula III or IV, wherein G¹ is

- a) -CH(CH₂phenyl)NHCO₂t-Bu,
b) -CH(CH₂phenyl)NHCOC₁₋₃ alkyl-G³,
c) -CH(CH₂phenyl)NHCOC₁₋₃ alkyl-CO₂R⁵, or
d)



wherein R³ is

- a) -C₅-C₆ alkyl, or
b) -C₃-C₆ alkyl-phenyl;

wherein R⁵ is -H;

wherein the configuration of the chiral center(s) is (S).

[0016] The compounds of the present invention are named according to the IUPAC or CAS nomenclature system.

[0017] The carbon atoms content of various hydrocarbon-containing moieties is indicated by a prefix designating the minimum and maximum number of carbon atoms in the moiety, i.e., the prefix C_i-C_j indicates a moiety of the integer "i" to the integer "j" carbon atoms, inclusive. Thus, for example, C₁-C₃ alkyl refers to alkyl of one to three carbon atoms, inclusive, or methyl, ethyl, propyl, and isopropyl, straight and branched forms thereof.

[0018] Also, the carbon atom content of various hydrocarbon-containing moieties of the present invention may be indicated by a subscripted integer representing the number of carbon and hydrogen atoms in the moiety, e.g., "C_nH_{2n}" indicates a moiety of the integer "n" carbon atoms, inclusive, and the integer "2n" hydrogen atoms, inclusive. Thus, for example, "C_nH_{2n}" wherein n is one to three carbon atoms, inclusive, and two to six hydrogen atoms, inclusive, or methyl, ethyl, propyl and isopropyl, and all isomeric, straight and branched forms thereof.

[0019] Examples of alkyl of one to nine carbon atoms, inclusive, are methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, and nonyl, and all isomeric forms thereof and straight and branched forms thereof.

[0020] Examples of alkenyl of one to five carbon atoms, inclusive, are ethenyl, propenyl, butenyl, pentenyl, all isomeric forms thereof, and straight and branched forms thereof.

[0021] By "halo" is meant the typical halogen atoms, such as fluorine, chlorine, bromine, and iodine.

[0022] The present invention encompasses all possible combinations of configurations at each of the possible chiral centers. The preferred configuration for the chiral center depicted in formula I is (S), and the preferred configuration for the chiral center present in R² (d and e) is (S).

[0023] The compounds of formulae I and II of the present invention are prepared as described in the Charts, Preparations and Examples below, or are prepared by methods analogous thereto, which are readily known and available to one of ordinary skill in the art of organic synthesis.

5 CHART A

[0024] Commercially available tyrosine benzyl ester A-1 is acylated with monomethyl succinate under standard amide coupling conditions employing 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) as the coupling reagent (Tet. Lett. 1993, 34:7685) to afford A-2. Alkylation of the phenol is effected with diethylchloromalonate in acetone with potassium carbonate as catalyst, conditions analogous to those previously described for alkylation of phenols (J. Am. Chem. Soc. 1951, 73:872). Standard hydrogenolysis of the benzyl ester A-3 affords A-4, which is then acylated with various amines (R^3NH_2) under the influence of EDC. The target acids A-5 are obtained by saponification.

15 CHART B

[0025] Commercially available Cbz-tyrosine (B-1) is coupled with n-pentylamine under standard EDC conditions, affording amide B-2. Alkylation of the phenol as described in Chart A gives ether B-3, which is then hydrogenolytically deprotected to obtain amine B-4, isolated as the corresponding HCl salt. Acylation of the free amine with various carboxylic acids R^2COOH is accomplished with EDC. Final saponification with dilute lithium hydroxide followed by acidification gives the sparingly soluble malonic acids B-5.

CHART C

[0026] Amine hydrochloride B-4 (from Chart B) is acylated with various isocyanates in the presence of triethylamine in methylene chloride to afford the corresponding urethanes. Saponification of the esters then provides the acids C-1.

CHART D

[0027] Amine B-4 (from Chart B) is converted to the corresponding isocyanate D-1 by reaction with diphosgene and Proton Sponge at 0 C (J. Org. Chem. 1996, 61:3883). Addition of N-benzylglycine ethyl ester followed by saponification then affords the desired urethane triacid D-2.

CHART E

[0028] Commercially available Boc-(L)-tyrosine E-1 is coupled with n-pentylamine, as described for Chart B, to give E-2. The Boc group is removed with HCl in acetic acid, and the resulting amine E-3 is coupled with a mono succinate ester as described for Chart A. The resulting phenol amides E-4 and E-5 are added directly to a dialkyl acetylenedicarboxylate in the presence of triethylamine (Aust. J. Chem. 1995, 48:677). Fumarate ester E-6 is hydrogenated with 10% palladium on carbon to give the saturated triacid E-8. Alternatively, fumarate ester E-7 is saponified to give the unsaturated triacid E-9.

CHART F

[0029] Amine hydrochloride E-3 (from Chart E) is reacted with succinic anhydride in the presence of triethylamine to afford the acid F-1. Sulfation of the phenol is effected with sulfur trioxide/pyridine complex in DMF (Int. J. Pep. Prot. Res. 1990, 35:566) and purification is accomplished with reverse phase HPLC to give F-2.

CHARTG

[0030] Previously described E-2 (from Chart E) is treated with trifluoromethanesulfonic anhydride in the presence of pyridine to afford triflate G-1. Palladium-catalyzed cross-coupling of G-1 with tributyl(vinyl)tin affords G-2. G-2 is then ozonized followed by reduction with dimethyl sulfide to give G-3. This aldehyde is then condensed with dibenzylmalonate in the presence of piperidine acetate to afford G-4. Deprotection of the Boc group with saturated HCl/HOAc affords G-5 which is subsequently reacted with succinic anhydride to afford G-6. Hydrogenation of G-6 with H_2 and 10% Pd/C gives final triacid G-7.

CHART H

[0031] Direct saponification of dibenzylester G-6 (from Chart G) affords the unsaturated triacid H-1.

CHART I

[0032] Alkylation of phenol E-4 (from Chart E) is accomplished by a carbenoid insertion reaction with di-*t*-butyl diazomalonate (Synthesis 1974, 347) catalyzed by rhodium acetate (J. Med. Chem. 1995, 38:4270), affording malonate ether I-1. Removal of the *t*-butyl esters is accomplished with trifluoroacetic acid in methylene chloride, and the benzyl ester is removed by hydrogenolysis, affording the desired triacid I-3.

CHART J

[0033] Amide E-2 (from Chart E) is alkylated on the phenolic hydroxyl with dibenzyl bromomalonate as described for Chart A (potassium carbonate/acetone) to give J-3. The Boc group is removed with HCl in acetic acid, affording the amine hydrochloride J-4. The free amine is added to various cyclic anhydrides in the presence of triethylamine, giving acids J-5. Hydrogenolysis of the benzyl esters then affords the desired triacids J-6.

CHART K

[0034] Chart K describes an alternative synthesis of A-5 (from Chart A) (now K-6 in Chart K) wherein benzyl esters are used as the protecting group for the malonate carboxyls instead of ethyl esters. Tyrosine *t*-butyl ester K-1 is acylated with monobenzyl succinate under the influence of EDC to afford amide K-2. Alkylation with dibenzyl bromomalonate under the conditions described in Chart A affords ether K-3. The *t*-butyl ester is removed with TFA in methylene chloride, giving carboxylic acid K-4, which is coupled with various amines using EDC as the coupling reagent. Final deprotection of K-5 is accomplished by hydrogenolysis to give K-6.

CHART L

[0035] Chart L describes an extension of Chart J wherein amine J-4 (from chart J) is coupled (EDC) with a protected amino acid to afford L-2. The Boc group is removed with HCl in acetic acid to give amine L-3. Addition to succinic anhydride followed by hydrogenolysis of the benzyl esters L-4 then provides the desired tetracids L-5.

CHART M

[0036] Cbz-tyrosine M-1 is coupled (EDC) with norleucine amide to provide M-2. Alkylation of the phenol with diethyl chloromalonate as described for Chart A gives ether M-3. The Cbz group is removed by hydrogenation, and the resulting free amine M-4 is acylated with succinic anhydride. Carboxylic acid M-5 is then saponified to give the target triacid M-6.

CHART N

[0037] Commercially available N-1 is condensed with dibenzylmalonate in the presence of piperidine acetate to afford N-2. N-2 is coupled to previously described J-4 (from Chart J) to afford N-3. Hydrogenation of N-3 leads to final tetraacid N-4.

CHART O

[0038] Direct hydrogenation of benzyl ester L-2 (from Chart L) gives the Boc-protected triacid O-1.

CHART P

[0039] Acylation of amine L-3 (from Chart L) with hexanoyl chloride gives amide P-1. Hydrogenation then removes the benzyl esters, providing triacid P-2.

CHART Q

[0040] Commercially available Q-1 is N-protected as the Boc derivative by reaction with Boc₂O, and the resulting compound is converted to amylamide Q-2 by coupling (EDC) with amylamine. Palladium catalyzed carbonylation with

carbon monoxide and methanol affords methyl ester Q-3. Alkylation of the phenolic oxygen with methylbromoacetate yields ether Q-4, which is N-deblocked with trifluoroacetic acid in methylene chloride and then acylated with succinic anhydride, leading to amide Q-5. Saponification under standard conditions then produced the desired triacid Q-6.

5 CHART R

[0041] Q-4 is deblocked with HCl/dioxane before coupling with 3-phenylpropanoic acid in the presence of EDC and saponified to afford R-4. Alternatively, Q-4 may be deblocked as before, followed by coupling with Boc-L-Phe to afford R-1. R-1 may be saponified directly to R-2, or the Boc group can be removed with HCl/dioxane, and the resulting amine can be coupled with an acid chloride or carboxylic acid to afford R-3 after saponification.

10 CHART S

[0042] S-1 is alkylated with diethylchloromalonate to afford S-2. Removal of the Boc group with HCl/dioxane followed by coupling with Boc-L-p-benzoyl-Phe gives S-3. Removal of the Boc group again followed by addition of succinic anhydride and saponification provides triacid S-4.

CHARTT

[0043] Iodotyrosine Q-2 is converted to nitrile T-1 by the action of zinc cyanide and Pd catalyst. Alkylation with methyl bromoacetate affords ether T-2, which is coupled with Boc-L-Phe after deblocking of the amine group with HCl. The nitrile is converted to the corresponding tetrazole T-4 with TMS-azide and catalytic dibutyltin oxide. Final saponification affords the acid T-5.

25 CHART U

[0044] Q-2 is carbonylated with carbon monoxide and palladium catalyst to afford esters U-1 and Q-3. The phenols are alkylated with methyl or benzyl bromoacetate to afford U-2 and U-3. The Boc group is removed with TFA, followed by coupling with Boc-L-Phe, affording amides U-4 and U-5. Catalytic hydrogenation removes the benzyl esters, providing U-6 and U-7. Coupling of the free carboxylic acids with hydroxylamine generates the hydroxamic acids U-8 and U-9, and the methyl esters are saponified with lithium hydroxide to provide acids U-10 and U-11.

CHART V

[0045] Ester V-1 is reduced with DIBAL to afford aldehyde V-2, which is subsequently converted by a Wittig reaction to olefin V-3. The phenol is alkylated with dibenzyl bromomalonate to afford ether V-4. Deprotection of the amine with TFA, followed by acylation of the free amine with mono-benzylsuccinate affords amide V-5. Saponification of the esters (LiOH) then provides the triacid V-6.

40 CHART W

[0046] Commercially available acid W-1 is amidated with n-pentylamine (EDC), and the resulting amide W-2 is catalytically hydrogenated to aniline W-3. The aniline is bis(alkylated) with methyl bromoacetate to afford W-4. Removal of the Boc group (TFA) followed by acylation of the amine with Boc-L-Phe affords W-5. Final saponification (LiOH) then provides the diacid W-6.

CHART X

[0047] Commercially available meta-iodotyrosine is esterified with benzyl alcohol before coupling with Boc-L-Phe, affording X-2. The iodine is carboxylated with CO under palladium catalysis, providing ester X-3, which is alkylated with methyl bromoacetate. The resulting ether X-4 is hydrogenated to remove the benzyl ester protecting group, and the resulting acid X-5 is reduced with sodium borohydride via the corresponding acyl imidazole to alcohol X-6. Final saponification of the esters affords the diacid X-7.

55 CHART Y

[0048] Commercially available methyl tyrosine Y-1 is protected as the N-Boc derivative under standard conditions before conversion of the carboxylic acid to amide Y-3. This amide is alkylated with dibenzyl bromomalonate to afford

ether Y-4. Boc cleavage with HCl is followed by acylation of the free amine with succinic anhydride, providing acid Y-5. Final saponification with LiOH affords triacid Y-6.

CHART Z

[0049] Meta fluorotyrosine Z-1 is converted to triacid Z-6 exactly as described in Chart Y.

CHART AA

[0050] 4-Hydroxybenzaldehyde AA-1 is alkylated with diethyl chloromalonate to afford ether AA-2. Mono ethyl malonate AA-3 is coupled with amylamine under standard conditions (DEPC) to afford amide AA-4. Hydrolysis of the ester with aq NaOH provides acid AA-5. Coupling of AA-5 with beta-alanine ethyl ester provides malondiamide AA-6, which is condensed with aldehyde AA-2 under Knoevenagel conditions. The resulting methyldene malondiamide AA-7 (a mixture of olefin isomers) is saturated by catalytic hydrogenation, and the ester AA-8 is saponified to triacid AA-9 with aq NaOH.

CHART BB

[0051] Amine B-4 (from Chart B) is acylated with the appropriate protected amino acid under the influence of EDC and triethylamine. The resulting amides BB-1 is directly saponified to give BB-3. Alternatively, where R⁶ is t-butylcarboxy (Boc), the Boc group is removed with HCl/acetic acid, and the resulting free amine acylated with succinic anhydride. Final saponification then affords the triacids BB-2.

CHART CC

[0052] Diol CC-1 (reference given in Example 142) is bis(alkylated) with ethyl bromoacetate, and the resulting bis (ether) CC-2 is hydrogenated to remove the benzyl ester. The carboxylic acid CC-3 is coupled with amylamine under standard conditions (DEPC) before cleavage of the Boc group with TFA. Free amine CC-5 is then coupled with Boc-L-Phe (DEPC), and the resulting amide CC-6 is saponified to the diacid CC-7.

CHART DD

[0053] Diester Q-4 (Chart Q) is reacted with TFA to cleave the Boc group and the free amine DD-1 is coupled (DEPC) with the appropriate amino acid (see Example 143) to afford amide DD-2. Saponification provides diacid DD-3.

CHART EE

[0054] Q-2 (Chart Q) is carbonylated with CO under palladium catalysis to afford ester EE-1. The phenol is alkylated with ethyl bromofluoroacetate/potassium carbonate to afford ether EE-2. Boc deprotection and amide coupling of the free amine with Boc-L-Phe under standard conditions affords diester EE-3, which is saponified to provide the diacid EE-4.

CHART FF

[0055] Acid X-5 (Chart X) is coupled with 4-phenylbutylamine under standard amide coupling conditions to provide FF-1. Saponification of the esters affords diacid FF-2.

[0056] Preferred methods of preparation are depicted in Charts A, B, BB, Q and R.

[0057] The present invention provides for compounds of formulae I and II or pharmacologically acceptable salts and/or hydrates thereof. Pharmacologically acceptable salts refers to those salts which would be readily apparent to a manufacturing pharmaceutical chemist to be equivalent to the parent compound in properties such as formulation, stability, patient acceptance and bioavailability. Examples of salts of the compounds of formula I include lithium, sodium and potassium.

[0058] Where R⁵ is other than H, the compounds would not be expected to have intrinsic activity, but would be expected to possess activity *in vivo* following hydrolysis by non-specific esterases to the corresponding carboxylic acids.

[0059] The compounds of the present invention are useful for treating patients with noninsulin-dependent diabetes mellitus (NIDDM) and related diseases. For this indication, these compounds may be administered by oral, intranasal, transdermal, subcutaneous and parenteral (including intramuscular and intravenous) routes in doses of 0.1 mg to 1000 mg/kg of body weight per day.

[0060] Those skilled in the art would know how to formulate the compounds of this invention into appropriate pharmaceutical dosage forms. Examples of the dosage forms include oral formulations, such as tablets or capsules, or parenteral formulations, such as sterile solutions.

[0061] When the compounds in this invention are administered orally, an effective amount is from about 0.1 mg to 100 mg per kg of body weight per day. Either solid or fluid dosage forms can be prepared for oral administration. Solid compositions, such as compressed tablets, are prepared by mixing the compounds of this invention with conventional ingredients such as talc, magnesium stearate, dicalcium phosphate, magnesium aluminum silicate, calcium sulfate, starch, lactose, acacia, methyl cellulose, or functionally similar pharmaceutical diluents and carriers. Capsules are prepared by mixing the compounds of this invention with an inert pharmaceutical diluent and placing the mixture into an appropriately sized hard gelatin capsule. Soft gelatin capsules are prepared by machine encapsulation of a slurry of the compounds of this invention with an acceptable inert oil such as vegetable oil or light liquid petrolatum.

[0062] Syrups are prepared by dissolving the compounds of this invention in an aqueous vehicle and adding sugar, aromatic flavoring agents and preservatives. Elixirs are prepared using a hydroalcoholic vehicle such as ethanol, suitable sweeteners such as sugar or saccharin and an aromatic flavoring agent. Suspensions are prepared with an aqueous vehicle and a suspending agent such as acacia, tragacanth, or methyl cellulose.

[0063] When the compounds of this invention are administered parenterally, they can be given by injection or by intravenous infusion. An effective amount is from about 0.1 mg to 100 mg per kg of body weight per day. Parenteral solutions are prepared by dissolving the compounds of this invention in aqueous vehicle and filter sterilizing the solution before placing in a suitable sealable vial or ampule. Parenteral suspensions are prepared in substantially the same way except a sterile suspension vehicle is used and the compounds of this invention are sterilized with ethylene oxide or suitable gas before it is suspended in the vehicle.

[0064] The exact route of administration, dose, or frequency of administration would be readily determined by those skilled in the art and is dependant on the age, weight, general physical condition, or other clinical symptoms specific to the patient to be treated.

[0065] The utility of representative compounds of the present invention has been demonstrated in the biological assays described below:

[0066] **PTP1 Assays:** A construct, which consisted of a C-terminal truncation of rat PTP1 (amino acid residues 1-322) (cloned from a rat brain library) with an N-terminal glutathione S-transferase (GST) tag and an adjacent thrombin cleavage site, was inserted into vector plasmid pGEX-2T and transformed into E.coli strain TG-1 under the control of a lac promoter (K. L. Guan and J. E. Dixon (1991) Eukaryotic proteins expressed in Escherichia coli: an improved thrombin cleavage and purification procedure of fusion proteins with glutathione S-transferase. *Analyt. Biochem.* 192: 262-267). The GST-fusion protein was purified on a glutathione agarose affinity column, the GST tag was cleaved with thrombin, and the active enzyme was recovered for use in an assay to identify PTP inhibitors.

[0067] The equivalent construct of human PTP1B (amino acid residues 1-321) (cloned from a human placental library), without the GST tag and thrombin cleavage site, was inserted into a pMB replicon and transformed into E. coli BL21(DE3), a strain containing a chromosomal copy of the gene for T7 RNA polymerase under control of a lacUV5 promoter. Expression of PTP1B was induced with isopropyl thiogalactose and the soluble protein was purified by ion exchange, hydrophobic interaction and gel exclusion chromatography for use in the assay to identify PTP inhibitors.

[0068] PTP1 activity is measured using either p-nitrophenol phosphate (pNPP) or a triphosphopeptide (that matches residues 1142 through 1153 of the β -subunit and the insulin receptor) as substrate in a 96-well microtiter plate format. An assay pH of 7.2 is used for standard assays (measured A_{405} =9800 at pH 7.2).

[0069] Human PTP1B, which is highly homologous to rat PTP1, was assayed exactly as described above for PTP1. The PTP1 inhibitors described here also inhibit PTP1B with similar or identical potencies.

[0070] Standard assays are conducted at room temperature in a total volume of 0.2 ml that contains Hepes buffer (50 mM, pH 7.2), NaCl (50 mM), EDTA (1 mM), DTT (1 mM), bovine serum albumin (1 mg/ml), pNPP (1 mM) and PTP1 (35 ng/ml). Compounds (2 μ l of 10 mM solutions) are pipetted into wells of microtiter plates followed by 198 μ l of premixed reaction mix (with PTP1 and pNPP added immediately before use). The rate of change in A_{405} is recorded for 60 min. Two wells on each plate contain DMSO controls and two wells contain sodium orthovanadate (1 mM) which inhibits PTP1-catalyzed hydrolysis of pNPP completely. Data are expressed as percent inhibition relative to the average of the DMSO controls measured on the same microtiter plate.

[0071] When triphosphopeptide ¹¹⁴²⁻¹¹⁵³ is used as substrate, the rate of release of inorganic phosphate is measured using a Malachite Green/phosphomolybdate reaction (A.A. Baykov, O.A. Evtushenko, and S.M. Avaeva (1988) A Malachite Green procedure for orthophosphate determination and its use in alkaline phosphatase-based enzyme immunoassay. *Anal. Biochem.* 171: 266-270.) in a microtiter plate format. Standard assays are conducted at room temperature in a total volume of 50 μ l that contains Hepes buffer (50 mM, pH 7.2), NaCl (50 mM), EDTA (1mM), DTT (1mM), bovine serum albumin (1mg/ml), triphosphopeptide ¹¹⁴²⁻¹¹⁵³ (200 μ M) and PTP1 (87 ng/ml). Reactions are terminated with the addition of 0.15 ml of Malachite Green/ ammonium molybdate reagent [10ml Malachite Green (0.44 g in 6N H₂SO₄), 2.5 ml ammonium molybdate (7.5% w/v), 0.2 ml Tween 20 (11% w/v)] that is diluted with 8 parts of water

immediately before use, and after 1 h absorbance at 650 nm is measured. The phosphate assay is calibrated using either KH_2PO_4 or pNPP (after ashing with $\text{Mg}(\text{NO}_3)_2$) which gives essentially identical standard curves. The phosphate assay is useful in the range of 1 to 10 nmol Pi.

[0072] The % inhibition of pNPP-hydrolysis by compounds of the present invention are listed in Table 1 and 2 below.

[0073] The following compounds of the present invention are preferred:

(S)-2-(Carboxymethoxy)-5-[2-[(3-carboxy-1-oxopropyl)amino]-3-oxo-3-(pentylamino)propyl]benzoic acid;
 5-[2(S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]-2-(carboxymethoxy)benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-2-[(dibenzylamino)carbonyl]amino]-3-oxo-3-(pentylamino)propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(3-phenylpropanoyl)amino]propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-2-[(2S)-2-[(3-carboxypropanoyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoic acid;
 5-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]-2-(carboxymethoxy)benzoic acid;
 5-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-(4-hydroxyphenyl)propanoyl]amino]-3-oxo-3-(pentylamino)propyl]-2-(carboxymethoxy)benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phenyl-2-[[2-(4H-1,2,4-triazol-3-ylsulfanyl)acetyl]amino]propanoyl]amino]propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phenyl-2-[[2-(5-sulfanyl-1H-1,2,3,4-tetraazol-1-yl)acetyl]amino]propanoyl]amino]propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phenyl-2-[[2-(1H-1,2,3-triazol-5-ylsulfanyl)acetyl]amino]propanoyl]amino]propyl]benzoic acid;
 2-[4-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]-2-(2H-1,2,3,4-tetraazol-5-yl)phenoxy]acetic acid;
 5-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]-2-(2-methoxy-2-oxoethoxy)benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-2-[(2S)-2-[[2-(1H-indol-3-yl)acetyl]amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phenyl-2-[(2-phenylacetyl)amino]propanoyl]amino]propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phenyl-2-[(4-phenylbutanoyl)amino]propanoyl]amino]propyl]benzoic acid;
 5-[(2S)-2-[(2S)-2-(acetyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]-2-(carboxymethoxy)benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-2-[(2S)-2-[(3-methoxypropanoyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-2-[(2S)-2-[(4-hydroxybutanoyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoic acid;
 5-[(2S,5S)-5-benzyl-13,13-dimethyl-4,7,11-trioxo-2-[(pentylamino)carbonyl]-12-oxa-3,6,10-triazatetradec-1-yl]-2-(carboxymethoxy)benzoic acid;
 5-[(2S,5S)-5-benzyl-4,7,11,11-tetraoxo-2-[(pentylamino)carbonyl]-11 λ 6-thia-3,6,10-triazadodec-1-yl]-2-(carboxymethoxy)benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-2-[(2S)-2-[[2-(3-hydroxyphenyl)acetyl]amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-2-[(2S)-2-[[2-(4-hydroxyphenyl)acetyl]amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-2-[(2S)-2-[[2-(4-methylphenyl)acetyl]amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phenyl-2-[[2-(4-(trifluoromethyl)phenyl)acetyl]amino]propanoyl]amino]propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-2-[(2S)-2-[[2-(4-methoxyphenyl)acetyl]amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoic acid;
 5-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]propanoyl]amino]-3-oxo-3-(pentylamino)propyl]-2-(carboxymethoxy)benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phenyl-2-[(3-phenylpropanoyl)amino]propanoyl]amino]propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-3-oxo-2-[(2R)-2-(2-oxo-1-pyrrolidinyl)-3-phenylpropanoyl]amino]-3-(pentylamino)

propyl]benzoic acid; and
 5-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-[(4-phenylbutyl)amino]propyl]-2-(carboxymethoxy)benzoic acid.

[0074] The invention also relates to the use of a compound as disclosed above in the manufacture of a medicament useful for the treatment and/or prevention of Non-Insulin Dependent Diabetes Mellitus.

DESCRIPTION OF PREFERRED EMBODIMENTS

EXAMPLE 1 (COMPARISON): (S)-4-Oxo-4-[[2-oxo-2-(pentylamino)-1-[[4-(sulfooxy)phenyl]methyl]ethyl]amino]butanoic acid (Formula F-2, Chart F)

[0075] PREPARATION OF E-2 (Chart E): To a 0 °C mixture of Boc-L-tyrosine (2.04 g) and amylamine (0.93 mL) in methylene chloride (30 mL) is added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (EDC) (1.53 g). The white mixture is stirred at 0 °C for 5 min and at room temp for 23 hrs. The resulting solution is diluted with methylene chloride (30 mL) and washed successively with 0.5 M HCl (40 mL), water (20 mL) and sat aq sodium bicarbonate (25 mL). The organic phase is dried over magnesium sulfate and concentrated to a foam (1.84 g), sufficiently pure to carry into the next step. An analytical sample is obtained by flash chromatography (1/1 ethyl acetate/hexane) as a glass.

[0076] Physical characteristics are as follows: ¹H NMR (CDCl₃) δ 7.01, 6.74, 5.90, 5.17, 4.21, 3.13, 2.93, 1.41, 1.1-1.4, 0.85; IR (mull) 3341, 3318, 3004, 1684, 1651, 1595, 1517, 1393, 1293, 1267, 1248, 1170, 1048, 828; MS (EI) m/z 350 (M+), 234, 233, 188, 180, 177, 147, 143, 136, 107, 57 cm⁻¹; HRMS (EI) found 350.2209. Anal. Found: C, 64.27; H, 8.54; N, 7.82.

[0077] PREPARATION OF E-3 (Chart E): To a solution of E-2 (5.28 g) in dioxane (40 mL) chilled in an ice bath is added a freshly prepared solution of HCl in dioxane (about 3 M, 25 mL). The solution is then stirred at room temp for 1.5 hrs when a TLC indicates the reaction is done. The solution is diluted rapidly with ether (350 mL) until no further precipitation is evident. The mixture is stirred vigorously until all insoluble material is adhering to the sides of the flask. After decanting the supernatant, the crude material is taken up in more ether (200 mL) and sonicated until a fine solid (required about 1 hr). Filtration gives a hygroscopic white powder (4.32 g).

[0078] Physical characteristics are as follows: ¹H NMR (DMSO) δ 9.3, 8.40, 8.25, 6.98, 6.68, 3.08, 2.9, 1.0-1.35, and 0.83.

[0079] PREPARATION OF F-1 (Chart F): Triethylamine (307 uL) is added to a 0 °C mixture of E-3 (287 mg) in methylene chloride (4 mL), causing rapid dissolution. To this solution is added succinic anhydride (100 mg), and the reaction is stirred at room temp for 25 h. The reaction is then diluted with ethyl acetate (20 mL) and washed successively with 0.5 M HCl (10 mL) and brine (10 mL). The organic phase is dried over magnesium sulfate and concentrated to a viscous oil (350 mg) that solidified on standing and is analytically pure.

[0080] Physical characteristics are as follows: ¹H NMR (DMSO) δ 9.1, 7.98, 7.74, 6.96, 6.60, 4.30, 2.9-3.1, 2.8, 2.60, 2.3, 1.1-1.4, 0.83; ¹³C NMR (CDCl₃) 175.1, 173.1, 172.2, 155.9, 129.8, 127.8, 114.8, 55.3, 39.2, 39.0, 36.7, 30.0, 28.8, 28.7, 28.5, 22.0, 13.0; IR (mull) 3296, 3102, 2728, 1715, 1642, 1615, 1596, 1548, 1516, 1401, 1239, 1173, 1117, 832, 722; MS (EI) m/z 350 (M+), 233, 177, 162, 147, 144, 143, 136, 107, 91, 55 cm⁻¹; MS (FAB) m/z 351 (M+H), 352, 351, 350, 236, 233, 136, 121, 107, 88, 43. HRMS (FAB) found 351.1928. Anal. Found: C, 60.34; H, 7.48; N, 7.72.

[0081] PREPARATION OF F-2 (Chart F): A solution of F-1 (100 mg) and pyridine sulfur trioxide complex (500 mg) in DMF:pyridine (1:1, 10 mL) is stirred under nitrogen at room temp for 20 hrs. NMR analysis of an aliquot indicates complete conversion to product. Solvent is removed under vacuum, leaving a solid that is purified by reverse phase HPLC.

[0082] Physical characteristics are as follows: ¹H NMR (DMSO) δ 8.03, 7.79, 7.06, 7.01, 4.33, 2.99, 2.87, 2.68, 2.3, 1.1-1.3, 0.83; ES MS m/z (negative ion) 429, 214.

EXAMPLE 2 (COMPARISON): (S)-[4-[2-[[1,4-Dioxo-4-(phenylmethoxy)butyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]propanedioic acid (Formula I-2, Chart I)

[0083] PREPARATION OF E-4 (Chart E): EDC (368 mg) is added to a 0 °C solution of E-3 (500 mg), (mono)benzyl succinate (368 mg) and triethylamine (270 uL) in methylene chloride (7 mL). The reaction is stirred at room temp for 24 h. After dilution with ethyl acetate (40 mL), the mixture is washed successively with 0.5 M HCl, water and sat aq sodium bicarbonate (20 mL each). The organic phase is dried over magnesium sulfate and concentrated to a white amorphous solid (694 mg) that is analytically pure.

[0084] Physical characteristics are as follows: [α]_D²⁵ -5.4° (c 0.017, methanol); ¹H NMR (CDCl₃) δ 7.33, 7.05, 6.74, 6.21, 5.90, 5.68, 5.13, 5.09, 4.54, 3.12, 3.06, 2.90, 2.4-2.85, 1.1-1.4, 0.86; IR (mull) 3379, 3285, 1706, 1640, 1617, 1552, 1517, 1339, 1271, 1218, 1196, 1181, 1174, 749, 694; MS (EI) m/z 440 (M+), 233, 226, 208, 147, 136, 108, 107,

91, 79, 77. Anal. Found: C, 67.89; H, 7.23; N, 6.38.

[0085] PREPARATION OF I-1 (Chart I): A solution of di-tert-butyl diazomalonate (396 mg) in benzene (2 mL) is added over 6 h to an 80 °C mixture of E-4 (342 mg) and rhodium (II) acetate (7 mg) in benzene (33 mL) via syringe pump, during which time most of the starting material goes into solution. The blue-green solution is stirred at that temp for an additional hour and then overnight at room temp. The reaction is filtered through a medium frit and then concentrated in vacuo. Flash chromatography (80 g silica, 70% ethyl acetate/hexane) provides the title material (232 mg) as a foam (R_f = 0.45).

[0086] Physical characteristics are as follows: ¹H NMR (CDCl₃) δ 7.33, 7.12, 6.89, 6.10, 5.81, 5.12, 5.08, 4.94, 4.52, 3.1, 2.6-2.95, 2.45, 1.49, 1.1-1.4, 0.86; MS (FAB) *m/z* 655 (M+H), 655, 599, 447, 92, 91, 88, 86, 57, 41, 29. HRMS (FAB) found 655.3588.

[0087] PREPARATION OF I-2 (Chart I): Trifluoroacetic acid (5 mL) is added to a solution of I-1 (225 mg) in methylene chloride (5 mL) with ice bath chilling. The solution is stirred at room temp for 2 h. Concentration *in vacuo* affords the title compound as a light amber foam, sufficiently pure to carry into the next step.

[0088] Physical characteristics are as follows: ¹H NMR (CD₃OD) δ 7.33, 7.15, 6.89, 5.23, 5.10, 4.48, 3.05, 2.80, 2.35-2.7, 1.1-1.5, 0.88; MS (FAB) *m/z* 543 (M+H), 544, 543, 542, 441, 92, 91, 88, 86, 43. HRMS (FAB) found 543.2332.

EXAMPLE 3 (COMPARISON): (S)-[4-[2-[(3-Carboxy-1-oxopropyl)amino]-3-oxo-3-(pentylamino)propyl]phenoxy]propanedioic acid (Formula 1-3, Chart I)

[0089] Crude I-2 (approx. 0.29 mmol) is dissolved in ethyl acetate and concentrated in vacuo three times to get rid of traces of trifluoroacetic acid. The residue is then dissolved in methanol (10 mL) and subjected to three cycles of evacuation and nitrogen purge at 0 °C before the introduction of 10% Pd/C (20 mg). The mixture is then hydrogenated at 1 atm for 1.5 h. The mixture is filtered through Celite and concentrated in vacuo. The crude glass is taken up in methylene chloride (40 mL) and sonicated until it is all suspended. Filtration gives a brittle white amorphous solid (116 mg) that is analytically pure (m.p. 117-120 °C, dec).

[0090] Physical characteristics are as follows: [α]_D²⁵ = -0.7° (c 0.0058, methanol); ¹H NMR (CD₃OD) δ 8.13, 7.81, 7.17, 6.90, 5.23, 4.48, 3.1, 2.81, 2.3-2.6, 1.2-1.5, 0.89; MS (FAB) *m/z* 453 (M+H), 453, 238, 194, 136, 133, 101, 88, 86, 55, 43. HRMS (FAB) found 453.1859. Anal. Found: C, 54.25; H, 6.31; N, 5.97.

EXAMPLE 4 (COMPARISON): (S)-[4-[2-[[[(Carboxymethyl)amino]carbonyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]propanedioic acid (Formula C-1 (R² = CH₂CO₂H), Chart C)

[0091] To a solution of amine hydrochloride B-4 (50 mg) and triethylamine (16 μL) in methylene chloride (1 mL) is added ethyl isocyanatoacetate (13 μL) neat. TLC analysis shows reaction is nearly done after 5 min. After stirring for a total of 1 hr at room temp., the reaction is diluted with ethyl acetate (5 mL) and washed successively with 1 M HCl, water, and brine (2 mL each). The organic layer is dried over magnesium sulfate and concentrated in vacuo to a colorless oil (58 mg).

[0092] Physical characteristics are as follows: ¹H NMR (CDCl₃) δ 7.11, 6.82, 6.57, 6.48, 6.11, 5.10, 4.49, 4.28, 4.15, 3.94, 3.82, 2.8-3.2, 1.1-1.4, 0.83; MS (ES⁺): 537.9 (M+H), 559.8 (M+Na).

[0093] To a solution of the crude triester from above in 3:1:1 THF:MeOH:water (1 mL) at room temp is added lithium hydroxide monohydrate (21 mg). The solution is stirred for 3 hrs. The solvent is evaporated in vacuo, and the residue is acidified with 1M HCl (1 mL). To the slightly cloudy mixture is added brine (2 mL), resulting in a copious ppt. The mixture is extracted with ethyl acetate, and the extracts are dried over magnesium sulfate. Concentration gives a glass (50 mg). This material is sonicated with methylene chloride (20 mL) for 30 min, and the resulting fine white solid is collected by filtration, giving the title product as an amorphous powder (32 mg).

[0094] Physical characteristics are as follows: ¹H NMR (DMSO) δ 7.84, 7.06, 6.79, 6.33, 6.29, 5.25, 4.25, 3.65, 2.85-3.05, 2.78, 2.63, 1.1-1.4, 0.82; MS (ES⁻): 451.7; HRMS (FAB) found 454.1820. Anal. Found: C, 51.50; H, 5.94; N, 8.89.

EXAMPLE 5 (COMPARISON): (S)-[4-[2-[[[(5-Carboxypentyl)amino]carbonyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]propanedioic acid (Formula C-1, Chart C)

[0095] By the two-step procedure described for Example 4, the title product is obtained as a foam (44 mg). Saponification is effected in 3:1 THF:MeOH (2 mL) with added 2.5 M aq LiOH (0.3 mL). Sonication does not produce a filtratable solid, so the methylene chloride is decanted, and the residue is dried under vacuum.

[0096] Physical characteristics are as follows: ¹H NMR (DMSO) δ 7.82, 7.04, 6.79, 6.00, 5.90, 5.25, 4.25, 2.8-3.0, 2.76, 2.65, 2.16, 1.45, 1.1-1.35, 0.82; MS (ES⁻): 508.1; HRMS (FAB) found 510.2446. Anal. Found: C, 54.63; H, 6.93; N, 8.02.

EXAMPLE 6 (COMPARISON): (S)-[4-[2-[[[(4-Carboxyphenyl)amino]carbonyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]propanedioic acid (Formula C-1, Chart C)

[0097] By the procedure described for Example 5 is obtained the title product as a glass that slowly solidifies (57 mg).

[0098] Physical characteristics are as follows: ¹H NMR (DMSO) δ 9.06, 8.05, 7.77, 7.42, 7.07, 6.81, 6.42, 5.26, 4.38, 3.01, 2.7-2.9, 1.1-1.4, 0.82; MS (ES⁻): 514.0; HRMS (FAB): Found = 516.1979.

EXAMPLE 7 (COMPARISON): (S)-[4-[2-[[[(2-Carboxyethyl)amino]carbonyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]propanedioic acid (Formula C-1, Chart C)

[0099] By the procedure described for Example 5 is obtained the title compound (43 mg) as a glass. Sonication in methylene chloride was not performed.

[0100] Physical characteristics are as follows: ¹H NMR (DMSO) δ 7.83, 7.04, 6.79, 6.11, 5.25, 4.22, 3.13, 2.96, 2.76, 2.61, 2.27, 1.1-1.4, 0.82; MS (ES⁻): 465.8; HRMS (FAB) found 468.1982.

EXAMPLE 8 (COMPARISON): (S)-5-[[[1-[4-(Dicarboxymethoxy)phenyl]methyl]-2-oxo-2-(pentylamino)ethyl]amino]carbonyl]amino]-1,3-benzene-dicarboxylic acid (Formula C-1, Chart C)

[0101] By the procedure described for Example 5 is obtained the title compound as a glass (61 mg).

[0102] Physical characteristics are as follows: ¹H NMR (DMSO) δ 8.15, 8.04, 7.99, 7.08, 6.82, 6.32, 5.26, 4.37, 2.7-3.1, 1.1-1.4, 0.82; MS (ES⁻): 557.8; HRMS (FAB) 560.1880. Anal. Found: C, 54.21; H, 5.28; N, 6.89.

EXAMPLE 9 (COMPARISON): (S)-N-[[[1-[4-(Dicarboxymethoxy)phenyl]methyl]-2-oxo-2-(pentylamino)ethyl]amino]carbonyl]-L-glutamic acid (Formula C-1, Chart C)

[0103] (S)-(-)-2-isocyanatoglutaric acid, diethyl ester (24 uL) is added to a 0°C mixture of B-4 (50 mg) and triethylamine (16 uL) in THF (1 mL). After 1 hr, 2.5 M aq LiOH (0.4 mL) is added directly to the reaction mixture, and the two-phase mixture is stirred vigorously for 1.5 hrs. The mixture is diluted with 1 M HCl (2 mL), saturated with solid NaCl, and extracted with ethyl acetate (3 x 2 mL). Drying of the extracts over magnesium sulfate and concentration leaves a glass (68 mg), which is sonicated with methylene chloride (20 mL) for 1 h. Filtration and drying in vacuo leaves the title compound as a white powder (52 mg).

[0104] Physical characteristics are as follows: ¹H NMR (DMSO) δ 7.84, 7.05, 6.79, 6.44, 6.14, 5.25, 4.22, 4.05, 2.95, 2.80, 2.68, 2.21, 1.85, 1.65, 1.1-1.4, 0.82; MS (ES⁻): 523.9. IR (mull) 3358, 2670, 2605, 1727, 1625, 1564, 1512, 1416, 1341, 1230, 1185, 1114, 855, 833, 805, cm⁻¹. HRMS (FAB) found 526.2051. Anal. Found: C, 51.54; H, 6.06; N, 7.85.

EXAMPLE 10 (COMPARISON): (S)-[4-[2-[[[(Carboxymethyl)(phenylmethyl)amino]carbonyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]propanedioic acid (Formula D-2, Chart D)

[0105] To a 0°C solution of B-4 (50 mg), and Proton Sponge (71 mg) in methylene chloride (1 mL) is added diphosgene (8 uL). The solution is stirred at 0°C for 45 min before dilution with more methylene chloride (2 mL) and washing successively with 1 M HCl, water, and sat aq sodium bicarbonate (2 mL each). The extracts are dried over magnesium sulfate and concentrated in vacuo, leaving D-1 as an oil.

[0106] Physical characteristics are as follows: ¹H NMR (CDCl₃) δ 7.17, 6.94, 6.15, 5.18, 4.30, 3.2-3.32, 2.92, 1.5, 1.3, 0.90.

[0107] To a solution of D-1 in THF (1 mL) is added N-benzylglycine ethyl ester (21 uL). The solution is stirred at room temp for 1.5 h before the addition of 2.5 M aq LiOH (0.3 mL). The resulting mixture is stirred vigorously at room temp for 1.5 h. The mixture is diluted with 1M HCl (2 mL), saturated with brine, and extracted with ethyl acetate. The extracts are dried over magnesium sulfate and concentrated in vacuo, leaving a glass (65 mg). Sonication with methylene chloride (20 mL) for 30 min followed by filtration afforded the title compound as a white solid (41 mg).

[0108] Physical characteristics are as follows: ¹H NMR (DMSO) δ 7.69, 7.25, 7.10, 6.77, 6.48, 5.25, 4.40, 4.27, 3.81, 2.7-3.1, 1.1-1.4, 0.84; MS (ES⁻): 541.9. HRMS (FAB) found 544.2293. Anal. Found: C, 58.10; H, 6.08; N, 7.56.

EXAMPLE 11 (COMPARISON): (S)-[4-[2-[[[(Carboxymethyl)[4-(phenylmethoxy)phenyl]methyl]amino]carbonyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]propanedioic acid (Refer to Chart D)

[0109] 4-(4-benzyloxy)benzylglycine methyl ester is prepared from 4-(benzyloxy)-benzaldehyde and glycine methyl ester by the method of Zydowsky et al (J. Org. Chem. 1988, 53:5607). Using the method described for Example 10, this material affords the title compound as a white powder (54 mg). Sonication with methylene chloride does not provide

the product as a filterable solid; it is isolated by decantation and drying of the insoluble residue.

[0110] Physical characteristics are as follows: ^1H NMR (DMSO) δ 7.69, 7.25-7.5, 7.10, 7.04, 6.91, 6.79, 6.45, 5.27, 5.06, 4.32, 4.25, 3.77, 3.0, 2.85, 2.75, 1.1-1.4, 0.84; MS (ES-): 647.9. IR (mull) 3346, 3063, 3032, 1732, 1612, 1587, 1538, 1511, 1422, 1341, 1297, 1230, 1177, 1113, 697, cm^{-1} . Anal. Found: C, 61.89; H, 6.05; N, 6.30.

EXAMPLES 12 - 24 (COMPARISON): (General Synthesis of Formula J-6, Chart J)

[0111] PREPARATION OF J-3: To a stirring solution of E-2 (2.0 g) in acetone (50 ml) is added K_2CO_3 (1.57 g) at ambient temperature. To the resulting heterogeneous mixture is added dibenzyl bromomalonate (2.89 g) and the mixture stirred at ambient temperature overnight. The resulting amber suspension is diluted with H_2O (100 ml) and extracted with EtOAc (2 x 100 ml). The combined organic layers are dried over MgSO_4 and solvent removed in vacuo. The residue is purified via SiO_2 flash chromatography (eluant 2: 1 EtOAc/hexane) to afford 1.26 g title compound as a white solid.

[0112] Physical characteristics are as follows: ^1H NMR (CDCl_3) δ 0.84, 1.16-1.38, 1.39, 2.96, 3.13, 4.21, 5.20, 5.08, 5.23, 5.85, 6.83, 7.07, 7.25; IR (mull) 3346, 3326, 1748, 1683, 1657, 1540, 1522, 1510, 1313, 1297, 1238, 1221, 1188, 1173, 698, cm^{-1} . MS (FAB) m/z 633 (MH^+), 633, 578, 577, 533, 515, 92, 91, 88, 86, 57. Anal. Found: C, 68.06; H, 6.91; N, 4.33.

[0113] PREPARATION OF J-4: To a stirring solution of J-3 (2.85 g) in HOAc (25 ml) at ambient temperature, is added 1.5N HCl/ HOAc (20 ml) and the resulting solution is stirred at ambient temperature for 2 h). The solvent is evaporated to 30 ml and triturated with Et_2O (400 ml). The resulting turbid suspension is stirred at ambient temperature for 30 min, sonicated and filtered to afford 2.50 g title compound as a white solid.

[0114] Physical characteristics are as follows: ^1H NMR (DMSO) δ 0.80, 1.18, 2.93, 3.04, 3.87, 5.20, 5.83, 6.91, 7.13, 7.32, 8.39; IR (liq.) 3035, 2957, 2932, 2872, 2861, 1763, 1748, 1661, 1511, 1500, 1456, 1224, 1185, 1167, 697, cm^{-1} . MS (FAB) m/z 533 (MH^+), 1067, 1066, 535, 534, 533, 418, 143, 92, 91, 88. Anal. Found: C, 64.74; H, 6.54; N, 4.88.

[0115] GENERAL PREPARATION OF J-5: To a stirring solution of J-4 (0.20 g) in CH_2Cl_2 (10 ml) is added triethylamine (0.078 g) at 0 $^\circ\text{C}$. The requisite cyclic anhydride (0.35 mmol) is added in one portion and the resulting solution allowed to stir for 16h while warming to ambient temperature. The solution is diluted with CH_2Cl_2 (50 ml) and washed with 10% HCl/ H_2O (2 x 50 ml). The combined organic phases are dried over MgSO_4 and solvent removed in vacuo to afford material suitable for subsequent transformations.

[0116] GENERAL PREPARATION OF J-6: To a solution of the requisite dibenzylester J-5 in MeOH (\sim .02 M) at ambient temperature is added 10% Pd/C (10 weight %) and the resulting mixture hydrogenated at atmospheric pressure for 3 h. The mixture is filtered through Celite and solvent removed to afford analytically pure material.

[0117] The following examples (12 - 24) are prepared by the general synthesis of J-6 outlined above, using the appropriate commercially available anhydrides (Chart J).

EXAMPLE 12 (COMPARISON) : (S)-[4-[2-[(4-Carboxy-1-oxobutyl)amino]-3-oxo-3-(pentylamino) propyl]phenoxy]propanedioic acid

[0118] 0.115 g as an amorphous white solid. ^1H NMR (DMSO) δ 0.84, 1.15-1.36, 1.61, 2.06, 2.68, 2.84, 3.00, 4.38, 5.25, 6.80, 7.13, 7.86, 8.00; IR (mull) 3305, 3070, 2729, 2669, 2599, 1730, 1626, 1551, 1512, 1341, 1231, 1185, 1112, 855, 831, cm^{-1} . MS (FAB) m/z 467 (MH^+), 481, 468, 467, 423, 238, 194, 91, 88, 59, 43. Anal. Found: C, 54.90; H, 6.52; N, 5.47.

EXAMPLE 13 (COMPARISON) : [1R-[1(S*),2]]-[4-[2-[(2-Carboxycyclohexyl)carbonyl]amino]-3-oxo-3-(pentylamino) propyl]phenoxy]propanedioic acid

[0119] 0.068 g as an amorphous white solid. ^1H NMR (DMSO) δ 0.84, 1.12-1.50, 1.80, 2.4-3.15, 3.35, 5.22, 6.65, 6.78, 7.0, 7.08, 7.22, 7.62, 7.75; IR (mull) 3336, 2731, 2674, 2595, 1727, 1630, 1537, 1512, 1338, 1296, 1229, 1184, 1130, 1103, 854, cm^{-1} . MS (FAB) m/z 507 (MH^+), 508, 507, 238, 194, 136, 102, 91, 88, 81, 43. Anal. Found: C, 57.50; H, 6.67; N, 5.33.

EXAMPLE 14 (COMPARISON) : (S)-[4-[2-[(Carboxymethoxy)acetyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]propanedioic acid

[0120] 0.096 g as an amorphous solid. ^1H NMR (DMSO) δ 0.83, 1.20-1.32, 2.76-3.01, 3.89, 4.03, 4.45, 5.23, 6.78, 7.08, 7.77, 7.95; IR (mull) 3340, 2730, 2670, 2596, 1745, 1632, 1543, 1512, 1346, 1230, 1184, 1144, 833, 721, 668, cm^{-1} . MS (FAB) m/z 469 (MH^+), 497, 483, 469, 107, 88, 86, 75, 43, 39, 23. Anal. Found: C, 52.07; H, 5.98; N, 5.67.

EXAMPLE 15 (COMPARISON) : (S)-[4-[2-[[[1-(Carboxymethyl)cyclopentyl]acetyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]propanedioic acid

[0121] 0.108 g as an amorphous solid. ¹H NMR (DMSO) δ 0.82, 1.20-1.47, 2.18-2.35, 2.60-2.92, 2.99, 4.40, 5.22, 7.79, 7.11, 7.83, 8.00; IR (mull) 3294, 3069, 2728, 2668, 2604, 1732, 1629, 1613, 1554, 1512, 1331, 1280, 1229, 1182, 1110 cm⁻¹. MS (FAB) *m/z* 462 (MH⁺), 535, 521, 143, 136, 109, 88, 81, 67, 43, 39. Anal. Found: C, 58.00; H, 6.87; N, 5.06.

EXAMPLE 16 (COMPARISON) : (S)-[4-[2-[(Carboxyacetyl)amino]-3-oxo-3-(pentylamino)propyl]phenoxy]propanedioic acid

[0122] 0.101 g as an amorphous solid. ¹H NMR (DMSO) δ 0.82, 1.19-1.31, 2.69, 2.85-3.21, 4.37, 5.23, 6.78, 7.09, 7.86, 8.27; IR (mull) 3326, 3079, 3041, 2591, 1740, 1634, 1557, 1513, 1441, 1424, 1231, 1185, 1115, 855, 833, cm⁻¹. MS (FAB) *m/z* 439 (MH⁺, 99), 453, 440, 439, 395, 331, 177, 133, 118, 88, 23. Anal. Found 53.18; H, 5.86; N, 5.91.

EXAMPLE 17 (COMPARISON) : (S)-[4-[2-[(4-Carboxy-3,3-dimethyl-1-oxobutyl)amino]-3-oxo-3-(pentylamino)propyl]phenoxy]propanedioic acid

[0123] 0.96 g as an amorphous solid. ¹H NMR (DMSO) δ 0.85, 1.11-1.40, 2.12, 2.67, 2.85, 2.98, 5.22, 5.73, 6.78, 7.11, 7.84, 7.97. IR (mull) 3296, 3069, 2729, 2670, 2606, 1727, 1612, 1557, 1512, 1329, 1231, 1184, 1115, 830, 720, cm⁻¹. Anal. Found: C, 55.71; H, 6.90; N, 5.32.

EXAMPLE 18 (COMPARISON) : (S)-[4-[2-[(2-Carboxybenzoyl)amino]-3-oxo-3-(pentylamino)propyl]phenoxy]propanedioic acid

[0124] 0.070 g as an amorphous solid. ¹H NMR (DMSO) δ 0.83, 1.22, 1.40, 2.73, 3.08, 4.50, 5.27, 6.83, 6.97, 7.15, 7.50, 7.70, 7.89, 8.52; IR (mull) 3341, 3065, 3033, 1720, 1627, 1599, 1580, 1538, 1512, 1488, 1271, 1232, 1185, 1147, 1103, cm⁻¹. MS (FAB) *m/z* 501 (MH⁺), 501, 154, 149, 137, 117, 109, 92, 59, 45 41. MS (FAB) *m/z* 501 (MH⁺), 501, 154, 149, 137, 117, 109, 59, 57, 43, 41. Anal. Found: C, 57.36; H, 5.74; N, 5.30.

EXAMPLE 19 (COMPARISON) : [2(S)]-[4-[2-[(3-Carboxybicyclo[2.2.2]oct-2-yl)carbonyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]propanedioic acid

[0125] 0.072 g as a white amorphous solid. ¹H NMR (DMSO) δ 0.83, 0.98-1.60, 1.76, 2.00, 2.47-3.11, 4.35, 5.20, 6.75, 7.06, 7.5-7.9; IR (mull) 3326, 1772, 1732, 1704, 1639, 1547, 1512, 1355, 1340, 1292, 1228, 1183, 1106, 1085, 907, cm⁻¹. MS (FAB) *m/z* 533 (MH⁺), 353, 309, 194, 136, 107, 88, 43, 41, 39, 23. Anal. Found: C, 59.26; H, 6.77; N, 4.84.

EXAMPLE 20 (COMPARISON) : (S)-[4-[2-[(1,1-Dimethylethoxy)carbonyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]propanedioic acid

[0126] 0.203 g as a low melting white solid. ¹H NMR (MeOH) δ 0.88, 1.16-1.58, 2.80, 2.97, 3.10, 4.19, 5.23, 6.89, 7.16, 7.80; IR (mull) 3326, 1738, 1693, 1639, 1614, 1588, 1512, 1393, 1297, 1233, 1169, 1114, 1085, 1051, 1022, cm⁻¹. MS (FAB) *m/z* 453 (MH⁺), 453, 409, 397, 370, 353, 194, 88, 57, 41, 29. Anal. Found: C, 59.08; H, 7.15; N, 6.23.

EXAMPLE 21 (COMPARISON) : (S)-3-[[[1-[4-(Dicarboxymethoxy)phenyl]methyl]-2-oxo-2-(pentylamino)ethyl]amino]carbonyl]pentanedioic acid

[0127] 0.037 g as an amorphous solid. ¹H NMR (DMSO) δ 0.83, 1.22, 2.24, 2.69, 2.98, 4.30, 5.21, 6.77, 7.08, 7.60, 8.10; HRMS (FAB) found 511.1933. Anal. Found: C, 51.39; H, 6.18; N, 5.42.

EXAMPLE 22 (COMPARISON) : [1(S)]-[4-[2-[(2-Carboxycyclopropyl)carbonyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]propanedioic acid

[0128] 0.068 g as an amorphous solid. ¹H NMR (DMSO) δ 0.82, 1.22, 1.92, 2.72, 2.98, 4.32, 5.24, 6.78, 7.09, 7.82, 8.40; IR (mull) 3330, 2730, 2598, 1729, 1612, 1554, 1512, 1229, 1184, 1114, 979, 855, 834, 808, 623, cm⁻¹. MS (FAB) *m/z* 465 (MH⁺), 466, 465, 421, 238, 194, 113, 107, 102, 88, 63. Anal. Found: C, 55.01; H, 6.18; N, 5.74.

EXAMPLE 23 (COMPARISON) : [2(S)]-[4-[2-[(3-Carboxybicyclo[2.2.1]hept-2-yl)carbonyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]propanedioic acid

[0129] 0.049 g. (quant). ¹H NMR (DMSO) δ 0.80, 1.21, 1.43-2.10, 2.12-2.42, 2.60-3.10, 4.34, 5.15, 6.75, 7.05, 7.60, 7.80; IR (mull) 3311, 2727, 2670, 1722, 1632, 1543, 1511, 1295, 1228, 1184, 1113, 1083, 846, 831, 721, cm⁻¹. MS (FAB) *m/z* 519 (MH⁺), 519, 353, 238, 194, 177, 167, 88, 67, 43, 23. Anal. Found: C, 58.39; H, 6.52; N, 4.81.

EXAMPLE 24 (COMPARISON) : [1R-[1(S*),2]]-[4-[2-[(2-Carboxycyclohexyl)carbonyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]propanedioic acid

[0130] 0.109 g as a waxy white solid. ¹H NMR (DMSO) δ 0.82, 1.20, 1.53-1.96, 2.40, 2.54-2.89, 3.00, 4.29, 5.23, 6.76, 7.08, 7.52, 7.90; IR (mull) 3307, 2729, 2671, 1727, 1631, 1543, 1511, 1299, 1226, 1184, 1113, 910, 855, 832, 721, cm⁻¹. MS (FAB) *m/z* 507 (MH⁺), 508, 507, 353, 238, 194, 143, 109, 88, 81, 43. Anal. Found: C, 56.11; H, 6.65; N, 5.31.

EXAMPLE 25 (COMPARISON) : N-[(1,1-Dimethylethoxy)carbonyl]-L-α-aspartyl-O-(dicarboxymethyl)-N-pentyl-L-tyrosinamide (Formula O-1 (n=1), Chart O)

[0131] PREPARATION OF L-2 (n=1, Chart L): To a stirring solution of J-4 (0.25 g) and triethylamine (0.044 g) is added Boc-Asp(OBn)OSu (0.185 g) at 0 °C. The resulting solution is stirred for 16 h allowing the mixture to warm to ambient temperature. The resulting solution is diluted with CH₂Cl₂ (50 ml) and washed with 10% aqueous HCl (3 x 50 ml). The organic layer is dried over MgSO₄ and solvent removed to afford 0.33 g title compound as a white solid. ¹H NMR (CDCl₃) δ 0.84, 1.15-1.43, 2.76, 2.93, 3.09, 4.41, 4.48, 5.10, 5.23, 5.29, 5.44, 5.91, 6.89, 6.85, 7.10, 7.30; IR (mull) 3308, 1744, 1694, 1646, 1527, 1511, 1500, 1355, 1280, 1223, 1169, 1115, 749, 738, 697, cm⁻¹. MS (FAB) *m/z* 838 (MH⁺), 783, 739, 515, 178, 92, 91, 88, 86, 57, 41. Anal. Found: C, 66.86; H, 6.59; N, 5.00.

[0132] PREPARATION OF O-1 (n=1): To a solution of L-2 (0.317 g) in MeOH (15 ml) at ambient temperature is added 10% Pd-C (50 mg) and the resulting mixture hydrogenated at atmospheric pressure for 2h. The mixture is filtered through Celite and solvent removed to afford 0.208 g title compound as a white solid. ¹H NMR (DMSO) δ 0.82, 1.12-1.40, 2.30-2.52, 2.70-3.06, 4.16, 4.32, 5.20, 6.76, 7.60, 7.12, 7.69, 7.78. IR (mull) 3334, 2729, 1726, 1668, 1637, 1512, 1395, 1341, 1280, 1233, 1180, 1165, 1114, 1053, 855, cm⁻¹. MS (FAB) *m/z* 568 (MH⁺), 512, 468, 88, 86, 57, 43, 41, 29, 23. Anal. Found: C, 52.55; H, 6.53; N, 7.08.

EXAMPLE 26 (COMPARISON) : N-(3-Carboxy-1-oxopropyl)-L-α-aspartyl-O-(dicarboxymethyl)-N-pentyl-L-tyrosinamide (Formula L-5, n=1, Chart L)

[0133] PREPARATION OF L-3: To a solution of L-2 (0.263 g) in HOAc (5 ml) is added 1.5 M HCl/ HOAc (5 ml) and the resulting solution allowed to stand for 2 h. The solvent is removed in vacuo to afford 0.24 g (quant) title compound as a white solid. ¹H NMR (DMSO) δ 0.79, 1.17, 2.94, 4.05, 4.40, 5.12, 5.19, 5.81, 6.86, 7.14, 7.34, 8.02, 8.70; IR (mull) 3300, 3034, 1744, 1671, 1649, 1554, 1510, 1500, 1305, 1285, 1221, 1188, 1099, 746, 697, cm⁻¹. MS (FAB) *m/z* 738 (MH⁺), 741, 740, 739, 178, 92, 91, 88, 86, 43.

[0134] PREPARATION OF L-4: To a stirring solution of L-3 (0.10 g) and triethylamine (0.028 g) in CH₂Cl₂ (8 ml) at 0 °C is added succinic anhydride (0.015 g, 0.13 mmol). The resulting solution is stirred for 16 h allowing the solution to warm to ambient temperature. The mixture is diluted with CH₂Cl₂ (50 ml) and washed with 10% aqueous HCl (2 x 50 ml). The organic layer is dried over MgSO₄ and solvent removed to afford 0.076 g title compound as a white solid. ¹H NMR (CDCl₃) δ 0.85, 1.23, 1.40, 2.37, 2.68, 3.14, 4.57, 4.72, 5.01, 5.16, 5.21, 6.36, 6.80, 7.05, 7.25; MS (FAB) *m/z* 838 (MH⁺), 839, 838, 533, 516, 515, 418, 178, 92, 91, 88. Anal. Found: C, 65.53; H, 6.01; N, 4.98.

[0135] PREPARATION OF L-5 (n=1): To a stirring solution of L-4 (0.07 g) in MeOH (5 ml) is added 10% Pd-C (10 mg). The resulting mixture is hydrogenated at atmospheric pressure for 2h and filtered through celite. The solvent is removed to afford 0.044 g title compound as a white amorphous solid. ¹H NMR (DMSO) δ 0.82, 1.20, 2.33, 2.43, 2.72, 2.96, 4.27, 4.42, 5.2, 6.74, 7.06, 7.60, 7.70, 8.30; IR (mull) 3326, 3069, 3036, 2601, 1726, 1638, 1544, 1512, 1407, 1342, 1231, 1183, 1115, 832, 637, cm⁻¹. MS (FAB) *m/z* 568 (MH⁺), 582, 569, 568, 238, 177, 102, 88, 88, 39, 23. Anal. Found: C, 51.15; H, 5.79; N, 6.64.

EXAMPLE 27 (COMPARISON) : N-[(1,1-Dimethylethoxy)carbonyl]-L-α-glutamyl-O-(dicarboxymethyl)-N-pentyl-L-tyrosinamide (Formula O-1 (n=2), Chart O)

[0136] By an analogous procedure as described for Example 25 is obtained 0.21 g title compound as a white amorphous solid. ¹H NMR (DMSO) δ 0.81, 1.18, 1.71, 2.12, 2.96-3.10, 3.82, 4.38, 5.19, 6.76, 6.96, 7.08, 7.77, 7.84; IR

(mull) 3316, 3067, 2730, 2604, 1720, 1639, 1512, 1395, 1341, 1232, 1168, 1106, 1056, 855, 832, cm^{-1} . MS (FAB) m/z 582 (MH^+), 133, 102, 88, 84, 57, 43, 41, 39, 29, 23. Anal. Found: C, 53.72; H, 6.75; N, 6.96.

EXAMPLE 28 (COMPARISON): N-(3-Carboxy-1-oxopropyl)-L- α -glutamyl-O-(dicarboxymethyl)-N-pentyl-L-tyrosinamide (Formula L-5 ($n=2$), Chart L)

[0137] By a procedure analogous to that described for Example 26 is obtained 0.082 g title compound as a white solid. ^1H NMR (DMSO) δ 0.82, 1.20, 1.32, 1.55-1.90, 2.14, 2.38, 2.72, 2.92, 4.11, 4.30, 5.21, 6.77, 7.67, 7.77, 8.08; IR (mull) 3316, 3069, 2730, 2605, 1722, 1634, 1546, 1512, 1414, 1341, 1231, 1183, 1116, 834, 721, cm^{-1} . MS (FAB) m/z 582 (MH^+), 582, 238, 133, 102, 102, 88, 84, 43, 39, 23. Anal. Found: C, 51.51; H, 6.00; N, 6.89.

EXAMPLES 29 - 35 (COMPARISON): (General Synthesis of Formula K-6, Chart K)

[0138] PREPARATION OF K-2: To a stirring solution of H-Tyr-OBu^t (3.0 g) in CH_2Cl_2 (150 ml) at 0 °C is added EDC (2.42 g) and mono-benzyl succinate (2.62 g). The mixture is stirred for 16 h allowing the solution to warm to ambient temperature. The resulting solution is washed with 10% HCl/ H_2O (150 ml), the solvent dried over Na_2SO_4 and evaporated in vacuo. The residue is purified via flash column chromatography (eluant 2:1 hexane/ EtOAc) to afford 5.06 g. ^1H NMR (CDCl_3) δ 1.41, 2.50, 2.67, 2.99, 4.68, 5.11, 5.54, 6.12, 6.71, 6.98, 7.33; IR (liq.) 3351, 2979, 1734, 1654, 1615, 1595, 1516, 1455, 1393, 1369, 1356, 1232, 1155, 752, 698, cm^{-1} . MS (EI) m/z 427 (M^+), 226, 209, 208, 164, 136, 108, 107, 92, 91, 57. Anal. Found: C, 67.17; H, 6.80; N, 3.27.

[0139] PREPARATION OF K-3: To a stirring solution of K-2 (3.55 g) in acetone (175 ml) at ambient temperature is added K_2CO_3 (2.29 g). Dibenzylbromomalonate (3.31 g) is added and the mixture stirred at ambient temperature overnight. The solvent is removed in vacuo and the residue suspended between EtOAc/ H_2O (100 ml each). The layers are shaken, the organic layer separated, dried over Na_2SO_4 , and the solvent removed. The residue is purified via flash column chromatography to afford 1.72 g title compound as a yellow oil. ^1H NMR (CDCl_3) δ 1.39, 2.49, 2.68, 3.00, 4.66, 5.11, 5.23, 5.24, 6.05, 6.82, 7.03, 7.33; MS (EI) m/z 709 (M^+), 368, 312, 108, 107, 91, 79, 77, 57, 56, 55. Anal. Found: C, 69.21; H, 6.15; N, 2.03.

[0140] PREPARATION OF K-4: To a stirring solution of K-3 (1.56 g) in CH_2Cl_2 (40 ml) is added trifluoroacetic acid (100 ml). The resulting solution is stirred for 2h and solvent removed in vacuo to afford 1.42 g (quant) title compound as a slightly pink oil. ^1H NMR (CDCl_3) δ 2.52, 2.67, 3.07, 4.81, 5.10, 5.20, 5.25, 6.43, 6.83, 7.40, 7.31, 7.80; IR (liq.) 3035, 1741, 1641, 1612, 1535, 1511, 1500, 1456, 1388, 1351, 1219, 1180, 1118, 751, 698, cm^{-1} . MS (FAB) m/z 654 (MH^+), 656, 655, 654, 564, 447, 446, 181, 107, 92, 91.

[0141] GENERAL PREPARATION OF K-5: To a stirring solution of K-4 in CH_2Cl_2 (0.035 M) at 0 °C is added EDC (1 eq) followed by the requisite amine. The mixture is stirred for 16h, allowing the solution to warm to ambient temperature. The mixture is diluted with CH_2Cl_2 (50 ml), washed with 10% HCl/ H_2O (2 x 50 ml) and saturated NaHCO_3 . The solvent is dried over Na_2SO_4 and removed in vacuo. Purification from SiO_2 flash column chromatography (eluant 1:1 EtOAc/ hexane) is done to afford title compound.

[0142] GENERAL PREPARATION OF K-6: To a stirring solution of requisite ester K-5 in MeOH (0.03M) is added 10% Pd-C (10% w/w). The resulting mixture is hydrogenated at atmospheric pressure for 3 h and filtered through celite. The solvent is removed in vacuo to afford desired material.

[0143] The following examples (29 - 35) are prepared by the general synthesis of K-6 outlined above, using the appropriate commercially available amines (Chart K).

EXAMPLE 29 (COMPARISON): (S)-[4-[2-[(3-Carboxy-1-oxopropyl)amino]-3-(hexylamino)-3-oxopropyl]phenoxy]propanedioic acid

[0144] 0.052 g as a waxy solid. ^1H NMR (DMSO) δ 0.82, 1.19, 2.32, 2.66, 2.90, 4.32, 5.22, 6.76, 7.09, 7.77, 8.03; IR (mull) 3308, 3069, 3034, 2730, 2597, 1728, 1630, 1550, 1511, 1402, 1341, 1229, 1183, 1111, 833, cm^{-1} . HRMS (FAB) found 467.2040. Anal. Found: C, 55.39; H, 6.56; N, 5.62.

EXAMPLE 30 (COMPARISON): (S)-[4-[2-[(3-Carboxy-1-oxopropyl)amino]-3-[(cyclohexylmethyl)amino]-3-oxopropyl]phenoxy]propanedioic acid

[0145] 0.048 g as a waxy solid. ^1H NMR (DMSO) δ 0.78, 1.12, 1.32, 1.62, 2.30, 2.62-2.92, 4.32, 5.22, 6.78, 7.09, 7.76, 8.02; IR (mull) 3318, 3068, 3034, 2730, 2665, 2599, 1728, 1629, 1550, 1512, 1402, 1350, 1228, 1184, 1108, cm^{-1} . MS (FAB) m/z 479 (MH^+), 480, 479, 435, 331, 177, 127, 114, 71, 57, 55. HRMS (FAB) found 479.2029. Anal. Found: C, 56.68; H, 6.46; N, 5.59.

EXAMPLE 31 (COMPARISON): (S)-[4-[2-[(3-Carboxy-1-oxopropyl)amino]-3-[(2,2-diethoxyethyl)amino]-3-oxopropyl]phenoxy]propanedioic acid

[0146] 0.062 g as a waxy solid. ¹H NMR (DMSO) δ 1.07, 2.30, 2.66, 2.85, 3.15, 3.42, 3.55, 4.38, 5.25, 5.73, 6.78, 7.12, 7.92, 8.06; IR (mull) 3327, 2728, 2669, 2606, 1732, 1638, 1546, 1512, 1351, 1230, 1183, 1116, 1053, 833, 721, cm⁻¹. MS (FAB) *m/z* 499 (MH⁺), 453, 251, 194, 177, 136, 107, 101, 88, 57, 23. Anal. Found: C, 50.80; H, 6.05; N, 5.57.

EXAMPLE 32 (COMPARISON): (S)-[4-[2-[(3-Carboxy-1-oxopropyl)amino]-3-[(3-methylbutyl)amino]-3-oxopropyl]phenoxy]propanedioic acid

[0147] 0.061 g as an amorphous solid. ¹H NMR (DMSO) δ 0.83, 1.22, 1.48, 2.32, 2.75, 2.82, 3.00, 4.35, 5.22, 6.78, 7.09, 7.74, 8.04; IR (mull) 3315, 3065, 2728, 2669, 2603, 1729, 1630, 1549, 1512, 1341, 1229, 1182, 1111, 833, 721, cm⁻¹. MS (FAB) *m/z* 453 (MH⁺), 475, 453, 431, 194, 136, 88, 86, 55, 43, 23. Anal. Found: C, 54.45; H, 6.36; N, 5.85.

EXAMPLE 33 (COMPARISON): (S)-[4-[2-[(3-Carboxy-1-oxopropyl)amino]-3-oxo-3-[[2-(1-piperidinyl)ethyl]amino]propyl]phenoxy]propanedioic acid

[0148] 0.047 g as a white solid. ¹H NMR (DMSO) δ 1.47, 1.66, 2.33, 2.76, 3.05, 4.30, 5.04, 6.64, 7.01, 8.02, 8.16; IR (mull) 3300, 3032, 2729, 2687, 2587, 1722, 1649, 1541, 1512, 1299, 1234, 1182, 1096, 833, 638, cm⁻¹. MS (FAB) *m/z* 494 (MH⁺), 495, 494, 450, 392, 133, 98, 71, 57, 45, 43. Anal. Found: C, 53.22; H, 6.63; N, 7.88.

EXAMPLE 34 (COMPARISON): (S)-[4-[2-[(3-Carboxy-1-oxopropyl)amino]-3-[[3-(4-morpholinyl)propyl]amino]-3-oxopropyl]phenoxy]propanedioic acid

[0149] 0.059 g as a waxy solid. ¹H NMR (DMSO) δ 2.32, 2.50-2.80, 2.97, 3.76, 4.31, 5.10, 6.68, 7.00, 7.79, 8.05; IR (mull) 3303, 2728, 2620, 1725, 1646, 1545, 1512, 1298, 1234, 1183, 1140, 1107, 1049, 834, 638, cm⁻¹. MS (FAB) *m/z* 510 (MH⁺), 511, 510, 466, 463, 308, 241, 177, 100, 57, 39. Anal. Found: C, 50.54; H, 6.12; N, 7.35.

EXAMPLE 35 (COMPARISON): (S)-[4-[2-[(3-Carboxy-1-oxopropyl)amino]-3-[[3-(1H-imidazol-1-yl)propyl]amino]-3-oxopropyl]phenoxy]propanedioic acid

[0150] 0.049 g as an amorphous solid. ¹H NMR (DMSO/DCI) δ 0.93, 1.80-1.93, 2.28, 3.03, 4.02, 5.00, 6.78, 7.12, 7.30, 7.63, 7.71; IR (mull) 3283, 3132, 3032, 1717, 1644, 1565, 1511, 1406, 1332, 1300, 1235, 1181, 1089, 834, 633, cm⁻¹. MS (FAB) *m/z* 491 (MH⁺), 505, 383, 357, 236, 222, 174, 127, 124, 118, 107.

EXAMPLE 36 (COMPARISON): N-(3-Carboxy-1-oxopropyl)-O-(dicarboxymethyl)-L-tyrosyl-L-norleucinamide (Formula M-6, Chart M)

[0151] PREPARATION OF M-2: To a stirring solution of CBZ-L-Tyr-OH (0.50 g) in CH₂Cl₂ (25 ml) at 0 °C is added H-L-Nle-NH₂·HCl (0.264 g), N-methyl morpholine (0.16 g) and EDC (0.303 g). The resulting turbid mixture is stirred for 16 h and allowed to warm to ambient temperature in which time a solid forms. The solid is collected; washed with CH₂Cl₂ and dried in vacuo to afford 0.374 g title compound as a white solid. ¹H NMR (DMSO) δ 0.82, 1.22, 1.40-1.80, 2.59, 2.87, 4.15, 4.92, 6.62, 7.02, 7.28, 7.41, 7.85.

[0152] PREPARATION OF M-3: To a stirring solution of M-2 (0.30 g) in acetone (30 ml) at ambient temperature is added diethyl chloromalonate (0.149 g) and K₂CO₃ (0.194 g). The resulting mixture is stirred for 16 h and solvent removed in vacuo. The residue is suspended between EtOAc/H₂O (100 ml each), the layers shaken, the organic layer separated, dried over Na₂SO₄, and solvent removed in vacuo. The residue is suspended in hexane, sonicated and filtered, to afford 0.295 g title compound as a white solid. ¹H NMR (DMSO) δ 0.83, 1.15, 1.50, 2.68, 2.95, 4.21, 4.92, 5.61, 6.83, 7.26, 7.49, 7.91; IR (mull) 3286, 1747, 1696, 1674, 1643, 1538, 1510, 1334, 1296, 1290, 1260, 1225, 1186, 1028, 698, cm⁻¹. MS (FAB) *m/z* 586 (MH⁺), 586, 569, 434, 384, 305, 265, 176, 92, 91, 86. Anal. Found: C, 61.09; H, 6.56; N, 7.03.

[0153] PREPARATION OF M-4: To a solution of M-3 (0.25 g) in MeOH is added 10% Pd-C (25 mg) and the mixture hydrogenated at atmospheric pressure for 3 h. The mixture is filtered through celite and solvent removed in vacuo to afford 0.168 g title compound as a clear yellow oil. ¹H NMR (DMSO) δ 0.89, 1.13, 1.62, 2.82, 2.75, 3.16, 3.60, 4.32, 5.16, 5.40, 6.11, 6.90, 7.13, 7.75; IR (liq.) 3327, 2959, 2935, 1767, 1746, 1666, 1612, 1511, 1445, 1371, 1298, 1226, 1183, 1097, 1027, cm⁻¹. MS (EI) *m/z* 451 (M⁺), 434, 295, 294, 266, 265, 186, 169, 141, 107, 86.

[0154] PREPARATION OF M-5: To a stirring solution of M-4 (0.157 g) in CH₂Cl₂ (5 ml) at 0 °C is added triethyl amine (0.039 g) followed by succinic anhydride (0.035 g). The resulting solution is stirred overnight allowing the solution to

warm to ambient temperature. CH_2Cl_2 (20 ml) is added and the organics washed with 10% $\text{HCl}/\text{H}_2\text{O}$ (2 x 50 ml). The organic layer is dried over Na_2SO_4 and solvent removed to afford 0.090 g title compound as a waxy solid. ^1H NMR (DMSO) δ 0.083, 1.17, 1.50, 1.65, 2.29, 2.65, 2.95, 4.20, 4.42, 5.60, 6.83, 6.97, 7.14, 7.82, 8.10; IR (mull) 3280, 3206, 1745, 1716, 1676, 1666, 1631, 1548, 1511, 1424, 1324, 1300, 1228, 1184, 1098, cm^{-1} . MS (FAB) m/z 552 (MH^+), 566,

553, 552, 434, 305, 294, 265, 131, 115, 86. Anal. Found: C, 55.96; H, 6.73; N, 7.41.
[0155] PREPARATION OF M-6: To a stirring solution of M-5 (0.050g) in THF/MeOH (1:3 v/v, 5 ml) is added LiOH/ H_2O (2.5 M, 0.18 ml). H_2O (3 ml) is added and the mixture is stirred for 3 h. The solution is diluted with H_2O (20 ml) and extracted with EtOAc (2 x 25 ml). The organic layers are combined, dried over Na_2SO_4 and the solvent removed to afford 0.036 g title compound as a waxy solid. ^1H NMR (DMSO) δ 0.83, 1.19, 1.53, 1.68, 2.30, 2.71, 2.95, 4.02, 4.40, 5.30, 6.79, 6.99, 7.09, 7.13, 7.80, 8.10; HRMS (FAB) found 496.1929.

EXAMPLE 37 (COMPARISON) : N-(1-Oxohexyl)-L- α -aspartyl-O-(dicarboxymethyl)-N-pentyl-L-tyrosinamide (Formula P-2, Chart P)

[0156] PREPARATION OF P-1: To a stirring solution of L-3 (0.25 g) in CH_2Cl_2 (8 ml) at 0 C is added hexanoyl chloride (0.043 g, 0.32). The resulting solution is stirred for 16 h, allowing the mixture to warm to ambient temperature. The solution is diluted with CH_2Cl_2 (20 ml) and washed with 10% $\text{HCl}/\text{H}_2\text{O}$ (3 x 50 ml). The organic layer is dried over Na_2SO_4 and solvent removed to afford 0.164 g title compound as a white solid. ^1H NMR (DMSO) δ 0.81, 1.20, 2.03, 2.54, 2.73, 2.79, 4.32, 4.60, 5.04, 5.19, 5.77, 6.82, 7.06, 7.31, 7.81, 8.11; IR (mull) 3286, 1765, 1739, 1663, 1639, 1543, 1511, 1499, 1299, 1276, 1223, 1170, 749, 733, 695, cm^{-1} . MS (FAB) m/z 836 (MH^+), 837, 533, 515, 418, 178, 92, 91, 88, 88, 43. Anal. Found: C, 68.75; H, 6.95; N, 5.01.

[0157] PREPARATION OF P-2: To a stirring solution of P-1 (0.175 g) in MeOH (15 ml) is added 10% Pd-C (25 mg). The mixture is hydrogenated at atmospheric pressure for 3 h, and filtered through celite. The solvent is removed to afford 0.113 g title compound as a white solid. ^1H NMR (DMSO) δ 0.82, 1.22, 1.44, 2.04, 2.38, 2.62, 2.63-3.10, 4.32, 4.51, 5.19, 6.75, 7.04, 7.74, 8.10; IR (mull) 3308, 3069, 3035, 2731, 2597, 1729, 1636, 1541, 1512, 1418, 1341, 1230, 1183, 1114, 637, cm^{-1} . MS (FAB) m/z 566 (MH^+), 588, 566, 177, 99, 88, 88, 71, 43, 39, 23. Anal. Found: C, 55.95; H, 7.05; N, 7.24.

EXAMPLE 38 (COMPARISON) : (S)-[[4-[3-[[1-[4-(Dicarboxymethoxy)phenyl]methyl]-2-oxo-2-(pentylamino)ethyl]amino]-3-oxopropyl]phenyl]methyl]propanedioic acid (Formula N-4, Chart N)

[0158] PREPARATION OF N-2: To a stirring suspension of 4-formyl cinnamic acid (N-1, 0.50 g) and dibenzylmalonate (0.806 g) is added piperidine (0.290 g) and acetic acid (3 drops). The mixture is refluxed for 16 h, cooled to ambient temperature, and solvent diluted with EtOAc (50 ml). The organics are washed with 10% $\text{HCl}/\text{H}_2\text{O}$ (2 x 50 ml), solvent dried over Na_2SO_4 and evaporated to afford 0.485 g title compound as a white solid. ^1H NMR (DMSO) δ 5.25, 5.27, 6.54, 7.35, 7.50, 7.75; IR (mull) 1724, 1696, 1680, 1628, 1442, 1426, 1414, 1315, 1260, 1224, 1215, 1202, 1185, 699, 694, cm^{-1} . MS (EI) m/z 442 (M^+), 442, 246, 245, 227, 127, 114, 92, 91, 77, 65. Anal. Found: C, 70.77; H, 4.87.

[0159] PREPARATION OF N-3: To a stirring suspension of N-2 (0.194 g) and J-4 (0.25 g) at 0 C is added triethyl amine (0.045 g) and EDC (0.085 g). The mixture is stirred for 16 h allowing the solution to warm to ambient temperature. The mixture is diluted with CH_2Cl_2 (50 ml) and washed with 10% $\text{HCl}/\text{H}_2\text{O}$ (2 x 50 ml). The organic layer is dried over Na_2SO_4 and evaporated to dryness. The residue is purified via flash column chromatography (eluant 1:1 EtOAc/ hexane) to afford 0.038 g title compound as a white solid. ^1H NMR (CDCl_3) δ 0.84, 1.25, 3.00, 3.20, 4.62, 5.21, 5.25, 5.27, 5.28, 5.51, 6.38, 6.87, 7.13, 7.32, 7.54, 7.73; MS (ES^+) 957.

[0160] PREPARATION OF N-4: To a stirring solution of N-3 (0.030 g) in MeOH/ THF (5:3 v/v, 8 ml) is added 10% Pd-C (10 mg). The mixture is hydrogenated at atmospheric pressure for 16 h. The solvent is filtered and evaporated to afford 0.016 g title compound as a white solid. ^1H NMR (DMSO) δ 0.82, 1.33, 2.28, 2.63, 2.80-3.10, 3.2-4.0, 4.40, 5.12, 6.75, 7.06, 7.73, 8.03; MS (FAB) m/z 601 (MH^+), 601, 219, 193, 177, 107, 88, 57, 43, 39, 23.

EXAMPLE 39 (COMPARISON) : (S)-[[4-[2-[(3-Carboxy-1-oxopropyl)amino]-3-oxo-3-(pentylamino)propyl]phenyl]methyl]propanedioic acid (Formula G-7, Chart G)

[0161] PREPARATION OF G-1: To a stirring solution of E-2 (5.0 g) and pyridine (2.58 g) in CH_2Cl_2 (50 ml) at 0 C is added trifluoromethane sulfonic anhydride (4.42 g) dropwise over 15 min. After addition, the mixture is stirred for 1 h at 0 C and diluted with CH_2Cl_2 (50 ml). The organics are washed with 10% $\text{HCl}/\text{H}_2\text{O}$ (2 x 50 ml), separated and dried over Na_2SO_4 . The solvent is filtered through a pad of SiO_2 (50 g) and washed with CH_2Cl_2 (75 ml). The solvent is removed to afford 2.54 g title compound as a yellow solid. ^1H NMR (CDCl_3) δ 0.86, 1.18-1.50, 1.40, 3.09, 4.27, 5.07, 5.90, 7.19, 7.28; IR (mull) 3346, 1683, 1651, 1540, 1523, 1504, 1429, 1271, 1252, 1211, 1169, 1144, 901, 638, 607, cm^{-1} . MS (EI) m/z 482 (M^+), 426, 409, 312, 268, 243, 232, 143, 135, 107, 57. Anal. Found: C, 50.10; H, 5.83; N, 5.88.

[0162] PREPARATION OF G-2: To stirring mixture of G-1 (0.60 g), LiCl (0.158 g) and tributylvinyl tin (0.591 g) in DMF (10 ml) at ambient temperature is added dichlorobis(triphenylphosphine)palladium(II) (0.087 g). The mixture is heated to 90 C and stirred for 16 h. The resulting black mixture is cooled to ambient temperature, poured into ice/ H₂O, and extracted with EtOAc (3 x 75 ml). The organic layers are combined, dried over Na₂SO₄, and solvent removed. The residue is purified via SiO₂ flash column chromatography (eluant 2:1 hexane/ EtOAc) to afford 0.365 g title compound as a white solid. ¹H NMR (CDCl₃) δ 0.84, 1.13-1.40, 1.36, 3.06, 4.23, 5.06, 5.22, 5.68, 5.70, 6.67, 7.15, 7.33; IR (mull) 3337, 1690, 1680, 1656, 1540, 1523, 1331, 1321, 1309, 1299, 1269, 1251, 1170, 906, 631, cm⁻¹. MS (EI) *m/z* 360 (M⁺). 304, 244, 243, 190, 187, 146, 143, 118, 117, 57.

[0163] PREPARATION OF G-3: O₃ (g) is bubbled through a stirring solution of G-2 (0.30 g) in CH₂Cl₂ (50 ml) at -78 C until the blue endpoint is observed. The reaction mixture is capped with a septum and stirred an additional 1.5 h at -78 C. Dimethylsulfide (0.78 g) is added at -78 C and the mixture allowed to warm to ambient temperature (over 1 h). The solvent is removed, the residue taken up in Et₂O (50 ml) and washed with H₂O (2 x 50 ml). The organic layer is dried over Na₂SO₄, solvent removed in vacuo, and residue purified via SiO₂ flash column chromatography (eluant 2:1 hexane/ EtOAc) to afford 0.172 g title compound as a white solid. ¹H NMR (CDCl₃) δ 0.85, 1.15-1.36, 1.39, 3.14, 4.30, 5.05, 5.08, 7.38, 7.81, 10.00; IR (mull) 3324, 1700, 1689, 1651, 1607, 1576, 1529, 1333, 1313, 1300, 1270, 1253, 1224, 1214, 1170, cm⁻¹. MS (EI) *m/z* 362 (M⁺), 306, 289, 243, 193, 192, 149, 148, 143, 120, 57. Anal. Found: C, 65.39; H, 8.43; N, 7.47.

[0164] PREPARATION OF G-4: G-3 (0.50 g), dibenzyl malonate (0.47 g), piperidine (0.023 g), and HOAc (3 drops) in benzene (15 ml) are heated to reflux for 2 h. The mixture is cooled to ambient temperature, diluted with EtOAc (50 ml) and washed with 10% HCl/H₂O (2 x 50 ml). The organic layer is dried over Na₂SO₄, the solvent removed, and the residue purified via SiO₂ flash column chromatography (eluant 2:1 hexane/ EtOAc) to afford 0.060 g title compound as a slightly yellow oil. ¹H NMR (CDCl₃) δ 0.84, 1.17-1.39, 1.40, 3.02, 3.13, 4.21, 5.00, 5.28, 5.67, 7.09, 7.32, 7.72; IR (mull) 3328, 1726, 1685, 1653, 1632, 1542, 1527, 1321, 1267, 1234, 1213, 1200, 1185, 749, 696, cm⁻¹. MS (EI) *m/z* 628 (M⁺), 571, 511, 420, 414, 386, 278, 143, 92, 91, 57. Anal. Found: C, 70.21; H, 7.05; N, 4.48.

[0165] PREPARATION OF G-5: To a stirring solution of G-4 (0.65 g) in HOAc (10 ml) is added 1.5 M HCl/HOAc (10 ml). The mixture is allowed to stand at ambient temperature for 2 h and the solvent removed to afford 0.58 g (quant) title compound as a slightly yellow amorphous solid. ¹H NMR (DMSO) δ 0.76, 1.03, 1.17, 2.89, 3.02, 3.94, 5.25, 5.29, 7.22, 7.36, 7.73, 8.34, 8.41; IR (liq.) 3065, 3034, 2957, 2933, 2872, 2861, 1730, 1669, 1629, 1499, 1456, 1260, 1204, 1185, 697, cm⁻¹. MS (FAB) *m/z* 529 (MH⁺), 1058, 531, 530, 529, 414, 143, 92, 91, 88, 43. Anal. Found: C, 67.22; H, 6.81; N, 4.94.

[0166] PREPARATION OF G-6: To a stirring solution of G-5 (0.507 g) in CH₂Cl₂ (15 ml) at 0 C is added triethylamine (0.20 g) followed by succinic anhydride (0.089 g). The mixture is stirred for 16 h allowing to warm to ambient temperature. The mixture is diluted with CH₂Cl₂ (50 ml) and washed with 10% HCl/ H₂O (2 x 50 ml). The organic layer is dried over Na₂SO₄ and solvent removed to afford 0.557 g title compound as a yellow amorphous solid. ¹H NMR (CDCl₃) δ 0.83, 1.11-1.28, 2.47, 2.63, 3.03, 4.57, 5.25, 5.27, 5.93, 6.88, 7.09, 7.27, 7.70; IR (mull) 3287, 3089, 3064, 3032, 1726, 1640, 1610, 1543, 1356, 1260, 1213, 1198, 1181, 745, 695, cm⁻¹. MS (FAB) *m/z* 629 (MH⁺), 631, 630, 629, 521, 414, 92, 91, 88, 86, 43. Anal. Found: C, 68.45; H, 6.50; N, 4.50.

[0167] PREPARATION OF G-7: To a stirring solution of G-6 (0.15 g) in MeOH (20 ml) is added 10% Pd-C (15 mg) and the mixture hydrogenated at atmospheric pressure for 3 h. The solvent is filtered through celite and removed in vacuo to afford 0.105 g title compound as an amorphous solid. ¹H NMR (DMSO) δ 0.82, 1.15-1.30, 2.31, 2.71, 2.96, 3.09, 4.35, 7.07, 7.77, 8.06; IR (mull) 3300, 2730, 2618, 1716, 1632, 1551, 1517, 1422, 1404, 1244, 1212, 1171, 956, 853, 655, cm⁻¹. MS (FAB) *m/z* 451 (MH⁺), 604, 452, 451, 407, 236, 192, 102, 88, 86, 43. Anal. Found: C, 57.00; H, 6.74; N, 6.06.

EXAMPLE 40 (COMPARISON): (S)-[4-[2-[(3-Carboxy-1-oxopropyl)amino]-3-oxo-3-(pentylamino) propyl]phenyl] methylene]propanedioic acid (Formula H-1, Chart H)

[0168] To a stirring solution of G-6 (0.15 g) in THF/ MeOH (3:1 v/v, 5 ml) at ambient temperature is added LiOH/ H₂O (2.5 M, 0.52 ml). The mixture is stirred for 3 h and acidified to pH- 4 with 10% HCl/ H₂O. The aqueous layer is extracted with EtOAc (2 x 50 ml), the organic layers combined and dried over Na₂SO₄. The solvent is removed to afford 0.062 g title compound as a waxy solid. ¹H NMR (DMSO) δ 0.82, 1.13-1.17, 2.30, 2.78, 2.96, 7.20, 7.29, 7.45, 7.81, 8.10; IR (mull) 3302, 3066, 3031, 2624, 1715, 1630, 1550, 1515, 1442, 1422, 1243, 1212, 1187, 699, 665, cm⁻¹. MS (FAB) *m/z* 449 (MH⁺), 481, 450, 449, 363, 177, 133, 118, 88, 86, 43.

EXAMPLES 41- 50 (COMPARISON): (General Synthesis of Formula A-5, Chart A)

[0169] PREPARATION OF A-2: To a mixture of L-tyrosine benzyl ester p-toluenesulfonate salt (5.0 g) and triethylamine in CH₂Cl₂ (25 mL) at 0 C is added EDC (2.2 g) and monomethyl succinate (1.5 g). The mixture is warmed to

room temperature and stirred overnight. The mixture is diluted with EtOAc (150 mL), and washed with 1 M HCl (50 mL), sat. NaHCO₃ (50 mL), and sat. NaCl (50 mL). The organic phase is dried (MgSO₄), and the solvent is removed under reduced pressure. The residue is purified by flash chromatography (SiO₂, 60% EtOAc/hexane) to provide 3.4 g of title compound as a colorless oil which slowly solidified to a white solid: ¹H NMR (CDCl₃) δ 7.33, 6.84, 6.65, 6.15, 5.62, 5.14, 4.87, 3.67, 3.01, 2.64, 2.47; MS (EI) *m/z* 385 (M⁺), 254, 209, 208, 147, 136, 132, 115, 107, 91, 55; Anal. Found: C, 65.41; H, 6.00; N, 3.61.

[0170] PREPARATION OF A-3: To a mixture of A-2 (2.64 g) and K₂CO₃ (1.9 g) in acetone (20 mL) is added diethyl chloromalonate (2.2 mL). The mixture is stirred vigorously for 18 h. The mixture is partitioned between EtOAc and H₂O. The organic phase is washed with sat. NaHCO₃ and sat. NaCl. After drying (MgSO₄), the solvent is removed under reduced pressure. The residue is purified by flash chromatography (150 g SiO₂, 50% EtOAc/hexane) to provide 2.9 g of A-3 as a colorless oil: ¹H NMR (CDCl₃) δ 7.37, 7.29, 6.90, 6.80, 6.06, 5.12, 4.86, 4.31, 3.67, 3.05, 2.62, 2.48, 1.30; MS (ES-) 542.

[0171] PREPARATION OF A-4: A Parr flask is charged with A-3 (150 mg), 10% Pd/C (25 mg) and abs. EtOH (25 mL), and the mixture is hydrogenated (35 psi) for 45 minutes. The mixture is filtered through Celite and concentrated to provide 110 mg (87%) of A-4 as a colorless oil: ¹H NMR (CDCl₃) δ 7.11, 6.88, 6.23, 5.17, 4.79, 4.30, 3.67, 3.15, 3.06, 2.63, 2.28, 1.30; MS (ES-) 452.

[0172] GENERAL PREPARATION OF A-5: To a mixture of A-4 (1 eq.) in CH₂Cl₂ (0.2 M) at 0 °C is added EDC (1 eq.) followed by the requisite amine. The reaction is warmed to room temperature and stirred for 18 h. The mixture is diluted with EtOAc and washed with 1 M HCl, sat. NaHCO₃, and sat. NaCl. The organic phase is dried (MgSO₄) and concentrated under reduced pressure. The residue is dissolved in THF (3 mL), and a solution of LiOH H₂O (6-8 eq.) in H₂O (1 mL) is added. The mixture is stirred for 2-4 h. The mixture is acidified with 1M HCl, and extracted with EtOAc (3x). The combined organic phase is washed with sat. NaCl and dried (MgSO₄). The solvent is removed *in vacuo* to provide A-5.

EXAMPLE 41 (COMPARISON): (S)-[4-[2-[(3-Carboxy-1-oxopropyl)amino]-3-[(3,3-diphenylpropyl) amino]-3-oxopropyl]phenoxy]propanedioic acid (Chart A, A-5)

[0173] ¹H NMR (DMSO) δ 8.04, 7.90, 7.25, 7.12, 6.77, 5.22, 4.34, 3.92, 2.89, 2.65, 2.31, 2.10; MS (ES+) 577; Anal. Found: C, 62.82; H, 5.77; N, 4.75.

EXAMPLE 42 (COMPARISON): (S)-[4-[3-[(1,3-Benzodioxol-5-ylmethyl)amino]-2-[(3-carboxy-1-oxopropyl)amino]-3-oxopropyl]phenoxy]propanedioic acid (Chart A, A-5)

[0174] ¹H NMR (DMSO) δ 8.32, 8.11, 7.10, 6.78, 5.59, 5.95, 5.23, 4.40, 4.14, 3.92, 2.70, 2.30; MS (ES-) 515.

EXAMPLE 43 (COMPARISON): (S)-[4-[2-[(3-Carboxy-1-oxopropyl)amino]-3-oxo-3-[(3-phenyl propyl)amino]propyl]phenoxy]propanedioic acid (Chart A, A-5)

[0175] ¹H NMR (DMSO) δ 8.10, 7.89, 7.25, 7.15, 6.80, 5.23, 4.33, 3.01, 2.89, 2.70, 2.32, 1.61; MS (ES-) 499; Anal. Found: C, 58.46; H, 5.68; N, 5.47.

EXAMPLE 44 (COMPARISON): (S)-[4-[2-[(3-Carboxy-1-oxopropyl)amino]-3-[(1-naphthalenyl methyl)amino]-3-oxopropyl]phenoxy]propanedioic acid (Chart A, A-5)

[0176] ¹H NMR (DMSO) δ 8.45, 8.17, 8.00, 7.92, 7.81, 7.41, 7.41, 7.25, 7.11, 6.77, 5.25, 4.70, 4.50, 3.94, 2.72, 2.32; MS (ES-) 521; HRMS (FAB) found 523.1722.

EXAMPLE 45 (COMPARISON): (S)-[4-[2-[(3-Carboxy-1-oxopropyl)amino]-3-(decylamino)-3-oxopropyl]phenoxy]propanedioic acid (Chart A, A-5)

[0177] ¹H NMR (DMSO) δ 8.06, 7.80, 7.10, 6.78, 5.24, 3.33, 2.98, 2.85, 2.65, 2.30, 1.21, 0.83; HRMS (FAB) found 523.2634; Anal. Found: C, 59.64; H, 7.88; N, 5.25.

EXAMPLE 46 (COMPARISON): (S)-[4-[3-[[2-[4-(Aminosulfonyl)phenyl]ethyl]amino]-2-[(3-carboxy-1-oxopropyl)amino]-3-oxopropyl]phenoxy]propanedioic acid (Chart A, A-5)

[0178] ¹H NMR (DMSO) δ 8.09, 8.00, 7.71, 7.35, 7.27, 5.27, 4.33, 2.84, 2.60-2.77, 2.31; MS (ES-) 564; Anal. Found: C, 48.88; H, 4.95; N, 7.04.

EXAMPLE 47 (COMPARISON) : (S)-[4-[2-[(3-Carboxy-1-oxopropyl)amino]-3-oxo-3-[[[4-(trifluoromethyl)phenyl]methyl]amino]propyl]-phenoxy]-propanedioic acid (Chart A, A-5)

[0179] ¹H NMR (DMSO) δ 8.50, 8.18, 7.62, 7.31, 7.12, 6.80, 5.27, 4.43, 7.32, 3.91, 3.71, 2.33; MS (ES-) 539; Anal. Found: C, 52.27; H, 4.68; N, 5.12.

EXAMPLE 48 (COMPARISON) : (S)-[4-[2-[(3-Carboxy-1-oxopropyl)amino]-3-oxo-3-[(2-phenoxy ethyl)amino]propyl]phenoxy]propanedioic acid (Chart A, A-5)

[0180] ¹H NMR (DMSO) δ 8.15, 8.09, 7.27, 7.10, 6.92, 7.76, 5.23, 4.40, 3.90, 2.87, 2.65, 2.30; MS (FAB) *m/z* 503 (MH⁺), 392, 391, 149, 113, 71, 69, 57, 55, 43, 41; HRMS (FAB) found 503.1656; Anal. Found: C, 57.51; H, 5.79; N, 4.96.

EXAMPLE 49 (COMPARISON) : (S)-[4-[2-[(3-Carboxy-1-oxopropyl)amino]-3-[[2-(4-hydroxyphenyl)ethyl]amino]-3-oxopropyl]phenoxy]propanedioic acid (Chart A, A-5)

[0181] ¹H NMR (DMSO) δ 8.06, 7.90, 7.09, 6.95, 6.79, 6.64, 5.27, 4.31, 3.15, 2.85, 2.63, 2.52, 2.33; MS (ES-) 501; HRMS (FAB) found 503.1656.

EXAMPLE 50 (COMPARISON) : (S)-[4-[2-[(3-Carboxy-1-oxopropyl)amino]-3-[[4-carboxyphenyl]methyl]amino]-3-oxopropyl]phenoxy]propanedioic acid (Chart A, A-5)

[0182] ¹H NMR (DMSO) δ 8.49, 8.20, 7.85, 7.27, 7.13, 6.80, 5.28, 4.45, 4.31, 2.95, 2.71, 2.34; MS (FAB) *m/z* 517 (MH⁺), 517, 391, 149, 113, 71, 69, 57, 55, 43, 41; HRMS (FAB) found 517.1451; Anal. Found: C, 54.60; H, 5.25; N, 4.82.

EXAMPLE 51 (COMPARISON) : (S)-[4-[2-[(4-Amino-1,4-dioxobutyl)amino]-3-oxo-3-(pentylamino)propyl]phenoxy]propanedioic acid (Chart B, B-5)

[0183] PREPARATION OF B-2: To a mixture of Cbz-Try-OH (2 g) in CH₂Cl₂ (75 mL) and DMF (5 mL) at 0 °C is added EDC (1.21 g). After a few minutes, amylamine (0.74 mL) is added, and the mixture is warmed to room temperature and stirred for 4.5 h. 10% HCl (50 mL) is added; and the phases are separated. The organic phase is washed with sat. NaCl (30 mL), dried (MgSO₄), and concentrated. The residue is purified by flash chromatography (75 g SiO₂, 50% EtOAc/hexane) to give 1.7 g of title compound as a white solid:

¹H NMR (CDCl₃) δ 7.34, 7.06, 6.74, 5.53, 5.35, 5.09, 5.05, 4.25, 3.14, 3.15, 2.92, 1.14-1.35, 0.86; MS (EI) *m/z* 384 (M⁺), 234, 233, 226, 177, 162, 147, 127, 107, 92, 91; Anal. Found: C, 68.53; H, 7.14; N, 7.25.

[0184] PREPARATION OF B-3: To a mixture of B-2 (200 mg) and K₂CO₃ (146 mg) in acetone (1.5 mL) is added diethyl chloromalonate (0.17 mL). The mixture is stirred vigorously for 18 h. The mixture is partitioned between EtOAc (10 mL) and H₂O (5 mL). The organic phase is washed with H₂O (1 x 5 mL) and sat. NaCl (1 x 5 mL). After drying (MgSO₄), the solvent is removed under reduced pressure. The residue is purified by flash chromatography (30 g SiO₂, 40% EtOAc/hexane) to provide 140 mg (49%) of title compound (B-3) as a white solid: ¹H NMR (CDCl₃) δ 7.33, 7.10, 6.88, 5.55, 5.31, 5.14, 5.08, 4.30, 3.12, 2.93, 1.30, 0.86; MS (ES-) 541.

[0185] PREPARATION OF B-4: To a solution of B-3 (2.9 g) in abs. EtOH (100 mL) and THF (10 mL) is added 10% Pd/C (0.29 g, moistened with abs. EtOH). The mixture is hydrogenated (40 psi) for 1 h. The mixture is filtered through Celite and concentrated under reduced pressure. The residue is dissolved in a 1 M solution of HCl in HOAc (10 mL). After stirring for several minutes, the mixture is concentrated to 2-3 mL. A large amount of Et₂O is added, and the mixture is cooled to 0 °C. The Et₂O is decanted from the oil which had settled on the flask. The oil is washed with Et₂O, and the Et₂O is decanted again. To the oil is added Et₂O again, and the mixture is sonicated. The oil gradually crystallizes, and the solid is collected to provide 2.0 g of B-4 as an off-white solid: ¹H NMR (DMSO) δ 8.39, 8.23, 7.14, 6.90, 5.63, 4.20, 3.88, 3.05, 2.93, 1.19, 1.1-1.4, 0.82; MS (ES+) 409.1.

[0186] PREPARATION OF B-5: To a suspension of succinamic acid (40 mg) in CH₂Cl₂ (1 mL) at 0 °C is added EDC (65 mg) and HOBt (46 mg), and the mixture is stirred for a few minutes. B-4 (150 mg) and triethylamine (48 mL) are added, and the mixture is warmed to room temperature and stirred overnight. The mixture is partitioned between EtOAc and 1 M HCl. The organic phase is washed with sat. NaHCO₃, sat. NaCl, and dried (MgSO₄). After the solvent is removed under reduced pressure, the residue is purified by flash chromatography (11 g SiO₂, 6% MeOH/CH₂Cl₂) to yield 72 mg of a white solid. To a solution of the solid dissolved in THF (3 mL) and MeOH (1 mL) is added a solution of LiOH H₂O (30 mg) in H₂O (1 mL). The mixture is stirred at room temperature for 2.5 h. The mixture is neutralized with 1 M HCl and extracted with EtOAc (3x). The combined organic phase is washed with sat. NaCl and dried (MgSO₄). The solvent is removed *in vacuo* to give 27 mg of B-5 as a white solid: ¹H NMR (DMSO) δ 8.05, 7.85, 7.27, 1.12, 6.79, 6.75, 5.26, 4.30, 2.89-3.03, 2.89-3.03, 2.60-2.71, 2.12-2.32, 1.11-1.38, 0.83; MS (ES-) 450; HRMS (FAB) found

452.2051.

EXAMPLES 52 - 54: COMPARISON (General Synthesis of Formulae BB-2 and BB-3, Chart BB)

[0187] GENERAL PREPARATION OF BB-1: To a mixture of the N-protected amino acid (0.34 mmol) in CH_2Cl_2 (1 mL) at 0 °C is added EDC (0.34 mmol). After stirring for a few minutes, the compound (B-4) (150 mg) and triethylamine (48 mL) are added, and the mixture is warmed to room temperature and stirred overnight. The mixture is partitioned between EtOAc and 1 M HCl. The organic phase is washed with sat. NaHCO_3 , sat. NaCl, and dried (MgSO_4). The solvent is removed under reduced pressure to provide BB-1 which is used directly in the next step.

[0188] GENERAL PREPARATION OF BB-2: BB-1 (0.34 mmol) is dissolved in a 1 M solution of HCl in acetic acid (4 mL), and stirred at room temperature for 3 h. The solvent is removed under reduced pressure, and the residue is dissolved in a mixture of triethylamine (0.75 mmol) in CH_2Cl_2 (1.5 mL). The mixture is cooled to 0 °C, and succinic anhydride (0.34 mmol) is added. The mixture is warmed to room temperature and stirred overnight. The mixture is partitioned between EtOAc and 1 M HCl, and the organic phase is washed with sat. NaCl and dried (MgSO_4). After the solvent is removed, the residue is dissolved in THF (3 mL), and a solution of LiOH \cdot H_2O (1-2 mmol) in H_2O (1 mL) is added. The mixture is stirred for 2-5 h. The mixture is neutralized with 1 M HCl and extracted with EtOAc (2x). The combined organic phase is washed with sat. NaCl and dried (MgSO_4). The solvent is removed *in vacuo* to provide BB-2.

[0189] GENERAL PREPARATION OF BB-3: Prepared by direct LiOH saponification of BB-1 with isolation as described in the general synthesis of BB-2.

EXAMPLE 52 (COMPARISON): N-[(Phenylmethoxy)carbonyl]-L- α -aspartyl-O-(dicarboxymethyl)-N-pentyl-L-tyrosinamide (Chart BB, BB-3)

[0190] Obtained 130 mg of title compound (BB-3) as a white solid. ^1H NMR (DMSO) δ 8.10, 7.80, 7.45, 7.31, 7.00, 6.78, 5.24, 5.00, 4.32, 2.80-3.00, 2.55-2.70, 1.10-1.35, 0.82; MS (FAB) m/z 602 (MH^+), 602, 392, 391, 149, 113, 91, 74, 71, 57, 43; HRMS (FAB) found 602.2363.

EXAMPLE 53 (COMPARISON): N-[(1,1-Dimethylethoxy)carbonyl]-D- α -aspartyl-O-(dicarboxymethyl)-N-pentyl-L-tyrosinamide (Chart BB, BB-3)

[0191] Obtained 90 mg of title compound (BB-3) as a white solid. ^1H NMR (DMSO) δ 8.04, 7.80, 7.09, 6.85, 7.78, 5.24, 4.35, 4.19, 2.81-3.09, 2.70, 1.34, 1.10-1.40, 0.82; MS (FAB) m/z 568 (MH^+), 468, 238, 194, 136, 133, 88, 57, 43, 41, 29; HRMS (FAB) found 568.2498; Anal. Found: C, 53.89; H, 6.58; N, 7.23.

EXAMPLE 54 (COMPARISON): 4-Benzoyl-N-(3-carboxy-1-oxopropyl)-L-phenylalanyl-O-(dicarboxymethyl)-N-pentyl-L-tyrosinamide (Chart BB, BB-2)

[0192] Obtained 169 mg of title compound (BB-2) as an off-white solid.

^1H NMR (DMSO) δ 8.12, 8.05, 7.75, 7.70-7.75, 7.35, 7.11, 6.81, 5.25, 4.51, 4.35, 2.70-3.05, 2.32, 1.10-1.35, 0.80; MS (FAB) m/z 704 (MH^+), 705, 704, 353, 238, 224, 194, 136, 107, 105, 88; HRMS (FAB) found 704.2804; Anal. Found: C, 62.15; H, 6.03; N, 5.87.

EXAMPLE 55 (COMPARISON): (S)-2-[4-[2-[(3-Carboxy-1-oxopropyl)amino]-3-oxo-3-(pentylamino)propyl] phenoxy]-2-butenedioic acid (Chart E, E-9)

[0193] To a mixture of E-5 (0.37 g) and triethylamine (0.17 mL) in THF (5 mL) is added dimethylacetylene dicarboxylate (0.25 mL), and the mixture is heated at 50 °C overnight. The mixture is diluted with Et_2O , and washed with 1 M HCl and sat. NaCl. The organic phase is dried (Na_2SO_4) and concentrated. The residue is purified by flash chromatography (SiO_2 , 90% EtOAc/hexane) to yield 0.41 g of E-7 as a 1:1 mixture of isomers. To a mixture of (E-7, Chart E) (100 mg) in MeOH (5 mL) is added a solution of LiOH \cdot H_2O (50 mg) in H_2O (1.5 mL), and the mixture is stirred at room temperature overnight. The reaction mixture is concentrated *in vacuo*, and the residue is dissolved in H_2O . After cooling to 0 °C, 1 M HCl is added until pH = 1: The solid which slowly precipitates over several hours at 0 °C is collected and dried to provide 30 mg of E-9 (6:1 mixture of isomers) as a slight yellow solid: ^1H NMR (MeOH) δ 7.18, 6.86, 6.55, 5.10, 4.49, 3.10, 2.87, 2.39-2.54, 1.27-1.45, 0.90; HRMS (FAB) found 465.1856; Anal. Found: C, 54.20; H, 6.04; N, 5.77.

EXAMPLE 56 (COMPARISON) : [2(S)]-[4-[2-[(3-Carboxy-1-oxopropyl)amino]-3-oxo-3-(pentylamino)propyl] phenoxy] butanedioic acid (Chart E, E-8)

[0194] To a suspension of E-4 (50 mg) and triethylamine (18 mL) in THF (0.3 mL) is added a solution of dibenzylacetylene dicarboxylate (35 mg) in THF (0.2 mL). The reaction mixture is heated at 50 °C for 20 h. The mixture is diluted with Et₂O, and washed with 1 M HCl and sat. NaCl. The organic phase is dried (Na₂SO₄), and the solvent is removed *in vacuo*. The residue is purified by flash chromatography (9 g SiO₂, 60% EtOAc/hexane) to provide 48 mg of E-6 as a 1:1 mixture of isomers. A mixture of E-6 (48 mg) and 10% Pd/C (5 mg) in MeOH (2 mL) is stirred under a hydrogen atmosphere (balloon) for 1 h. The mixture is filtered through Celite and concentrated to provide 27 mg of E-8 as a glass:
¹H NMR (DMSO) δ 8.02, 7.78, 7.08, 6.76, 4.92, 4.32, 2.6-3.0, 2.25-2.35, 1.1-1.4, 0.83; MS (FAB) *m/z* 467 (MH⁺), 489, 468, 467, 349, 252, 136, 107, 88, 86, 43; HRMS (FAB) found 467.2040.

EXAMPLE 57 (COMPARISON) : (R)-[4-[2-[(3-Carboxy-1-oxopropyl)amino]-3-oxo-3-(pentylamino)propyl]phenoxy] propanedioic acid (Chart J, J-6)

[0195] Prepared by general method of Chart J from N-t-Boc-D-tyrosine. ¹H NMR (MeOH) δ 7.83, 7.16, 6.90, 5.21, 4.48; MS (ES-) 451; Anal. Found: C, 54.36; H, 6.41; N, 6.22.

EXAMPLE 58: (S)-2-(Carboxymethoxy)-5-[2-[(3-carboxy-1-oxopropyl)amino]-3-oxo-3-(pentylamino)propyl]benzoic acid (Chart Q, Q-6)

[0196] PREPARATION OF Q-2: To a stirring mixture of 3-iodo-L-tyrosine (10.0 g) in dioxane (100 mL), H₂O (50 mL) and 1 M aqueous NaOH (50 mL) is added di-*t*-butyl dicarbonate (7.8 g) at 0° C. The mixture is stirred for 2 h allowing the solution to warm to ambient temperature, and is then washed with EtOAc (2 x 50 mL). The water layer is separated and carefully acidified with 4 M NaHSO₄ · H₂O in a beaker, and is then extracted with EtOAc (2 x 100 mL). The organic layer is dried (NaSO₄) and concentrated to afford 15.1 g of the crude acid as a yellow oil. The acid is suspended in CH₂Cl₂ (200 mL) and cooled with ice to 0 °C. 1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC, 6.2 g) is added and the mixture is stirred for 10 min at 0° C. 1-Pentylamine (3.8 mL) is added, and the mixture is stirred for 16 h allowing the solution to warm to ambient temperature. The mixture is washed with 10% aqueous HCl (2 x 100 mL), the organic layer dried (MgSO₄), and concentrated. The residue is purified by column chromatography (SiO₂, EtOAc), which furnishes 10.0 g of Q-2 as a white solid. ¹H NMR 400 MHz (CDCl₃) δ 0.87, 1.21, 1.30, 1.41, 1.43, 2.93, 3.09-3.20, 4.17, 5.20, 6.00, 6.85, 7.05, 7.50; ¹³C NMR (CDCl₃) δ 13.95, 22.28, 28.26, 28.89, 29.04, 37.34, 39.54, 54.80, 80.37, 85.24, 115.04, 130.47, 130.86, 139.00, 154.27, 155.52, 171.01. Anal. Found: C, 47.5; H, 6.1.

[0197] PREPARATION OF Q-3: Triethylamine (0.61 mL) is added to a stirring suspension of Q-2 (1.05 g), palladium (II) acetate (14 mg) and 1,1'-bis(diphenylphosphino)-ferrocene (DPPF, 73 mg) in DMF/MeOH 4:1 (5 mL). The mixture is saturated with CO (1 atm) and stirred at 70 °C for 16 h. The mixture is extracted with EtOAc (5 mL), and the organic layer is washed with 10% aqueous HCl (2 x 2 mL), dried (MgSO₄) and concentrated. The residue is purified by column chromatography (SiO₂, EtOAc/*n*-hexane 1:2), which furnishes 0.54 g of Q-3 as a white solid. ¹H NMR 500 MHz (CDCl₃) δ 0.86, 1.17, 1.25, 1.36, 1.41, 2.97, 3.12-3.21, 3.92, 4.21, 5.11, 5.81, 6.91, 7.30, 7.67; ¹³C NMR (CDCl₃) δ 13.86, 22.24, 28.26, 28.90, 29.05, 37.77, 39.47, 52.23, 54.69, 80.37, 112.25, 117.82, 127.54, 130.34, 136.76, 160.56, 170.32, 170.74. Anal. Found: C, 61.6; H, 7.7.

[0198] PREPARATION OF Q-4: A mixture of Q-3 (259 mg), methyl bromoacetate (66 mL) and freshly ground K₂CO₃ (96 mg) is suspended in acetone (5 mL). After being stirred for 24 h at ambient temperature, TLC (EtOAc/*n*-hexane 1:1) indicates that not all starting material has been consumed, and additional methyl bromoacetate (60 mL, 0.63 mmol) is added. After stirring for 24 h, H₂O (2 mL) is added and the mixture is extracted with EtOAc (3 mL). The organic layer is dried (MgSO₄) and concentrated. The residue is purified by column chromatography (SiO₂, EtOAc/*n*-hexane 1:1), which furnishes 163 mg of Q-4 as a white solid. ¹H NMR 500 MHz (CDCl₃) δ 0.87, 1.21, 1.28, 1.38, 1.41, 3.00, 3.16, 3.79, 3.88, 4.23, 4.70, 5.10, 6.81, 7.29, 7.66; ¹³C NMR (CDCl₃) δ 13.86, 22.21, 28.22, 28.88, 29.02, 37.51, 39.46, 52.04, 52.20, 55.87, 66.68, 80.19, 114.61, 121.21, 130.35, 132.59, 134.24, 154.25, 156.44, 165.97, 168.95, 170.69. Anal. Found: C, 60.6; H, 7.5.

[0199] PREPARATION OF Q-5: Trifluoroacetic acid (0.38 mL) is carefully added to a stirring solution of Q-4 (159 mg) in CH₂Cl₂ (3 mL) at 0 °C. The mixture is stirred for 4h allowing the solution to warm to ambient temperature. The volatiles are removed *in vacuo* and the residue is partitioned between EtOAc (3 mL) and saturated NaHCO₃ (3 mL). The organic layer is dried (MgSO₄), and concentrated to dryness to afford 130 mg of the crude amine as a colorless oil. The amine is dissolved in CH₂Cl₂ (3 mL) and cooled with ice to 0 °C. Succinic anhydride (33 mg) and triethylamine (101 mL) is added and the mixture is stirred for 16 h allowing the solution to warm to ambient temperature. The mixture is diluted with CH₂Cl₂ (3 mL) and the organic phase is washed with 10 % aqueous HCl (2 x 3 ml), dried (MgSO₄), and concentrated. The residue is purified by column chromatography (SiO₂, mobile impurities are eluted with 5% MeOH

in CH₂Cl₂, and then 5% MeOH/1% HOAc in CH₂Cl₂ to bring of product). The collected fractions are concentrated, and the remaining HOAc is removed by azeotrope with toluene on a rotavapor and then drying over night under high vacuum, which furnishes 109 mg of Q-5 as a white solid. ¹H NMR 400 MHz (MeOH-d₄) δ 0.95, 1.29, 1.38, 1.48, 2.43-2.64, 2.91, 3.13-3.22, 3.82, 3.92, 4.55, 4.83, 7.00, 7.43, 7.72, 7.97; ¹³C NMR (MeOH-d₄) δ 14.75, 23.78, 30.35, 30.52, 30.57, 31.81, 38.19, 40.99, 52.95, 53.00, 56.62, 67.31, 115.68, 122.18, 132.22, 133.74, 135.87, 158.12, 168.53, 171.27, 173.51, 174.98, 176.81. MS (ESI) 481 (M+H). Anal. Found: C, 57.3; H, 6.7.

[0200] PREPARATION OF Q-6: A solution of Q-5 (87 mg) and 2.5 M aqueous LiOH (435 mL) in THF/MeOH/H₂O 3:1:1 (3 mL) is stirred at ambient temperature for 16 h. The reaction mixture is acidified with 10% aqueous HCl and extracted with EtOAc (4 x 2 mL). The organic layer is dried (MgSO₄) and concentrated to dryness which furnished 68 mg of Q-6 as a white solid. ¹H NMR 400 MHz (MeOH-d₄) δ 0.66, 1.00, 1.08, 2.13-2.40, 2.65, 2.84-2.93, 4.28, 4.58, 6.79, 7.20, 7.56, 7.69; ¹³C NMR (MeOH-d₄) δ 14.75, 23.78, 30.34, 30.47, 30.52, 31.74, 38.22, 40.87, 56.58, 67.67, 115.90, 121.69, 132.74, 134.38, 136.40, 158.14, 169.50, 172.65, 173.41, 174.97, 176.76. MS (ESI) 453 (M+H). Anal. Found: C, 54.3; H, 6.3

[0201] Examples 59-64 were prepared according to the general procedure described for BB-2.

EXAMPLE 59 (COMPARISON) : 2-{4-[(2S)-2-[(2S)-3-[4-(benzyloxy)phenyl]-2-[(3-carboxypropanoyl)amino]propanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy}malonic acid (Chart BB, BB-2)

[0202] ¹H NMR (DMSO-d₆) δ 8.05 (d, 1 H), 7.95 (d, 1 H), 7.66 (t, 1 H), 7.36 (m, 5 H), 7.09 (t, 4 H), 6.82 (dd, 4 H), 5.25 (s, 1 H), 5.02 (s, 2 H), 4.34 (m, 2 H), 2.97 (m, 2 H), 2.95-2.90 (m, 2 H), 2.90 (dd, 1 H), 2.61 (dd, 1 H), 2.30 (m, 4 H), 1.33-1.13 (m, 6 H), 0.82 (t, 3 H); MS (FAB) m/z (rel. intensity) 706 (MH⁺, 36), 707 (15), 706 (36), 353 (15), 238 (13), 226 (28), 91 (99), 88 (16), 57 (20), 55 (15), 43 (20); HRMS (FAB) calcd for C₃₇H₄₃N₃O₁₁+H₁ 706.2975; found 706.2986.

EXAMPLE 60 (COMPARISON) : 2-{4-[(2S)-2-[(2R)-3-(4-benzoylphenyl)-2-[(3-carboxypropanoyl)amino]propanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy}malonic acid (Chart BB, BB-2)

[0203] ¹H NMR (DMSO-d₆) δ 8.38 (d, 1 H), 8.10 (d, 1 H), 7.90 (t, 1 H), 7.67 (m, 3 H), 7.55 (m, 4 H), 7.25 (d, 2 H), 7.25 (d, 2 H), 7.79 (d, 2 H), 5.21 (s, 1 H), 4.52 (m, 1 H), 4.38 (m, 1 H), 3.00 (m, 2 H), 2.75-2.90 (m, 2 H), 2.60 (dd, 2 H), 2.27 (m, 4 H), 1.21 (m, 6 H), 0.83 (t, 3 H); MS (FAB) m/z (rel. intensity) 704 (MH⁺, 99), 705 (42), 704 (99), 353 (20), 238 (35), 224 (70), 219 (27), 194 (19), 107 (28), 105 (34), 88 (49); HRMS (FAB) calcd for C₃₇H₄₁N₃O₁₁+H₁ 704.2819, found 704.2822.

EXAMPLE 61 (COMPARISON) : (Chart BB, BB-2) 2-{4-[(2S)-2-[(2S)-2-[(3-carboxypropanoyl)amino]-3-{4-[(2,6-dichlorobenzyl)oxy]phenyl]propanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy}malonic acid

[0204] ¹H NMR (DMSO-d₆) δ 8.10 (d, 1 H), 7.95 (d, 1 H), 7.67 (t, 1 H), 7.54 (m, 2 H), 7.45 (dd, 1 H), 7.11 (m, 4 H), 6.91 (d, 2 H), 6.81 (d, 2 H), 5.26 (s, 1 H), 5.15 (s, 2 H), 4.35 (m, 2 H), 2.98 (m, 2 H), 2.85 (m, 2 H), 2.73 (dd, 1 H), 2.62 (dd, 1 H), 2.33 (m, 4 H), 1.33-1.12 (m, 6 H), 0.81 (t, 3 H); MS (FAB) m/z (rel. intensity) 774 (MH⁺, 99), 776 (70), 775 (53), 774 (99), 391 (97), 294 (48), 161 (54), 159 (80), 149 (63), 136 (39), 88 (46); HRMS (FAB) calcd for C₃₇H₄₁Cl₂N₃O₁₁+H₁ 774.2196, found 774.2224.

EXAMPLE 62 (COMPARISON) : (Chart BB, BB-2) 2-{4-[(2S)-2-[(2S)-3-[4-(tert-butoxy)phenyl]-2-[(3-carboxypropanoyl)amino]propanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy}malonic acid

[0205] ¹H NMR (DMSO-d₆) δ 8.05 (d, 1 H), 7.90 (d, 1 H), 7.62 (t, 1 H), 7.10 (d, 2 H), 6.94 (d, 2 H), 6.80 (d, 2 H), 6.58 (d, 2 H), 5.26 (s, 1 H), 4.33 (m, 2 H), 2.95 (m, 2 H), 2.90-2.70 (m, 3 H), 2.51 (dd, 1 H), 2.40 (m, 9 H), 2.32 (m, 4 H), 1.35-1.10 (m, 6 H), 0.82 (t, 3 H); MS (FAB) m/z (rel. intensity) 672 (MH⁺, 2), 616 (56), 149 (99), 136 (55), 135 (67), 71 (48), 69 (48), 57 (83), 55 (60), 43 (66), 41 (48); HRMS (FAB) calcd for C₃₄H₄₅N₃O₁₁+H₁ 672.3132, found 672.3110.

EXAMPLE 63 (COMPARISON) : (Chart BB, BB-2) 2-{4-[(2S)-2-[(2S)-2-[(3-carboxypropanoyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy}malonic acid

[0206] ¹H NMR (DMSO-d₆) δ 8.10 (d, 1 H), 7.96 (d, 1 H), 7.69 (t, 1 H), 7.15 (m, 7 H), 6.80 (d, 2 H), 5.26 (s, 1 H), 4.35 (m, 2 H), 2.92 (m, 4 H), 2.71 (m, 2 H), 2.80 (m, 4 H), 1.25 (m, 6 H), 0.82 (t, 3 H); MS (FAB) m/z (rel. intensity) 600 (MH⁺, 35), 600 (35), 155 (22), 149 (50), 136 (20), 120 (99), 88 (27), 73 (32), 71 (23), 57 (34), 43 (26); HRMS (FAB) calcd for C₃₀H₃₇N₃O₁₀+H₁ 600.2557, found 600.2579.

EXAMPLE 64 (COMPARISON) : (Chart BB, BB-2) 2-[4-[(2S)-2-[[[(2S)-2-[(3-carboxypropanoyl)amino]-3-(4-methoxyphenyl)propanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0207] ¹H NMR (DMSO-*d*₆) δ 8.02 (d, 1 H), 7.90 (d, 1 H), 7.64 (t, 1 H), 7.09 (m, 4 H), 6.78 (m, 4 H), 5.25 (s, 1 H), 4.35 (m, 2 H), 3.68 (s, 3 H), 2.98 (m, 2 H), 2.9--2.70 (m, 3 H), 2.60 (dd, 1 H), 2.30 (m, 4 H), 1.35-1.10 (m, 6 H), 0.82 (t, 3 H); MS (FAB) *m/z* (rel. intensity) 630 (MH⁺, 81), 631 (28), 630 (81), 353 (25), 250 (25), 238 (23), 177 (30), 161 (28), 150 (99), 121 (44), 88 (39); HRMS (FAB) calcd for C₃₁H₃₉N₃O₁₁+H₁ 630.2662, found 630.2661.

[0208] Examples 65-71 were prepared according to the general procedure described for BB-3.

EXAMPLE 65 (COMPARISON) : (Chart BB, BB-3) 2-[4-[(2S)-2-[[[(2S)-3-(4-benzoylphenyl)-2-[(tert-butoxycarbonyl)amino]propanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0209] ¹H NMR (DMSO-*d*₆) δ 8.00 (d, 1 H), 7.88 (t, 1 H), 7.66 (m, 5 H), 7.53 (t, 2 H), 7.12 (d, 2 H), 7.00 (d, 1 H), 6.80 (d, 2 H), 5.23 (s, 1 H), 4.40 (m, 1 H), 4.30 (m, 1 H), 2.70-3.05 (m, 6 H), 1.10-1.30 (m, 15 H), 0.80 (t, 3 H); MS (FAB) *m/z* (rel. intensity) 704 (MH⁺, 6), 648 (30), 238 (23), 224 (49), 194 (30), 136 (20), 105 (33), 88 (99), 57 (88), 43 (24), 41 (22); HRMS (FAB) calcd for C₃₈H₄₅N₃O₁₀+H₁ 704.3183, found 704.3171.

EXAMPLE 66 (COMPARISON) : (Chart BB, BB-3) 2-[4-[(2S)-2-[[[(2S)-2-[(tert-butoxycarbonyl)amino]-3-[4-[(2,6-dichlorobenzyl)oxy]phenyl]propanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0210] ¹H NMR (DMSO-*d*₆) δ 7.87 (m, 2 H), 7.53 (m, 2 H), 7.46 (m, 1 H), 7.11 (d, 2 H), 6.91 (d, 2 H), 6.89 (m, 1 H), 6.80 (d, 2 H), 5.23 (s, 1 H), 5.15 (s, 2 H), 4.40 (m, 1 H), 4.05 (m, 1 H), 2.97 (m, 2 H); 2.78 (m, 3 H), 2.60 (m, 1 H), 1.35-1.14 (m, 15 H), 0.81 (t, 3 H); MS (FAB) *m/z* (rel. intensity) 774 (MH⁺, 8), 296 (28), 294 (41), 238 (39), 194 (31), 161 (52), 159 (81), 136 (31), 88 (56), 57 (99), 41 (24); HRMS (FAB) calcd for C₃₈H₄₅Cl₂N₃O₁₀+H₁ 774.2560, found 774.2557.

EXAMPLE 67 (COMPARISON) : (Chart BB, BB-3) 2-[4-[(2S)-2-[[[(2S)-2-[(tert-butoxycarbonyl)amino]-3-[4-(tert-butoxy)phenyl]propanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0211] ¹H NMR (DMSO-*d*₆) δ 7.90 (d, 2 H), 7.09 (t, 2 H), 6.85 (d, 1 H), 6.81 (t, 2 H), 5.23 (s, 1 H), 4.40 (m, 1 H), 4.08 (m, 1 H), 2.97 (m, 2 H), 2.77 (m, 3 H), 2.60 (m, 1 H), 1.27-1.14 (m, 24 H), 0.82 (t, *J* = 7 Hz, 3 H); MS (FAB) *m/z* (rel. intensity) 672 (MH⁺, 2), 238 (25), 194 (22), 192 (19), 136 (79), 107 (39), 88 (48), 57 (99), 41 (26), 39 (23), 29 (25); HRMS (FAB) calcd for C₃₅H₄₉N₃O₁₀+H₁ 672.3496, found 672.3491.

EXAMPLE 68 (COMPARISON) : (Chart BB, BB-3) 2-[4-[(2S)-2-[[[(2S)-2-[(tert-butoxycarbonyl)amino]-3-(4-methoxyphenyl)propanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0212] ¹H NMR (DMSO-*d*₆) δ 7.85 (m, 2 H), 7.09 (m, 4 H), 6.78 (m, 5 H), 5.22 (s, 1 H), 4.40 (m, 1 H), 4.02 (m, 1 H), 3.68 (s, 3 H), 2.97 (m, 2 H), 3.90-3.70 (m, 3 H), 2.60 (m, 1 H), 1.34-1.12 (m, 15 H), 0.82 (t, 3 H); MS (FAB) *m/z* (rel. intensity) 630 (MH⁺, 11), 574 (24), 238 (29), 194 (31), 177 (34), 161 (29), 150 (99), 136 (24), 121 (65), 88 (64), 57 (87); HRMS (FAB) calcd for C₃₂H₄₃N₃O₁₀+H₁ 630.3026, found 630.3015.

EXAMPLE 69 (COMPARISON) : (Chart BB, BB-3) 2-[4-[(2S)-2-[[[(2S)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0213] ¹H NMR (DMSO-*d*₆) δ 7.87 (m, 2 H), 7.17 (m, 5 H), 6.90 (d, 1 H), 6.80 (d, 2 H), 5.22 (s, 1 H), 4.41 (m, 1 H), 4.10 (m, 1 H), 2.97 (m, 2 H), 2.83 (m, 2 H), 2.70-2.50 (m, 2 H), 1.34-1.12 (m, 15 H), 0.82 (t, *J* = 3 Hz, H); MS (FAB) *m/z* (rel. intensity) 600 (MH⁺, 27), 600 (27), 544 (33), 238 (22), 136 (22), 133 (22), 120 (99), 88 (61), 57 (87), 43 (18), 41 (20); HRMS (FAB) calcd for C₃₁H₄₁N₃O₉+H₁ 600.2921, found 600.2923.

EXAMPLE 70 (COMPARISON) : (Chart BB, BB-3) 2-[4-[(2S)-2-[[[(2S)-2-[(tert-butoxycarbonyl)amino]propanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0214] ¹H NMR (DMSO-*d*₆) δ 7.80 (m, 1 H), 7.69 (d, 1 H), 7.08 (d, 2 H), 6.95 (d, 1 H), 6.77 (d, 2 H), 5.22 (s, 1 H), 4.35 (m, 1 H), 3.80 (m, 1 H), 2.95 (m, 2 H), 2.95-2.85 (m, 1 H), 2.75 (m, 1 H), 1.35-1.15 (m, 15 H), 1.05 (d, 3 H), 0.82 (t, 3 H); MS (FAB) *m/z* (rel. intensity) 524 (MH⁺, 13), 468 (39), 238 (25), 136 (21), 133 (20), 88 (99), 86 (19), 57 (71), 44 (58), 41 (19), 29 (18); HRMS (FAB) calcd for C₂₅H₃₇N₃O₉+H₁ 524.2607, found 524.2612.

EXAMPLE 71 (COMPARISON) : (Chart BB, BB-3) 2-[4-[(2S)-2-[(2S)-3-[4-(benzyloxy)phenyl]-2-[(tert-butoxycarbonyl)(methyl)amino]propanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0215] Prepared from B-4 and Boc-N-Me-Try(Bzl)-OH by general procedure for BB-3. ¹H NMR (DMSO-*d*₆) δ 7.85 (br m, 2 H), 7.35 (m, 5 H), 7.10 (m, 4 H), 7.88 (br m, 2 H), 6.78 (d, 2 H), 5.24 (s, 1 H), 5.02 (s, 2 H), 5.72 (br m, 1 H), 4.40 (br m, 1 H), 2.95 (m, 4 H), 2.80-7.70 (m, 2 H), 2.42 (br s, 3 H), 1.25 (m, 15 H), 0.83 (t, 3 H); MS (FAB) *m/z* (rel. intensity) 720 (MH⁺, 3), 620 (15), 253 (9), 241 (12), 240 (73), 238 (10), 237 (9), 91 (99), 57 (51), 41 (13), 29 (9); HRMS (FAB) calcd for C₃₉H₄₉N₃O₁₀+H₁ 720.3496, found 720.3511.

GENERAL PROCEDURE FOR THE PREPARATION OF BB-4 (Chart BB):

[0216] Where R6 is t-butyloxycarbonyl (Boc), the Boc group is removed with HCl/acetic acid or HCl/dioxane, and the resulting amine is acylated with the appropriate acid chloride, isocyanate, sulfonyl chloride, or carboxylic acid via standard procedures. Final saponification (as described for BB-2) affords the diacids BB-4.

EXAMPLE 72 (COMPARISON) : (Chart BB, BB-4) 2-[4-[(2S)-2-[(2S)-3-(4-benzoylphenyl)-2-[[3-(4-hydroxyphenyl)propanoyl]amino]propanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0217] ¹H NMR (DMSO-*d*₆) δ 8.10 (d, 1 H), 8.05 (d, 1 H), 7.83 (t, 1 H), 7.65 (m, 3 H), 7.60 (m, 2 H), 7.51 (t, 2 H), 7.41 (d, 2 H), 7.11 (d, 2 H), 6.90 (d, 2 H), 6.80 (d, 2 H), 6.58 (d, 2 H), 5.24 (s, 1 H), 4.57 (m, 1 H), 4.48 (m, 1 H), 2.96 (m, 3 H), 2.80 (m, 3 H), 2.50 (m, 2 H), 2.25 (m, 2 H), 1.31-1.10 (m, 6 H), 0.79 (t, 3 H); MS (FAB) *m/z* (rel. intensity) 752 (MH⁺, 45), 753 (21), 752 (45), 353 (41), 238 (25), 224 (99), 136 (23), 107 (74), 88 (39), 57 (20), 43 (19); HRMS (FAB) calcd for C₄₂H₄₅N₃O₁₀+H₁ 752.3183, found 752.3186.

EXAMPLE 73 (COMPARISON) : (Chart BB, BB-4) 2-[4-[(2S)-2-[(2S)-2-(acetylamino)-3-(4-benzoylphenyl)propanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0218] ¹H NMR (DMSO-*d*₆) δ 8.06 (d, 2 H), 7.79 (t, 1 H), 7.68 (dd, 2 H), 7.62 (m, 3 H), 7.55 (t, 2 H), 7.35 (d, 2 H), 7.10 (d, 2 H), 6.79 (d, 2 H), 5.24 (s, 1 H), 4.65 (m, 1 H), 4.38 (m, 1 H), 3.00 (m, 3 H), 2.88-3.70 (m, 3 H), 1.74 (s, 3 H), 1.33-1.10 (m, 6 H), 0.80 (t, 3 H); MS (FAB) *m/z* (rel. intensity) 646 (MH⁺, 99), 647 (42), 646 (99), 353 (28), 238 (42), 224 (58), 194 (24), 136 (15), 105 (27), 88 (45), 43 (15); HRMS (FAB) calcd for C₃₅H₃₉N₃O₉+H₁ 646.2764, found 646.2770.

EXAMPLE 74 (COMPARISON) : (Chart BB, BB-4) 2-[4-[(2S)-2-[(2S)-2-[(tert-butylamino)carbonyl]amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0219] ¹H NMR (DMSO-*d*₆) δ 7.94 (d, 1 H), 7.80 (t, 1 H), 7.18 (m, 3 H), 7.08 (m, 4 H), 6.78 (d, 2 H), 5.73 (d, 1 H), 5.23 (s, 1 H), 4.35 (m, 1 H), 4.22 (m, 1 H), 2.97 (m, 2 H), 2.85 (m, 2 H), 2.65 (m, 2 H), 1.38-1.14 (m, 6 H), 1.13 (s, 9 H), 0.82 (t, 3 H); MS (FAB) *m/z* (rel. intensity) 599 (MH⁺, 6), 500 (13), 247 (9), 121 (10), 120 (99), 102 (9), 88 (15), 58 (7), 57 (15), 43 (7), 41 (7); HRMS (FAB) calcd for C₃₁H₄₂N₄O₈+H₁ 599.3080, found 599.3088.

EXAMPLE 75 (COMPARISON): (Chart BB, BB-4) 2-[4-[(2S)-2-[(2S)-2-[(methylsulfonyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0220] ¹H NMR (DMSO-*d*₆) δ 8.33 (d, 1 H), 7.86 (t, 1 H), 7.41 (d, 1 H), 7.23 (m, 5 H), 7.14 (d, 2 H), 6.79 (d, 2 H), 5.24 (s, 1 H), 4.45 (m, 1 H), 4.01 (m, 1 H), 2.99 (m, 2 H), 2.85 (m, 2 H), 2.65 (m, 2 H), 2.20 (s, 3 H), 1.35-1.15 (m, 6 H), 0.83 (t, 3 H); MS (FAB) *m/z* (rel. intensity) 578 (MH⁺, 50), 578 (50), 238 (29), 198 (20), 136 (27), 120 (59), 119 (24), 118 (30), 91 (28), 88 (99), 43 (29); HRMS (FAB) calcd for C₂₇H₃₅N₃O₉S+H₁ 578.2172, found 578.2197.

EXAMPLE 76 (COMPARISON) : (Chart BB, BB-4) 2-[4-[(2S)-2-[(2S)-2-[[3-(diethylamino)propanoyl]amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid hydrochloride

[0221] ¹H NMR (DMSO-*d*₆) δ 8.39 (d, 1 H), 8.23 (d, 1 H), 7.86 (t, 1 H), 7.21 (m, 5 H), 7.11 (d, 2 H), 6.78 (d, 2 H), 5.25 (s, 1 H), 4.52 (m, 1 H), 4.39 (m, 1 H), 3.10 (t, 2 H), 2.95 (m, 10 H), 2.75-2.60 (m, 2 H), 1.33-1.15 (m, 6 H), 1.10 (t, 6 H), 0.82 (t, 3 H); MS (FAB) *m/z* (rel. intensity) 627 (MH⁺, 19), 627 (19), 583 (13), 123 (37), 120 (10), 105 (32), 103 (32), 91 (29), 86 (99), 58 (9), 57 (12); HRMS (FAB) calcd for C₃₃H₄₆N₄O₈+H₁ 627.3394, found 627.3402.

EXAMPLE 77 (COMPARISON): (Chart BB, BB-4) 2-(4-((2S,5S)-5-benzyl-13,13-dimethyl-4,7,11-trioxo-2-[(pentylamino)carbonyl]-12-oxa-3,6,10-triazatetradec-1-yl)phenoxy)malonic acid

[0222] ^1H NMR (DMSO- d_6) δ 8.03 (d, 1 H), 7.98 (d, 1 H), 7.76 (t, 1 H), 7.17 (m, 5 H), 7.10 (d, 2 H), 6.80 (d, 2 H), 7.60 (br s, 1 H), 5.24 (s, 1 H), 4.40 (m, 2 H), 2.98 (m, 4 H), 2.88 (m, 2 H), 3.75-3.62 (m, 2 H), 2.17 (m, 2 H), 1.34 (s, 9 H), 1.33-1.13 (m, 6 H), 0.82 (t, 3 H); MS (FAB) m/z (rel. intensity) 671 (MH^+ , 6), 572 (13), 571 (39), 219 (11), 191 (13), 136 (11), 121 (10), 120 (99), 88 (16), 57 (33), 41 (11); HRMS (FAB) calcd for $\text{C}_{34}\text{H}_{46}\text{N}_4\text{O}_{10}+\text{H}_1$ 671.3292, found 671.3300.

EXAMPLE 78 (COMPARISON): (Chart BB, BB-4) 2-(4-((2S)-2-(((2S)-2-[(benzylsulfonyl)amino]-3-phenylpropanoyl)amino)-3-oxo-3-(pentylamino)propyl)phenoxy)malonic acid

[0223] ^1H NMR (DMSO- d_6) δ 8.25 (d, 1 H), 7.82 (t, 1 H), 7.50 (d, 1 H), 7.23 (m, 9 H), 7.11 (m, 3 H), 6.76 (d, 2 H), 5.13 (s, 1 H), 4.45 (m, 1 H), 4.10 (m, 1 H), 3.80 (d, 1 H), 3.65 (d, 1 H), 3.90 (m, 4 H), 3.80-3.63 (m, 2 H), 1.33-1.10 (m, 6 H), 0.81 (t, 3 H); MS (FAB) m/z (rel. intensity) 654 (MH^+ , 31), 654 (31), 210 (26), 120 (40), 91 (99), 88 (31), 69 (21), 57 (21), 55 (20), 43 (26), 41 (15); HRMS (FAB) calcd for $\text{C}_{33}\text{H}_{39}\text{N}_3\text{O}_9\text{S}+\text{H}_1$ 654.2485, found 654.2488.

EXAMPLE 79 (COMPARISON): (Chart BB, BB-4) 2-(4-((2S)-2-(((2S)-2-[(3-methoxypropanoyl)amino]-3-phenylpropanoyl)amino)-3-oxo-3-(pentylamino)propyl)phenoxy)malonic acid

[0224] ^1H NMR (DMSO- d_6) δ 7.98 (t, 2 H), 7.73 (br s, 1 H), 7.17 (m, 5 H), 7.08 (d, 2 H), 6.80 (d, 2 H), 5.23 (s, 1 H), 4.45 (m, 1 H), 4.35 (m, 1 H), 3.38 (m, 2 H), 3.11 (s, 3 H), 2.96 (m, 4 H), 2.72 (m, 2 H), 2.25 (m, 2 H), 1.22 (m, 6 H), 0.83 (t, 3 H); MS (FAB) m/z (rel. intensity) 586 (MH^+ , 69), 587 (26), 586 (69), 542 (18), 353 (28), 251 (19), 234 (31), 206 (35), 121 (18), 120 (99), 88 (24); HRMS (FAB) calcd for $\text{C}_{30}\text{H}_{39}\text{N}_3\text{O}_9\text{S}+\text{H}_1$ 586.2764, found 586.2757.

EXAMPLES 80-##: (General Synthesis of R-1, R-2, R-3 and R-4, Chart R)

[0225] GENERAL PREPARATION OF R-1: Q-4 (0.50 g, 1.0 mmol) was dissolved in a 4 M solution of HCl in dioxane (5 mL). Stirred at room temperature for 2 h. The solvent was removed under reduced pressure. The residue was dissolved in CH_2Cl_2 (5 mL) and triethylamine (0.43 mL, 3.1 mmol), and N-(tert-Butoxycarbonyl)-L-phenylalanine (0.28 g, 1.0 mmol) was added. After the mixture was cooled to 0°C , EDC (0.20 g, 1.0 mmol) and HOBT (0.14 g, 1.0 mmol) was added. The mixture was warmed to room temperature and stirred overnight. The mixture was partitioned between EtOAc and 1 M HCl. The organic phase was washed with sat. NaHCO_3 , sat. NaCl and dried (MgSO_4) and concentrated to a glassy solid (0.54 g). The residue was purified by flash chromatography (40 g SiO_2 60% EtOAc/heptane) to obtain 0.42 g (67%) of R-1 (R = PhCH_2) as a white powder. PNU-178773 ^1H NMR (CDCl_3) δ 7.48 (br s, 1 H), 7.30 (m, 3 H), 7.18 (m, 3 H), 6.76 (d, 1 H), 6.32 (br s, 1 H), 6.00 (br s, 1 H), 4.90 (br s, 1 H), 4.68 (s, 3 H), 4.55 (m, 1 H), 4.25 (m, 1 H), 3.88 (s, 3 H), 3.78 (s, 3 H), 3.07 (m, 4 H), 2.90 (m, 2 H), 1.40-1.20 (m, 15 H), 0.86 (t, 3 H);

MS (ESI-) for $\text{C}_{33}\text{H}_{45}\text{N}_3\text{O}_9$ m/z 626 (M-H) $^-$; HRMS (FAB) calcd for $\text{C}_{33}\text{H}_{45}\text{N}_3\text{O}_9+\text{H}_1$ 628.3234, found 628.3233.

[0226] GENERAL PREPARATION OF R-2: R-2 was prepared by LiOH saponification of R-1 with isolation as described in the general synthesis of BB-2.

EXAMPLE 80: (Chart R, R-2) 5-[(2S)-2-(((2S)-2-[(tert-butoxycarbonyl)amino]propanoyl)amino)-3-oxo-3-(pentylamino)propyl]-2-(carboxymethoxy)benzoic acid

[0227] ^1H NMR (DMSO- d_6) δ 7.87 (br s, 1 H), 7.54 (d, 1 H), 7.52 (br s, 1 H), 7.29 (dd, 1 H), 6.99 (d, 1 H), 6.89 (d, 1 H), 4.71 (s, 2 H), 4.39 (m, 1 H), 3.84 (m, 1 H), 2.97 (m, 3 H), 2.77 (m, 1 H), 1.40-1.15 (m, 15 H), 1.07 (d, 3 H), 0.84 (t, 3 H); MS (ESI-) for $\text{C}_{25}\text{H}_{37}\text{N}_3\text{O}_9$ m/z 522 (M-H) $^-$; MS (FAB) m/z (rel. intensity) 524 (MH^+ , 1), 546 (45), 525 (22), 446 (16), 425 (23), 424 (99), 88 (60), 57 (56), 44 (46), 41 (18), 29 (17); HRMS (FAB) calcd for $\text{C}_{25}\text{H}_{37}\text{N}_3\text{O}_9+\text{NA}_1$ 546.2427, found 546.2438.

EXAMPLE 81: (Chart R, R-2) 5-[(2S)-2-(((2S)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropanoyl)amino)-3-oxo-3-(pentylamino)propyl]-2-(carboxymethoxy)benzoic acid

[0228] ^1H NMR (DMSO- d_6) δ 7.92 (br s, 1 H), 7.54 (br s, 1 H), 7.30 (d, 1 H), 7.18 (m, 6 H), 6.90 (d, 1 H), 4.68 (s, 2 H), 4.40 (m, 1 H), 4.05 (m, 1 H), 2.98 (m, 2 H), 2.83 (m, 2 H), 2.65 (m, 2 H), 1.33-1.10 (m, 15 H), 0.82 (t, 3 H); MS (FAB) m/z (rel. intensity) 600 (MH^+ , 21), 622 (15), 600 (21), 501 (30), 500 (99), 238 (11), 120 (73), 88 (34), 57 (52), 43 (12), 41 (14); HRMS (FAB) calcd for $\text{C}_{31}\text{H}_{41}\text{N}_3\text{O}_9+\text{H}_1$ 600.2921, found 600.2923.

[0229] GENERAL PREPARATION OF R-3: R-3 was prepared by removal of the Boc group from R-1 with HCl/dioxane

followed by coupling with the appropriate carboxylic acid with EDC and HOBT as described for R-1. Final saponification then affords R-3.

EXAMPLE 82: (Chart R, R-3) 2-(carboxymethoxy)-5-[(2S)-2-(((2S)-2-[(3-carboxypropanoyl)amino]-3-phenylpropanoyl)amino)-3-oxo-3-(pentylamino)propyl]benzoic acid

[0230] ^1H NMR (DMSO- d_6) δ 8.10 (d, 1 H), 8.00 (d, 1 H), 7.78 (t, 1 H), 7.53 (d, 1 H), 7.29 (dd, 1 H), 7.16 (m, 5 H), 6.90 (d, 1 H), 4.71 (s, 2 H), 4.37 (m, 2 H), 2.95 (m, 4 H), 2.70 (m, 2 H), 2.29 (m, 4 H), 1.35-1.15 (m, 6 H), 0.82 (t, 3 H); MS (FAB) m/z (rel. intensity) 600 (MH $^+$, 20), 600 (20), 353 (12), 248 (11), 238 (12), 220 (15), 131 (9), 121 (10), 120 (99), 88 (32), 43 (11); HRMS (FAB) calcd for $\text{C}_{30}\text{H}_{37}\text{N}_3\text{O}_{10}+\text{H}_1$ 600.2557, found 600.2564.

EXAMPLE 83: (Chart R, R-3) 2-(carboxymethoxy)-5-[(2S)-2-(((2S)-2-[[2-(1H-indol-3-yl)acetyl]amino]-3-phenylpropanoyl)amino)-3-oxo-3-(pentylamino)propyl]benzoic acid

[0231] ^1H NMR (DMSO- d_6) δ 8.09 (d, 1 H), 7.97 (d, 1 H), 7.88 (t, 1 H), 7.53 (br s, 1 H), 7.31 (m, 3 H), 7.14 (br s, 5 H), 7.01 (m, 2 H), 6.90 (t, 2 H), 4.71 (s, 2 H), 4.48 (m, 1 H), 4.39 (m, 1 H), 3.46 (d, 2 H), 2.97 (m, 4 H), 2.75 (m, 2 H), 1.35-1.15 (m, 6 H), 0.84 (t, 3 H); MS (ESI-) for $\text{C}_{36}\text{H}_{40}\text{N}_4\text{O}_8$ MS m/z 655 (M-H) $^-$; MS (EI) m/z (rel. intensity) 304 (67), 157 (62), 131 (59), 130 (99), 128 (37), 117 (46), 103 (52), 91 (94), 77 (68), 55 (37); HRMS (FAB) calcd for $\text{C}_{36}\text{H}_{40}\text{N}_4\text{O}_8+\text{H}_1$ 657.2924, found 657.2894.

EXAMPLE 84: (Chart R, R-3) 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-(((2S)-3-phenyl-2-[(2-phenylacetyl)amino]propanoyl)amino)propyl]benzoic acid

[0232] ^1H NMR (DMSO- d_6) δ 8.20 (d, 1 H), 8.10 (d, 1 H), 7.90 (t, 1 H), 7.55 (d, 1 H), 7.29 (dd, 1 H), 7.18 (m, 8 H), 6.90 (d, 2 H), 6.90 (d, 1 H), 4.71 (s, 2 H), 4.45 (m, 2 H), 3.33 (m, 2 H), 2.97 (m, 4 H), 2.73 (m, 2 H), 1.35-1.15 (m, 6 H), 0.85 (t, 3 H); MS (ESI-) for $\text{C}_{34}\text{H}_{39}\text{N}_3\text{O}_8$ m/z 616 (M-H) $^-$; MS (EI) m/z (rel. intensity) 265 (40), 120 (56), 118 (26), 117 (29), 92 (50), 91 (99), 89 (30), 77 (28), 65 (54), 51 (25); HRMS (FAB) calcd for $\text{C}_{34}\text{H}_{39}\text{N}_3\text{O}_8+\text{H}_1$ 618.2815, found 618.2799.

EXAMPLE 85: (Chart R, R-3) 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-(((2S)-3-phenyl-2-[(4-phenylbutanoyl)amino]propanoyl)amino)propyl]benzoic acid

[0233] ^1H NMR (DMSO- d_6) δ 8.00 (br d, 2 H), 7.90 (br s, 1 H), 7.55 (br s, 1 H), 7.20 (m, 11 H), 6.90 (br d, 1 H), 4.70 (s, 2 H), 4.45 (m, 2 H), 2.97 (m, 4 H), 2.75 (m, 2 H), 2.36 (t, 2 H), 2.03 (t, 2 H), 1.62 (m, 2 H), 1.36-1.10 (m, 6 H), 0.83 (t, 3 H); MS (ESI-) for $\text{C}_{36}\text{H}_{43}\text{N}_3\text{O}_8$ m/z 644 (M-H) $^-$; MS (EI) m/z (rel. intensity) 189 (41), 120 (63), 119 (82), 104 (41), 91 (93), 73 (49), 65 (50), 64 (99), 63 (73), 59 (35); HRMS (FAB) calcd for $\text{C}_{36}\text{H}_{43}\text{N}_3\text{O}_8+\text{H}_1$ 646.3128, found 646.3113.

EXAMPLE 86: (Chart R, R-3) 5-[(2S)-2-(((2S)-2-(acetyl)amino)-3-phenylpropanoyl)amino)-3-oxo-3-(pentylamino)propyl]-2-(carboxymethoxy)benzoic acid

[0234] ^1H NMR (DMSO- d_6) δ 8.04 (m, 2 H), 7.84 (t, 1 H), 7.53 (d, 1 H), 7.30 (dd, 1 H), 7.19 (m, 5 H), 6.91 (d, 1 H), 4.71 (m, 2 H), 2.95 (m, 4 H), 2.70 (m, 2 H), 1.73 (s, 3 H), 1.35-1.15 (m, 6 H), 0.85 (t, 3 H); MS (ESI-) for $\text{C}_{28}\text{H}_{35}\text{N}_3\text{O}_8$ m/z 540 (M-H), MS (EI) m/z (rel. intensity) 120 (22), 91 (99), 86 (14), 73 (82), 65 (18), 59 (17), 58 (18), 57 (15), 55 (17), 51 (19); HRMS (FAB) calcd for $\text{C}_{28}\text{H}_{35}\text{N}_3\text{O}_8+\text{H}_1$ 542.2502, found 542.2507.

EXAMPLE 87: (Chart R, R-3) 2-(carboxymethoxy)-5-[(2S)-2-(((2S)-2-[(3-methoxypropanoyl)amino]-3-phenylpropanoyl)amino)-3-oxo-3-(pentylamino)propyl]benzoic acid

[0235] ^1H NMR (DMSO- d_6) δ 8.01 (d, 2 H), 7.85 (t, 1 H), 7.52 (d, 1 H), 7.30 (dd, 1 H), 7.22 (m, 5 H), 6.93 (d, 1 H), 4.69 (s, 2 H), 4.02 (m, 2 H), 3.39 (t, 2 H), 3.12 (s, 3 H), 2.96 (m, 4 H), 2.76 (m, 2 H), 2.27 (t, 2 H), 1.37-1.15 (m, 6 H), 0.85 (t, 3 H); MS (ESI-) for $\text{C}_{30}\text{H}_{39}\text{N}_3\text{O}_9$ m/z 584 (M-H) $^-$; MS (EI) m/z (rel. intensity) 120 (72), 119 (30), 91 (95), 86 (54), 84 (78), 73 (33), 57 (34), 55 (51), 51 (99), 50 (34); HRMS (FAB) calcd for $\text{C}_{30}\text{H}_{39}\text{N}_3\text{O}_9+\text{H}_1$ 586.2764, found 586.2757.

EXAMPLE 88: (Chart R, R-3) 2-(carboxymethoxy)-5-[(2S)-2-(((2S)-2-[(4-hydroxybutanoyl)amino]-3-phenylpropanoyl)amino)-3-oxo-3-(pentylamino)propyl]benzoic acid

[0236] ^1H NMR (DMSO- d_6) δ 7.99 (m, 2 H), 7.86 (br s, 1 H), 7.53 (d, 1 H), 7.31 (m, 1 H), 7.19 (m, 5 H), 6.90 (d, 1 H), 4.71 (s, 2 H), 4.40 (m, 2 H), 3.30 (m, 2 H), 2.95 (m, 4 H), 2.72 (m, 2 H), 2.05 (m, 2 H), 1.50 (m, 2 H), 1.38-1.15 (m,

6 H), 0.84 (t, 3 H); MS (ESI-) for $C_{30}H_{39}N_3O_9$ m/z 584 (M-H)⁻; MS (FAB) m/z (rel. intensity) 586 (MH⁺, 1), 587 (15), 188 (11), 131 (10), 120 (99), 118 (18), 88 (21), 79 (14), 77 (11), 59 (10), 43 (14); HRMS (FAB) calcd for $C_{30}H_{39}N_3O_9+H_1$ 586.2764, found 586.2791.

5 EXAMPLE 89: (Chart R, R-3) 5-[(2S,5S)-5-benzyl-13,13-dimethyl-4,7,11-trioxo-2-[(pentylamino)carbonyl]-12-oxa-3,6,10-triazatetradec-1-yl]-2-(carboxymethoxy)benzoic acid

10 [0237] ¹H NMR (DMSO- d_6) δ 8.03 (t, 1 H), 7.86 (t, 1 H), 7.56 (d, 1 H), 7.30 (dd, 1 H), 7.21 (m; 5 H), 6.92 (d, 1 H), 6.58 (br s, 1 H), 4.72 (s, 2 H), 4.42 (m, 2 H), 2.95 (m, 6 H), 2.73 (m, 2 H), 2.17 (m, 2 H), 1.36 (s, 9 H), 1.35-1.15 (m, 6 H), 0.84 (t, 3 H); MS (ESI-) for $C_{34}H_{46}N_4O_{10}$ m/z 669 (M-H)⁻; MS (FAB) m/z (rel. intensity) 671 (MH⁺, 2), 673 (16), 672 (38), 573 (24), 572 (67), 133 (12), 121 (12), 120 (99), 89 (17), 88 (20), 57 (24); HRMS (FAB) calcd for $C_{34}H_{46}N_4O_{10}+H_1$ 671.3292, found 671.3324.

15 EXAMPLE 90: (Chart R, R-3) 5-[(2S,5S)-5-benzyl-4,7,11,11-tetraoxo-2-[(pentylamino)carbonyl]-11 λ 6~thia-3,6,10-triazadodec-1-yl]-2-(carboxymethoxy)benzoic acid

20 [0238] ¹H NMR (DMSO- d_6) δ 8.10 (m, 2 H), 7.85 (t, 1 H), 7.56 (d, 1 H), 7.32 (dd, 1 H), 7.20 (m, 5 H), 6.90 (m, 2 H), 4.72 (s, 2 H), 4.45 (m, 2 H), 2.98 (m, 6 H), 2.82 (s, 3 H), 2.71 (m, 2 H), 2.27 (m, 2 H), 1.35-1.15 (m, 6 H), 0.85 (t, 3 H); MS (ESI-) for $C_{30}H_{40}N_4O_{10}S$ m/z 647 (M-H)⁻; MS (FAB) m/z (rel. intensity) 649 (MH⁺, 41), 650 (15), 649 (41), 297 (14), 269 (25), 133 (15), 131 (12), 120 (99), 88 (30), 79 (40), 43 (18); HRMS (FAB) calcd for $C_{30}H_{40}N_4O_{10}S+H_1$ 649.2543, found 649.2544.

25 EXAMPLE 91: (Chart R, R-3) 2-(carboxymethoxy)-5-[(2S)-2-[(2S)-2-[[2-(3-hydroxyphenyl)acetyl]amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoic acid

30 [0239] ¹H NMR (DMSO- d_6) δ 8.10 (m, 2 H), 7.86 (t, 1 H), 7.55 (d, 1 H), 7.30 (d, 1 H), 7.16 (m, 5 H), 7.00 (t, 1 H), 6.90 (d, 1 H), 6.59 (m, 2 H), 6.49 (d, 1 H), 4.72 (s, 2 H), 4.42 (m, 2 H), 3.32 (m, 2 H), 2.95 (m, 4 H), 2.75 (m, 2 H), 1.35-1.15 (m, 6 H), 0.85 (t, 3 H); MS (ESI+) for $C_{34}H_{39}N_3O_9$ m/z 634 (M+H)⁺; MS (FAB) m/z (rel. intensity) 634 (MH⁺, 7), 120 (99), 107 (31), 91 (27), 74 (46), 69 (28), 57 (29), 55 (28), 43 (36), 41 (26), 23 (28); HRMS (FAB) calcd for $C_{34}H_{39}N_3O_9+H_1$ 634.2764, found 634.2787.

EXAMPLE 92: (Chart R, R-3) 2-(carboxymethoxy)-5-[(2S)-2-[(2S)-2-[[2-(4-hydroxyphenyl)acetyl]amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoic acid

35 [0240] ¹H NMR (DMSO- d_6) δ 8.05 (t, 2 H), 7.86 (t, 1 H), 7.55 (d, 1 H), 7.30 (dd, 1 H), 7.17 (m, 5 H), 6.91 (d, 1 H), 6.84 (d, 2 H), 6.58 (d, 2 H), 4.72 (s, 2 H), 4.45 (m, 2 H), 3.35 (m, 2 H), 2.98 (m, 4 H), 2.75 (m, 2 H), 1.35-1.15 (m, 6 H), 0.85 (t, 3 H); MS (ESI+) for $C_{34}H_{39}N_3O_9$ m/z 634 (M+H)⁺; MS (FAB) m/z (rel. intensity) 634 (MH⁺, 10), 219 (15), 120 (99), 107 (32), 91 (20), 88 (22), 57 (16), 55 (13), 43 (22), 41 (15), 23 (19); HRMS (FAB) calcd for $C_{34}H_{39}N_3O_9+H_1$ 634.2764, found 634.2769.

40 EXAMPLE 93: (Chart R, R-3) 2-(carboxymethoxy)-5-[(2S)-2-[(2S)-2-[[2-(4-methylphenyl)acetyl]amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoic acid

45 [0241] ¹H NMR (DMSO- d_6) δ 8.11 (d, 1 H), 8.06 (d, 1 H), 7.86 (t, 1 H), 7.55 (d, 1 H), 7.28 (dd, 1 H), 7.18 (m, 5 H), 7.00 (d, 2 H), 6.92 (m, 3 H), 4.71 (s, 2 H), 4.42 (m, 2 H), 3.33 (m, 2 H), 2.98 (m, 4 H), 2.73 (m, 3 H), 2.24 (s, 3 H), 1.35-1.15 (m, 6 H), 0.85 (t, 3 H); MS (FAB) m/z (rel. intensity) 632 (MH⁺, 25), 632 (25), 280 (13), 252 (13), 121 (12), 120 (99), 105 (57), 103 (12), 91 (15), 88 (15), 23 (14); HRMS (FAB) calcd for $C_{35}H_{41}N_3O_8+H_1$ 632.2972, found 632.2980.

50 EXAMPLE 94: (Chart R, R-3) 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[[2-(2-[[4-(trifluoromethyl)phenyl]acetyl]amino)propanoyl]amino]propyl]benzoic acid

55 [0242] ¹H NMR (DMSO- d_6) δ 8.32 (d, 1 H), 8.13 (d, 1 H), 7.87 (t, 1 H), 7.57 (m, 3 H), 7.28 (m, 3 H), 7.16 (m, 5 H), 6.90 (d, 1 H), 4.72 (s, 2 H), 4.51 (m, 1 H), 4.41 (m, 1 H), 4.48 (m, 2 H), 2.98 (m, 4 H), 2.75 (m, 2 H), 1.35-1.15 (m, 6 H), 0.84 (t, 3 H); MS (FAB) m/z (rel. intensity) 686 (MH⁺, 28), 686 (28), 159 (23), 139 (36), 121 (17), 120 (99), 105 (25), 103 (26), 91 (30), 88 (26), 23 (19); HRMS (FAB) calcd for $C_{35}H_{38}F_3N_3O_8+H_1$ 686.2689, found 686.2719.

EXAMPLE 95: (Chart R, R-3) 2-(carboxymethoxy)-5-[(2S)-2-[(2S)-2-[[2-(4-methoxyphenyl)acetyl]amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoic acid

[0243] ¹H NMR (DMSO-*d*₆) δ 8.08 (t, 2 H), 7.86 (t, 1 H), 7.55 (d, 1 H), 7.29 (dd, 1 H), 7.17 (m, 5 H), 6.98 (d, 2 H), 6.89 (d, 1 H), 6.76 (d, 2 H), 4.71 (s, 2 H), 4.43 (m, 2 H), 3.70 (s, 3 H), 3.20 (m, 2 H), 2.98 (m, 4 H), 2.75 (m, 2 H), 1.35-1.15 (m, 6 H), 0.85 (t, 3 H); MS (FAB) *m/z* (rel. intensity) 648 (MH⁺, 28), 648 (28), 148 (16), 139 (29), 123 (18), 121 (83), 120 (99), 105 (21), 103 (21), 91 (25), 88 (16); HRMS (FAB) calcd for C₃₅H₄₁N₃O₉+H⁺ 648.2921, found 648.2915.

EXAMPLE 96: (Chart R, R-3) 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phenyl-2-[(3-phenylpropanoyl)amino]propanoyl] amino]propyl]benzoic acid

[0244] ¹H NMR (DMSO-*d*₆) δ 8.03 (t, 2 H), 7.85 (t, 1 H), 7.56 (d, 1 H), 7.31 (dd, 1 H), 7.14 (m, 10 H), 6.91 (d, 1 H), 4.71 (s, 2 H), 4.45 (m, 2 H), 2.96 (m, 4 H), 2.72 (m, 4 H), 2.32 (t, 2 H), 1.35-1.15 (m, 6 H), 0.84 (t, 3 H); MS (ESI-) for C₃₅H₄₁N₃O₈ *m/z* 630 (M-H)⁻; MS (FAB) *m/z* (rel. intensity) 632 (MH⁺, 3), 634 (15), 633 (38), 353 (7), 335 (6), 280 (12), 252 (10), 131 (7), 120 (99), 88 (18), 79 (17); HRMS (FAB) calcd for C₃₅H₄₁N₃O₈+H⁺ 632.2972, found 632.2986.

EXAMPLE 97: (Chart R, R-4) 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(3-phenylpropanoyl)amino]propyl]benzoic acid

[0245] Prepared from Q-4 and hydrocinnamic acid according to the general procedure for R-1.

[0246] ¹H NMR (DMSO-*d*₆) δ 8.08 (d, 1 H), 7.91 (t, 1 H), 7.55 (d, 1 H), 7.19 (m, 3 H), 7.13 (t, 3 H), 6.87 (d, 1 H), 4.70 (s, 2 H), 4.40 (m, 1 H), 2.98 (m, 2 H), 2.85 (dd, 1 H), 2.68 (m, 3 H), 2.32 (t, 2 H), 1.35-1.15 (m, 6 H), 0.83 (t, 3 H); MS (FAB) *m/z* (rel. intensity) 485 (MH⁺, 99), 971 (10), 970 (16), 638 (8), 486 (30), 485 (99), 398 (9), 238 (23), 105 (8), 91 (11), 88 (70); HRMS (FAB) calcd for C₂₆H₃₂N₂O₇+H⁺ 485.2288, found 485.2303.

EXAMPLE 98 (COMPARISON): PNU-176703 (Chart S, S-4) 2-[4-[2-[(2S)-3-(4-benzoylphenyl)-2-[(3-carboxypropanoyl)amino]propanoyl]amino]ethyl]phenoxy]malonic acid

[0247] PREPARATION OF S-2: To a suspension of N-(tert-Butoxycarbonyl)tyramine (S-1, 0.20 g, 0.84 mmol) and K₂CO₃ (0.23 g, 1.7 mmol) in acetone (3 mL) was added diethyl chloromalonate (0.27 mL, 1.7 mmol). The mixture was stirred vigorously at room temperature for 24 h. The mixture was partitioned between EtOAc and H₂O. The organic phase was washed with sat. NaCl and dried (MgSO₄). After the solvent was removed, the residue was purified by flash chromatography (45 g SiO₂, 3% EtOAc/CH₂Cl₂) to provide 0.24 g of S-2 as a colorless oil.

[0248] ¹H NMR (CDCl₃) δ 7.11 (d, 2 H), 6.90 (d, 2 H), 5.16 (s, 1 H), 4.51 (br s, 1 H), 4.32 (m, 4 H), 3.33 (q, 2 H), 2.73 (t, 2 H), 1.42 (s, 9 H), 1.30 (t, 6 H); MS (ESI-) for C₂₀H₂₉NO₇ *m/z* 394 (M-H)⁻.

[0249] PREPARATION OF S-3: S-2 (239 mg, 0.6 mmol) was dissolved in 1 M HCl /acetic acid (4 mL) and stirred at room temperature for 4 h. The solvent was removed under reduced pressure. The residue was dissolved in CH₂Cl₂ (2 mL), and triethylamine (0.25 mL, 1.8 mmol) and Boc-p-Bz-Phe-OH (222 mg, 0.60 mmol) was added. The mixture was cooled to 0°C, and EDC (115 mg, 0.6 mmol) and HOBT (81 mg, 0.6 mmol) was added. The mixture was warmed to room temperature and stirred overnight. The mixture was partitioned between EtOAc and 1 M HCl. The organic phase was washed with sat. NaHCO₃, sat. NaCl, and dried (MgSO₄). After the solvent was removed under reduced pressure, the residue was purified by flash chromatography (25 g SiO₂, 15% EtOAc/CH₂Cl₂) to provide 108 mg of S-3 as a colorless glass.

[0250] ¹H NMR (CDCl₃) δ 7.76 (t, 4 H), 7.59 (t, 1 H), 7.46 (t, 2 H), 7.31 (d, 2 H), 7.01 (d, 2 H), 6.87 (d, 2 H), 5.90 (br s, 1 H), 5.16 (s, 1 H), 4.95 (br s, 1 H), 4.30 (q, 4 H), 3.40 (m, 2 H), 3.12 (m, 2 H), 2.66 (m, 2 H), 1.40 (s, 9 H), 1.30 (t, 6 H); MS (ESI-) for C₃₆H₄₂N₂O₉ *m/z* 645 (M-H)⁻.

[0251] PREPARATION OF S-4: S-3 (108 mg, 0.17 mmol) was dissolved in 1 M HCl /acetic acid (2 mL) and stirred at room temperature for 3 h. The solvent was removed under reduced pressure. The residue was dissolved in CH₂Cl₂ (2 mL) and triethylamine (70 μL, 0.51 mmol). The mixture was cooled to 0°C, and succinic anhydride (17 mg, 0.17 mmol) was added. The mixture was warmed to room temperature and stirred overnight. The mixture was partitioned between EtOAc and 1 M HCl. The organic phase was washed with sat. NaCl, dried (MgSO₄), and concentrated. The residue was dissolved in THF (3 mL), and a solution of LiOH · H₂O (28 mg, 0.68 mmol) in H₂O (1 mL) was added. The mixture was stirred at room temperature for 3 h. The mixture was acidified (1 M HCl) and extracted with EtOAc. The organic phase was washed sat. NaCl and dried (MgSO₄). The solvent was removed *in vacuo* to provide 89 mg of S-4 as an off-white solid.

[0252] ¹H NMR (DMSO-*d*₆) δ 8.18 (d, 1 H), 8.04 (t, 1 H), 7.65 (m, 5 H), 7.55 (t, 2 H), 7.37 (d, 2 H), 7.10 (d, 2 H), 6.82 (d, 2 H), 5.28 (s, 1 H), 4.45 (m, 1 H), 3.22 (m, 2 H), 3.03 (dd, 1 H), 2.80 (dd, 1 H), 2.57 (t, 2 H), 2.31 (m, 4 H); MS (FAB)

m/z (rel. intensity) 591 (MH^+ , 27), 591 (27), 391 (83), 149 (99), 113 (29), 71 (48), 69 (37), 57 (74), 55 (39), 43 (52), 41 (35); HRMS (FAB) calcd for $C_{31}H_{30}N_2O_{10}+H_1$ 591.1978, found 591.1981.

EXAMPLE 99 (COMPARISON): (Chart A, A-5) 2-[4-((2S)-2-[(3-carboxypropanoyl)amino]-3-[(1S)-1-(hydroxymethyl)-3-methylbutyl]amino]-3-oxopropyl]phenoxy]malonic acid

[0253] Prepared by the general procedure described for A-5.

[0254] 1H NMR (DMSO- d_6) δ 7.99 (d, 1 H), 7.43 (d, 1 H), 7.11 (d, 2 H), 6.78 (d, 2 H), 5.25 (s, 1 H), 4.35 (m, 1 H), 7.72 (m, 1 H), 3.25 (dd, 1 H), 3.12 (dd, 1 H), 2.87 (dd, 1 H), 2.65 (dd, 1 H), 2.31 (m, 4 H), 1.52 (m, 1 H), 1.26 (m, 2 H), 0.82 (q, 6 H); MS (FAB) m/z (rel. intensity) 483 (MH^+ , 91), 484 (22), 483 (91), 139 (25), 123 (18), 118 (99), 105 (17), 103 (15), 91 (18), 86 (16), 55 (15); HRMS (FAB) calcd for $C_{22}H_{30}N_2O_{10}+H_1$ 483.1978, found 483.1999.

[0255] Examples 100-113 were prepared according to the general procedure described for BB-3.

EXAMPLE 100 (COMPARISON): (Chart BB, BB-3) 2-[4-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0256] 1H NMR (DMSO- d_6) δ 8.22 (bd, $J=8$ Hz, 1 H), 7.8 (bt, 1 H), 7.0-7.3 (m, 7 H), 6.80 (d overlapping m, $J=8$ Hz, 3 H), 5.21 (s, 1 H); 4.39 (m, 1 H), 4.15 (m, 1 H), 2.6-3.1 (m, 6 H), 1.1-1.4 (m, 6 H), 1.31 (s, 9 H), 0.86 (t, $J=7$ Hz, 3 H).

EXAMPLE 101 (COMPARISON): (Chart BB, BB-3) 2-[4-[(2S)-2-[(2R)-3-(benzylsulfanyl)-2-[(tert-butoxycarbonyl)amino]propanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0257] 1H NMR (DMSO- d_6) δ 7.98 (d, $J=8$ Hz, 1 H), 7.85 (bt, $J=6$ Hz, 1 H), 7.15-7.3 (m, 5 H), 7.07 (d, $J=8$ Hz, 2 H), 6.76 (d, $J=8$ Hz, 2 H), 5.22 (s, 1 H), 4.38 (m, 1 H), 4.12 (m, 1 H), 3.71 (bs, 2 H), 2.4-3.0 (m, 6 H), 1.1-1.4 (m, 6 H), 1.38 (s, 9 H), 0.81 (t, $J=7$ Hz, 3 H);

MS (FAB) m/z (rel. intensity) 646 (MH^+ , 11), 590 (16), 546 (21), 238 (17), 194 (18), 166 (22), 136 (27), 91 (99), 88 (69), 57 (60), 43 (16).

HRMS (FAB) calcd for $C_{32}H_{43}N_3O_9S+H_1$ 646.2798, found 646.2769.

EXAMPLE 102 (COMPARISON): (Chart BB, BB-3) 2-[4-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-(2-naphthyl)propanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0258] 1H NMR (DMSO- d_6) δ 7.99 (bd, $J=8$ Hz, 1 H), 7.8 (m, 3 H), 7.66 (bs, 1 H), 7.3-7.5 (m, 3 H), 7.1 (d, $J=8$ Hz, 2 H), 6.80 (d, $J=8$ Hz, 2 H), 5.24 (s, 1 H), 4.4 (m, 1 H), 4.17 (m, 1 H), 2.7-3.1 (m, 6 H), 1.1-1.4 (m, 6 H), 1.22 (s, 9 H), 0.82 (t, $J=7$ Hz, 3 H);

Anal. Calcd for $C_{35}H_{43}N_3O_9$: C, 64.70; H, 6.67; N, 6.47.

Found: C, 64.76; H, 6.86; N, 6.14.

MS (ESI-) for $C_{35}H_{43}N_3O_9$ m/z 648 ($M-H$) $^-$.

EXAMPLE 103 (COMPARISON): (Chart BB, BB-3) 2-[4-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-(1-naphthyl)propanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0259] 1H NMR (DMSO- d_6) δ 8.08 (bd, $J=8$ Hz, 1 H), 7.9 (m, 2 H), 7.75 (d, $J=7$ Hz, 1 H), 7.5 (m, 2 H), 7.35 (t, $J=7$ Hz, 1 H), 7.25 (m, 1 H), 7.15 (d, $J=8$ Hz, 2 H), 6.82 (d, $J=8$ Hz, 2 H), 5.24 (s, 1 H), 4.44 (m, 1 H), 4.2 (m, 1 H), 2.7-3.1 (m, 6 H), 1.1-1.4 (m, 6 H), 1.22 (s, 9 H), 0.82 (t, $J=7$ Hz, 3 H);

MS (FAB) m/z (rel. intensity) 650 (MH^+ , 37), 606 (38), 391 (27), 294 (83), 170 (80), 153 (90), 141 (39), 136 (27), 88 (83), 57 (99).

Anal. Calcd for $C_{35}H_{43}N_3O_9 \cdot 0.6 H_2O$: C, 63.64; H, 6.75; N, 6.36.

Found: C, 63.94; H, 6.86; N, 5.97.

EXAMPLE 104 (COMPARISON): (Chart BB, BB-3) 2-[4-[(2S)-2-[(2S)-6-[(benzyloxy)carbonyl]amino]-2-[(tert-butoxycarbonyl)amino]hexanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0260] 1H NMR (DMSO- d_6) δ 7.82 (bt, $J=7$ Hz, 1 H), 7.72 (d, $J=8$ Hz, 1 H), 7.32 (m, 5 H), 7.17 (bt, $J=6$ Hz, 1 H), 7.08 (d, $J=8$ Hz, 2 H), 6.87 (d, $J=7$ Hz, 1 H), 6.78 (d, $J=8$ Hz, 2 H), 5.22 (s, 1 H), 4.98 (s, 2 H), 4.35 (m, 1 H), 3.77 (m, 1 H), 2.7-3.1 (m, 6 H), 1.1-1.5 (m, 12 H), 1.35 (s, 9 H), 0.82 (t, $J=7$ Hz, 3 H);

MS (FAB) m/z (rel. intensity) 715 (MH^+ , 5), 615 (10), 133 (8), 92 (9), 91 (99), 88 (16), 84 (24), 57 (31), 43 (6), 41 (8), 29 (7).

HRMS (FAB) calcd for $C_{36}H_{50}N_4O_{11}+H_1$ 715.3554, found 715.3562.

Anal. Calcd for $C_{36}H_{50}N_4O_{11} \cdot 1.2 H_2O$: C, 58.72; H, 7.17; N, 7.61.

Found: C, 58.68; H, 7.05; N, 7.35.

EXAMPLE 105 (COMPARISON): (Chart BB, BB-3) 2-{4-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-4-(methylsulfonyl)butanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy}malonic acid

[0261] 1H NMR (DMSO- d_6) δ 7.84 (bt, J = 6 Hz, 1 H), 7.75 (d, J = 8 Hz, 1 H), 7.09 (d, J = 8 Hz, 2 H), 6.77 (d, J = 8 Hz, 2 H), 7.01 (d, J = 7 Hz, 1 H), 5.21 (s, 1 H), 4.38 (m, 1 H), 3.90 (m, 1 H), 2.7-3.1 (m, 4 H), 2.31 (bt, J = 7 Hz, 2 H), 1.97 (s, 3 H), 1.66 (m, 2 H), 1.1-1.4 (m, 6 H), 1.35 (s, 9 H), 0.82 (t, J = 7 Hz, 3 H);

MS (FAB) m/z (rel. intensity) 584 (MH^+ , 12), 484 (33), 238 (29), 194 (30), 104 (68), 88 (99), 61 (43), 57 (99), 56 (33), 43 (26), 41 (31).

HRMS (FAB) calcd for $C_{27}H_{41}N_3O_9S+H_1$ 584.2642, found 584.2620.

EXAMPLE 106 (COMPARISON): (Chart BB, BB-3) 2-{4-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-4-(methylsulfonyl)butanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy}malonic acid

[0262] 1H NMR (DMSO- d_6) δ 7.9 (bm, 2 H), 7.10 (d, J = 8 Hz, 2 H), 7.05 (bm, 1 H), 6.78 (d, J = 8 Hz, 2 H), 5.23 (s, 1 H), 4.4 (m, 1 H), 3.95 (m, 1 H), 2.5-3.1 (m, 6 H), 2.47 (s, 3 H), 1.8 (m, 2 H), 1.1-1.4 (m, 6 H), 1.36 (s, 9 H), 0.82 (t, J = 7 Hz, 3 H);

[0263] MS (FAB) m/z (rel. intensity) 600 (M^+ , 48), 556 (41), 500 (40), 100 (30), 88 (50), 59 (64), 57 (71), 56 (99), 41 (32).

[0264] Anal. Calcd for $C_{27}H_{41}N_3O_{10}S \cdot 0.6 H_2O$: C, 53.12; H, 6.97; N, 6.88.

[0265] Found: C, 53.13; H, 6.95; N, 6.72.

EXAMPLE 107 (COMPARISON): (Chart BB, BB-3) 2-{4-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-(2,3,4,5,6-pentafluorophenyl)propanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy}malonic acid

[0266] 1H NMR (DMSO- d_6) δ 7.98 (bm, 1 H), 7.91 (bt, J = 6 Hz, 1 H), 7.10 (bd, J = 8 Hz, 2 H), 6.78 (d, J = 8 Hz, 2 H), 5.21 (s, 1 H), 4.4 (m, 1 H), 4.15 (m, 1 H), 2.6-3.1 (m, 6 H), 1.28 (s, 9 H), 1.1-1.4 (m, 6 H), 0.82 (t, J = 7 Hz, 3 H);

MS (FAB) m/z (rel. intensity) 690 (MH^+ , 14), 294 (28), 210 (24), 136 (29), 133 (49), 88 (95), 86 (33), 57 (99), 43 (18), 41 (29), 29 (27).

HRMS (FAB) calcd for $C_{31}H_{36}F_5N_3O_9+H_1$ 690.2449, found 690.2457.

Anal. Calcd for $C_{31}H_{36}F_5N_3O_9 \cdot 0.48 H_2O$: C, 53.32; H, 5.34; N, 6.02.

Found: C, 53.33; H, 5.49; N, 5.70.

EXAMPLE 108 (COMPARISON): (Chart BB, BB-3) 2-{4-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-4-methylpentanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy}malonic acid

[0267] 1H NMR (DMSO- d_6) δ 7.83 (bt, J = 6 Hz, 1 H), 7.69 (bd, J = 8 Hz, 1 H), 7.07 (d, J = 8 Hz, 2 H), 6.91 (m, 1 H), 6.77 (d, J = 8 Hz, 2 H), 5.19 (s, 1 H), 4.38 (m, 1 H), 3.82 (m, 1 H), 2.6-3.0 (m, 4 H), 1.35 (s, 9 H), 1.1-1.6 (m, 9 H), 0.8 (overlapping t and d, 9 H); MS (ESI-) for $C_{28}H_{43}N_3O_9$ m/z 564.1 ($M-H$).

Anal. Calcd for $C_{28}H_{43}N_3O_9$: C, 59.45; H, 7.66; N, 7.43.

Found: C, 59.56; H, 7.71; N, 7.06.

EXAMPLE 109 (COMPARISON): (Chart BB, BB-3) 2-{4-[(2S)-2-[(2S)-3-(benzyloxy)-2-[(tert-butoxycarbonyl)amino]propanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy}malonic acid

[0268] 1H NMR (DMSO- d_6) δ 7.94 (bd, J = 6 Hz, 1 H), 7.83 (bt, 1 H), 7.31 (bs, 5 H), 7.09 (d, J = 8 Hz, 2 H), 6.95 (bd, J = 8 Hz, 1 H), 6.78 (d, J = 8 Hz, 2 H), 5.22 (s, 1 H), 4.45 (s, 2 H), 4.4 (m, 1 H), 4.2 (m, 1 H), 3.51 (m, 2 H), 2.6-3.1 (m, 4 H), 1.1-1.4 (m, 15 H), 0.84 (t, J = 7 Hz, 3 H);

MS (ESI-) for $C_{32}H_{43}N_3O_{10}$ m/z 628.1 ($M-H$).

Anal. Calcd for $C_{32}H_{43}N_3O_{10}$: C, 61.04; H, 6.88; N, 6.67.

Found: C, 60.98; H, 7.14; N, 6.33.

EXAMPLE 110 (COMPARISON): (Chart BB, BB-3) 2-{4-[(2S)-2-[(2S)-4-amino-2-[(tert-butoxycarbonyl)amino]-4-oxobutanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy}malonic acid

[0269] ¹H NMR (DMSO-*d*₆) δ 7.85 (bt, *J* = 6 Hz, 1 H), 7.80 (d, *J* = 8 Hz, 1 H), 7.3 (bs, 1 H), 7.08 (d, *J* = 8 Hz, 2 H), 6.9 (m, 2 H), 6.78 (d, *J* = 8 Hz, 2 H), 5.21 (s, 1 H), 4.3 (m, 1 H), 4.16 (m, 1 H), 2.65-3.1 (m, 4 H), 2.40 (dd, *J* = 15, 7 Hz, 1 H), 2.28 (dd, *J* = 15, 8 Hz), 1.35 (s, 9 H), 1.1-1.4 (m, 6 H), 0.83 (t, *J* = 7 Hz, 3 H);

[0270] MS (FAB) *m/z* (rel. intensity) 567 (MH⁺, 19), 467 (43), 238 (26), 194 (30), 88 (76), 87 (47), 73 (34), 57 (99), 43 (27), 41 (31), 29 (33).

EXAMPLE 111 (COMPARISON): (Chart BB, BB-3) 2-{4-[(2S)-2-[(2-[(benzyloxy)carbonyl]amino) acetyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy}malonic acid

[0271] ¹H NMR (DMSO-*d*₆) δ 8.00 (d, *J* = 8 Hz, 1 H), 7.88 (bt, 1 H), 7.32 (m, 5 H), 7.10 (d, *J* = 8 Hz, 2 H), 6.79 (d, *J* = 8 Hz, 2 H), 5.23 (s, 1 H), 5.00 (s, 2 H), 4.38 (m, 1 H), 3.4-3.7 (m, 2 H), 2.6-3.1 (m, 4 H), 1.1-1.4 (m, 6 H), 0.82 (t, *J* = 7 Hz, 3 H); MS (ESI-) for C₂₇H₃₃N₃O₉ *m/z* 542.2 (M-H)⁻.

Anal. Calcd for C₂₇H₃₃N₃O₉ • 0.4 H₂O: C, 58.87; H, 6.19; N, 7.63.

Found: C, 58.86; H, 6.33; N, 7.41.

EXAMPLE 112 (COMPARISON): (Chart BB, BB-3) 2-{4-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-(4-hydroxyphenyl)propanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy}malonic acid

[0272] ¹H NMR (DMSO-*d*₆) δ 7.8-8.0 (m, 2 H), 7.10 (d, *J* = 8 Hz, 2 H), 6.94 (d, *J* = 8 Hz, 2 H), 6.79 (d, *J* = 8 Hz, 2 H), 6.59 (d overlapping m, *J* = 8 Hz, 3 H), 5.22 (s, 1 H), 4.38 (m, 1 H), 4.0 (m, 1 H), 2.5-3.1 (m, 6 H), 1.1-1.4 (m, 6 H), 0.82 (t, *J* = 7 Hz, 3 H);

MS (FAB) *m/z* (rel. intensity) 616 (MH⁺, 77), 616 (77), 560 (66), 516 (25), 336 (32), 238 (37), 194 (25), 136 (98), 133 (25), 88 (92), 57 (99).

HRMS (FAB) calcd for C₃₁H₄₁N₃O₁₀+H₁ 616.2870, found 616.2860.

Anal. Calcd for C₃₁H₄₁N₃O₁₀ • 1.1 H₂O: C, 58.52; H, 6.86; N, 6.61.

Found: C, 58.52; H, 6.80; N, 6.52.

EXAMPLE 113 (COMPARISON): (Chart BB, BB-3) 2-{4-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-4-phenylbutanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy}malonic acid

[0273] ¹H NMR (DMSO-*d*₆) δ 7.84 (t, *J* = 6 Hz, 1 H), 7.77 (d, *J* = 8 Hz, 1 H), 7.21 (d, *J* = 8 Hz, 2 H), 7.1 (m, 6 H), 6.79 (d, *J* = 8 Hz, 2 H), 5.17 (s, 1 H), 4.4 (m, 1 H), 3.8 (m, 1 H), 2.7-3.0 (m, 4 H), 2.45 (m, 2 H), 1.71 (m, 2 H), 1.37 (s, 9 H), 1.1-1.4 (m, 6 H), 0.79 (t, *J* = 7 Hz, 3 H);

MS (FAB) *m/z* (rel. intensity) 614 (MH⁺, 27), 558 (53), 514 (37), 238 (52), 194 (41), 134 (85), 117 (38), 91 (78), 88 (90), 57 (99), 41 (31).

HRMS (FAB) calcd for C₃₂H₄₃N₃O₉+H₁ 614.3077, found 614.3073.

[0274] Examples 114-115 were prepared by the general procedure described for R-2 (Chart R).

EXAMPLE 114 (COMPARISON): (Chart R, R-2) 5-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]-2-(carboxymethoxy)benzoic acid

[0275] ¹H NMR (DMSO-*d*₆) δ 8.30 (d, *J* = 8 Hz, 1 H), 7.84 (bt, *J* = 6 Hz, 1 H), 7.58 (bs, 1 H), 7.27 (bd, *J* = 8 Hz, 1 H), 7.15 (m, 5 H), 6.88 (d, *J* = 9 Hz, 1 H), 6.79 (d, *J* = 8 Hz, 1 H), 4.66 (s, 2 H), 4.38 (m, 1 H), 4.11 (m, 1 H), 2.5-3.1 (m, 6 H), 1.1-1.4 (m, 6 H), 1.27 (s, 9 H), 0.83 (t, *J* = 7 Hz, 3 H); MS (ESI-) for C₃₁H₄₁N₃O₉ *m/z* 598.4 (M-H)⁻.

HRMS (FAB) calcd for C₃₁H₄₁N₃O₉+H₁ 600.2921, found 600.2930.

EXAMPLE 115: (Chart R, R-2) 5-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-(4-hydroxyphenyl)propanoyl]amino]-3-oxo-3-(pentylamino)propyl]-2-(carboxymethoxy)benzoic acid

[0276] ¹H NMR (DMSO-*d*₆) δ 7.85 (m, 2 H), 7.52 (bs, 1 H), 7.28 (dm, *J* = 8 Hz, 1 H), 6.9 (m, 3 H), 6.78 (d, *J* = 8 Hz, 1 H), 6.59 (d, *J* = 8 Hz, 2 H), 4.68 (s, 2 H), 4.39 (m, 1 H), 3.97 (m, 1 H), 2.5-3.1 (m, 6 H), 1.1-1.4 (m, 6 H), 1.28 (s, 9 H), 0.82 (t, *J* = 7 Hz, 3 H); MS (ESI) for C₃₁H₄₁N₃O₁₀ *m/z* 614.3 (M-H)⁻.

HRMS (FAB) calcd for C₃₁H₄₁N₃O₁₀+H₁ 616.2870, found 616.2866.

[0277] Examples 116-124 were prepared by procedures analogous to that described for Example 10 (Chart D).

EXAMPLE 116 (COMPARISON) : 2-[4-[(2S)-2-[[[(1-carboxy-2-phenylethyl)amino]carbonyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0278] Major diastereomer. ¹H NMR (DMSO-*d*₆) δ 7.79 (bt, *J* = 7 Hz, 1 H), 7.1-7.3 (m, 5 H), 7.04 (d, *J* = 8 Hz, 2 H), 6.78 (d, *J* = 8 Hz, 2 H), 6.35 (d, *J* = 7 Hz, 1 H), 6.30 (d, *J* = 7 Hz, 1 H), 5.26 (s, 1 H), 4.23 (m, 2 H), 2.6-3.1 (m, 6 H), 1.1-1.4 (m, 6 H), 0.82 (t, *J* = 7 Hz, 3 H); MS (FAB) *m/z* (rel. intensity) 544 (MH⁺, 99), 545 (31), 544 (99), 353 (24), 238 (21), 194 (18), 166 (23), 136 (16), 120 (42), 88 (20), 43 (17). Minor diastereomer (apparent peaks): ¹H NMR (DMSO-*d*₆) δ 6.96 (d, *J* = 8 Hz), 5.22 (s);

HRMS (FAB) calcd for C₂₇H₃₃N₃O₉+H₁ 544.2295, found 544.2310.

Anal. Calcd for C₂₇H₃₃N₃O₉ • 1.9 H₂O: C, 56.13; H, 6.42; N, 7.27.

Found: C, 56.13; H, 6.06; N, 7.26.

EXAMPLE 117 (COMPARISON) : 2-[4-[(2S)-2-[[[benzyl(4-carboxybenzyl)amino]carbonyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0279] ¹H NMR (DMSO-*d*₆) δ 7.85 (d, *J* = 8 Hz, 2 H), 7.24 (m, 5 H), 7.06 (m, 4 H), 6.77 (d, *J* = 8 Hz, 2 H), 5.27 (s, 1 H), 4.4 (m, 5 H), 2.7-3.1 (m, 4 H), 1.1-1.4 (m, 6 H), 0.83 (t, *J* = 7 Hz, 3 H); MS (FAB) *m/z* (rel. intensity) 620 (MH⁺, 25), 621 (9), 620 (25), 139 (10), 135 (13), 107 (7), 105 (8), 103 (8), 92 (8), 91 (99), 43 (7).

HRMS (FAB) calcd for C₃₃H₃₇N₃O₉+H₁ 620.2607, found 620.2621.

Anal. Calcd for C₃₃H₃₇N₃O₉ • 0.81 H₂O: C, 62.49; H, 6.14; N, 6.63.

Found: C, 62.49; H, 6.10; N, 6.73.

EXAMPLE 118 (COMPARISON) : 2-[4-[(2S)-2-[[[4-(carboxymethyl)benzyl][3-(trifluoromethyl)benzyl]amino]carbonyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0280] ¹H NMR (DMSO-*d*₆) δ 7.76 (bt, *J* = 7 Hz, 1 H), 7.4-7.6 (m, 4 H), 7.3 (bd, *J* = 7 Hz, 1 H), 7.15 (d, *J* = 8 Hz, 2 H), 7.07 (d, *J* = 9 Hz, 2 H), 7.01 (d, *J* = 8 Hz, 2 H), 6.75 (d, *J* = 9 Hz, 2 H), 5.24 (s, 1 H), 4.2-4.5 (m, 5 H), 3.52 (s, 2 H), 2.7-3.1 (m, 4 H), 1.1-1.4 (m, 6 H), 0.83 (t, *J* = 7 Hz, 3 H);

MS (FAB) *m/z* (rel. intensity) 702 (MH⁺, 21), 702 (21), 324 (14), 322 (8), 159 (27), 150 (10), 149 (99), 107 (11), 105 (19), 104 (18), 91 (10).

HRMS (FAB) calcd for C₃₅H₃₈F₃N₃O₉+H₁ 702.2638, found 702.2637.

Anal. Calcd for C₃₅H₃₈F₃N₃O₉ • 0.44 H₂O: C, 59.24; H, 5.52; N, 5.92.

Found: C, 59.24; H, 5.56; N, 5.89.

EXAMPLE 119 (COMPARISON) : 2-[4-[(2S)-2-[[[1-[4-(benzyloxy)benzyl]-2-hydroxy-2-oxoethyl]amino]carbonyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0281] ¹H NMR (major isomer peaks)(DMSO-*d*₆) δ 7.3-7.5 (m, 5 H), 7.0-7.1 (m, 4 H), 6.88 (d, *J* = 8 Hz, 2 H), 6.78 (d, *J* = 8 Hz, 2 H), 5.26 (s, 1 H), 5.04 (s, 2 H), 4.2 (m, 2 H), 2.6-3.0 (m, 6 H), 1.1-1.4 (m, 6 H), 0.81 (t, *J* = 7 Hz, 3 H);

MS (FAB) *m/z* (rel. intensity) 544 (MH⁺, 99), 545 (31), 544 (99), 353 (24), 238 (21), 194 (18), 166 (23), 136 (16), 120 (42), 88 (20), 43 (17).

HRMS (FAB) calcd for C₂₇H₃₃N₃O₉+H₁ 544.2295, found 544.2310.

EXAMPLE 120 (COMPARISON) : 2-[4-[(2S)-2-[[[4-(aminosulfonyl)benzyl][3-(trifluoromethyl)benzyl]amino]carbonyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0282] ¹H NMR (DMSO-*d*₆) δ 7.83 (bt, *J* = 6 Hz, 1 H), 7.72 (d, *J* = 8 Hz, 2 H), 7.45-7.6 (m, 3 H), 7.3 (m, 4 H), 7.08 (d, *J* = 8.5 Hz, 2 H), 6.76 (d, *J* = 8.5 Hz, 2 H), 6.67 (bd, *J* = 8 Hz, 1 H), 5.27 (s, 1 H), 4.3-4.55 (m, 5 H), 2.7-3.1 (m, 4 H), 1.1-1.4 (m, 6 H), 0.83 (t, *J* = 7 Hz, 3 H);

MS (FAB) *m/z* (rel. intensity) 723 (MH⁺, 37), 724 (15), 723 (37), 345 (16), 170 (8.1), 159 (99), 107 (28), 106 (17), 91 (29), 88 (16), 43 (29).

HRMS (FAB) calcd for C₃₃H₃₇F₃N₄O₉S+H₁ 723.2311, found 723.2315.

Anal. Calcd for C₃₃H₃₇F₃N₄O₉S • 2.2 H₂O: C, 51.99; H, 5.48; N, 7.48.

Found: C, 51.99; H, 5.35; N, 7.34.

EXAMPLE 121 (COMPARISON) : 2-{4-[(2S)-2-[[[(3-carboxybenzyl)[3-(trifluoromethyl)benzyl]amino] carbonyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy}malonic acid

[0283] ¹H NMR (DMSO-*d*₆) δ 7.75 (m, 3 H), 7.2-7.6 (m, 6 H), 7.05 (d, *J* = 8 Hz, 2 H), 6.73 (d, *J* = 8 Hz, 2 H), 6.6 (bd, *J* = 8 Hz, 1 H), 5.23 (s, 1 H), 4.2-4.6 (m, 5 H), 3.0 (m, 2 H), 2.7-2.9 (m, 2 H), 1.1-1.4 (m, 6 H), 0.82 (t, *J* = 7 Hz, 3 H); MS (FAB) *m/z* (rel. intensity) 688 (MH⁺, 19), 689 (8), 688 (19), 601 (7), 310 (15), 160 (6), 159 (38), 136 (11), 135 (99), 91 (11), 43 (8).

HRMS (FAB) calcd for C₃₄H₃₆F₃N₃O₉+H₁ 688.2482, found 688.2489:

Anal. Calcd for C₃₄H₃₆F₃N₃O₉ • 1.05 H₂O: C, 57.80; H, 5.44; N, 5.95.

Found: C, 57.79; H, 5.21; N, 5.77.

EXAMPLE 122 (COMPARISON) : 2-{4-[(2S)-2-[[[benzyl[1-(carboxymethyl)-3-phenylpropyl] amino]carbonyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy}malonic acid

[0284] NMR analysis indicated a 1:1 mixture of diastereomers. ¹H NMR (DMSO-*d*₆) δ 7.7 (m, 1 H), 7.0-7.3 (m, 10 H), 6.93 (t, *J* = 8 Hz, 2 H), 6.7 (overlapping t, *J* = 8 Hz, 2 H), 5.19, 5.14 (two s, 1 H), 4.2-4.5 (m, 3 H), 2.2-3.1 (m, 9 H), 1.6 (m, 2 H), 1.1-1.4 (m, 6 H), 0.81, 0.80 (two t, *J* = 7 Hz, 3 H);

MS (FAB) *m/z* (rel. intensity) 662 (MH⁺, 27), 663 (11), 662 (27), 575 (9), 327 (6), 285 (8), 284 (34), 282 (6), 238 (8), 92 (9), 91 (99).

HRMS (FAB) calcd for C₃₆H₄₃N₃O₉+H₁ 662.3077, found 662.3080.

Anal. Calcd for C₃₆H₄₃N₃O₉ • 0.7 H₂O: C, 64.12; H, 6.64; N, 6.23.

Found: C, 64.12; H, 6.62; N, 6.11.

EXAMPLE 123 (COMPARISON) : 2-{4-[(2S)-2-[[[dibenzylamino]carbonyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy}malonic acid

[0285] ¹H NMR (DMSO-*d*₆) δ 7.71 (bt, *J* = 6 Hz, 1 H), 7.25 (m, 6 H), 7.05 (m, 6 H), 6.77 (d, *J* = 8.6 Hz, 2 H), 5.27 (s, 1 H), 4.38, 4.25 (ABq, *J* = 17 Hz, 4 H), 4.35 (m, 1 H), 2.7-3.1 (m, 4 H), 1.1-1.4 (m, 6 H), 0.84 (t, *J* = 7 Hz, 3 H).

MS (FAB) *m/z* (rel. intensity) 576 (MH⁺, 51), 577 (17), 576 (51), 575 (8), 489 (9), 241 (9), 177 (8), 92 (10), 91 (99), 63 (9), 43 (8).

HRMS (FAB) calcd for C₃₂H₃₇N₃O₇+H₁ 576.2709, found 576.2706.

Anal. Calcd for C₃₂H₃₇N₃O₇ • 0.85 H₂O: C, 65.04; H, 6.00; N, 7.11.

Found: C, 65.03; H, 6.51; N, 7.29.

EXAMPLE 124 (COMPARISON) : 2-{4-[(2S)-2-[[[4-(tert-butoxycarbonyl)-1-piperazinyl]carbonyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy}malonic acid

[0286] ¹H NMR (DMSO-*d*₆) δ 7.85 (bt, *J* = 6 Hz, 1 H), 7.15 (d, *J* = 8.6 Hz, 2 H), 6.79 (d, *J* = 8.6 Hz, 2 H), 5.28 (s, 1 H), 4.16 (m, 1 H), 3.20 (bs, 8 H), 2.55-3.1 (m, 4 H), 1.38 (s, 9 H), 1.1-1.4 (m, 6 H), 0.83 (t, *J* = 7 Hz, 3 H);

MS (FAB) *m/z* (rel. intensity) 565 (MH⁺, 60), 566 (18), 565 (60), 157 (41), 131 (23), 129 (16), 113 (45), 87 (61), 57 (99), 43 (19), 41 (16).

HRMS (FAB) calcd for C₂₇H₄₀N₄O₉+H₁ 565.2873, found 565.2896.

Anal. Calcd for C₂₇H₄₀N₄O₉ • 0.58 H₂O: C, 56.38; H, 7.21; N, 9.74.

Found: C, 56.39; H, 6.93; N, 9.43.

EXAMPLE 125 (COMPARISON) : 2-{4-[(2S)-2-[[[3-carboxyanilino]carbonyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy}malonic acid

[0287] Prepared by a procedure analogous to that described for Example 4 (Chart C).

¹H NMR (DMSO-*d*₆) δ 8.86 (s, 1 H), 8.03 (bt, *J* = 6 Hz, 1 H), 8.00 (s, 1 H), 7.45 (m, 2 H), 7.30 (t, *J* = 8 Hz, 1 H), 7.08 (d, *J* = 9 Hz, 2 H), 6.82 (d, *J* = 9 Hz, 2 H), 6.30 (2, *J* = 8 Hz, 1 H), 5.27 (s, 1 H), 4.36 (m, 1 H), 2.7-3.1 (m, 4 H), 1.1-1.4 (m, 6 H), 0.83 (t, *J* = 7 Hz, 3 H);

MS (FAB) *m/z* (rel. intensity) 516 (MH⁺, 99), 517 (29), 516 (99), 515 (15), 414 (20), 353 (19), 194 (14), 136 (12), 107 (13), 88 (22), 43 (13).

HRMS (FAB) calcd for C₂₅H₂₉N₃O₉+H₁ 516.1982, found 516.1965.

EXAMPLE 126: 2-(carboxymethoxy)-5-[(2S)-2-[[[(dibenzylamino)carbonyl]amino]-3-oxo-3-(pentylamino)propyl]benzoic acid

[0288] Prepared by a procedure analogous to that described for Example 10 (Chart D), using Q-4 as a starting material instead of B-4 as follows. Q-4 (54 mg, 0.11 mmole) was dissolved in 4 M HCl/dioxane (1 mL) and stirred at room temp. for 1 h. The solution was concentrated in vacuo, and the resulting residue was taken up in dry THF (2 mL). To the mixture was added triethylamine (47 μ L, 0.34 mmol), and the reaction was cooled to 0 °C before the addition of diphosgene (7 μ L, 0.06 mmol). The reaction was stirred at 0 °C for 15 minutes before the addition of dibenzylamine (29 μ L, 0.15 mmol). Stirring was continued at room temperature for 3 h, followed by the addition of 2.5 M aq LiOH (0.6 mL). The mixture was stirred vigorously for 2 h, at which point MS analysis indicated saponification was complete. The reaction was acidified with 1 M HCl (3 mL), saturated with solid NaCl, and extracted with ethyl acetate. Drying of the extracts over MgSO₄ and concentration in vacuo left a glass (67 mg). The crude material was sonicated with CH₂Cl₂ (20 mL) for 30 min, diluted with hexane (approx. 5 mL), and allowed to stand at room temperature for 1 h, affording a white powder (43 mg, 68% overall). ¹H NMR (DMSO-*d*₆) δ 7.82 (t, *J* = 7 Hz, 1 H), 7.62 (bs, 1 H), 7.15-7.3 (m, 7 H), 7.06 (d, *J* = 7 Hz, 4 H), 6.83 (d, *J* = 9 Hz, 1 H), 6.54 (d, *J* = 7 Hz, 1 H), 4.71 (s, 2 H), 4.35 (m, 1 H), 4.35, 4.25 (ABq, *J* = 16 Hz, 4 H), 2.7-3.1 (m, 4 H), 1.1-1.4 (m, 6 H), 0.84 (t, *J* = 7 Hz, 3 H); IR (mull) 3289 (b), 3088, 3064, 3030, 1735 (s), 1614 (s), 1585 (s), 1536 (s), 1496 (s), 1438 (s), 1340, 1301, 1247 (s), 1153, 700, cm⁻¹.

MS (FAB) *m/z* (rel. intensity) 576 (MH⁺, 40), 577 (14), 576 (40), 490 (4), 489 (14), 198 (7), 196 (6), 106 (6), 92 (8), 91 (99), 43 (5).

HRMS (FAB) calcd for C₃₂H₃₇N₃O₇+H₁ 576.2709, found 576.2704.

Anal. Calcd for C₃₂H₃₇N₃O₇ • 0.84 H₂O: C, 65.06; H, 6.60; N, 7.11.

Found: C, 65.06; H, 6.47; N, 7.24.

EXAMPLE 127 (COMPARISON): 2-[4-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-(1H-indol-3-yl)propanoyl]amino]-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0289] Prepared according to the general procedure described for BB-3 (Chart BB).

[0290] ¹H NMR (DMSO-*d*₆) δ 10.75 (s, 1 H), 7.84 (d, *J* = 8 Hz, 1 H), 7.80 (bt, 1 H), 7.51 (d, *J* = 7 Hz, 1 H), 7.29 (d, *J* = 8 Hz, 1 H), 6.95-7.15 (m, 4 H), 6.94 (t, *J* = 7 Hz, 1 H), 6.79 (d overlapping m, *J* = 8 Hz, 3 H), 5.22 (s, 1 H), 4.4 (m, 1 H), 4.12 (m, 1 H), 2.6-3.1 (m, 6 H), 1.1-1.4 (m, 6 H), 1.28 (s, 9 H), 0.82 (t, *J* = 7 Hz, 3 H);

MS (FAB) *m/z* (rel. intensity) 639 (MH⁺, 7), 194 (13), 186 (17), 170 (25), 159 (42), 131 (13), 130 (99), 88 (27), 57 (48), 43 (12), 41 (14).

HRMS (FAB) calcd for C₃₃H₄₂N₄O₉+H₁ 639.3030, found 639.3026.

Anal. Calcd for C₃₃H₄₂N₄O₉ • 0.72 H₂O: C, 60.82; H, 6.72; N, 8.60.

Found: C, 60.81; H, 6.70; N, 8.35.

EXAMPLE 128: (Chart R, R-3) 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[[[(2S)-3-phenyl-2-[[2-(4H-1,2,4-triazol-3-ylsulfanyl)acetyl]amino]propanoyl]amino]propyl]benzoic acid

[0291] A solution of R-1 (R = (S)-CH₂Ph) 0.5 g, 0.8 mmol) and 4N HCl/dioxane (20 mL) was stirred at room temp for 1.5 h. The reaction was concentrated to dryness in vacuo, and the resulting residue was taken up in DMF (50 mL). To the solution was added sequentially diisopropylethylamine (1.03 g, 0.8 mmol), 1,2,4-triazole-5-ylthioacetic acid (Drysdale et al, *J. Med. Chem.* **1992**, *35*, 2573)(159 mg, 1 mmol), HOBT monohydrate (135 mg, 1 mmol) and EDC hydrochloride (153 mg, 0.8 mmol). The mixture was stirred overnight at room temp. Solvent was removed in vacuo, and the residue was partitioned taken up in water and extracted with ethyl acetate. The organic extracts were dried over MgSO₄ and concentrated in vacuo. The residue was triturated with ether and dried again in vacuo, leaving the crude diester (190 mg) as a white solid. The diester was saponified by dissolving in DMF (10 mL) and adding 5.0 mL of 1.0 N aq NaOH. After stirring for 2 h, the reaction was neutralized by the addition of 1.0 N aq HCl (5.0 mL). The solvent was removed in vacuo, and the residue was taken up in water. The insoluble solid was filtered, air-dried and washed with ether, affording the title compound (136 mg) as an off-white solid after drying in vacuo. ¹H NMR (DMSO-*d*₆) δ 8.37 (bs, 1 H), 8.23 (d, *J* = 8 Hz, 1 H), 8.14 (d, *J* = 8 Hz, 1 H), 7.81 (bt, 1 H), 7.43 (bs, 1 H), 7.25 (bd, *J* = 7 Hz, 1 H), 7.1 (m, 5 H), 7.01 (d, *J* = 7 Hz, 1 H), 4.56 (bs, 2 H), 4.45 (m, 1 H), 4.37 (m, 1 H), 3.81, 3.72 (ABq, *J* = 15 Hz, 2 H), 2.8-3.1 (m, 4 H), 2.6-2.8 (m, 2 H), 1.1-1.4 (m, 6 H), 0.82 (t, *J* = 7 Hz, 3 H); MS (ESI-) for C₃₀H₃₆N₆O₈S *m/z* 639.2 (M-H)⁻.

HRMS (FAB) calcd for C₃₀H₃₆N₆O₈S +H₁ 641.2393, found 641.2388.

EXAMPLE 129: (Chart R, R-3) 2-(carboxymethoxy)-5-((2S)-3-oxo-3-(pentylamino)-2-(((2S)-3-phenyl-2-[[2-(5-sulfanyl-1H-1,2,3,4-tetraazol-1-yl)acetyl]amino]propanoyl)amino]propyl)benzoic acid

[0292] Prepared by a procedure analogous to that described for Example 128. ¹H NMR (DMSO-*d*₆) δ 8.55 (d, *J* = 7 Hz, 1 H), 8.22 (d, *J* = 8 Hz, 1 H), 7.77 (bt, *J* = 6 Hz, 1 H), 7.55 (d, *J* = 1 Hz, 1 H), 7.29 (dd, *J* = 7, 1 Hz, 1 H), 7.17 (m, 5 H), 6.90 (d, *J* = 7 Hz, 1 H), 4.94, 4.84 (ABq, *J* = 15 Hz, 2 H), 4.70 (bs, 2 H), 4.5 (m, 1 H), 4.37 (m, 1 H), 2.8-3.1 (m, 4 H), 2.65-2.8 (m, 2 H), 1.1-1.4 (m, 6 H), 0.82 (t, *J* = 7 Hz, 3 H); MS (ESI-) for C₂₉H₃₅N₇O₈S *m/z* 640.2 (M-H)⁻.

HRMS (FAB) calcd for C₂₉H₃₅N₇O₈S + H₁ 642.2346, found 642.2322.

EXAMPLE 130: (Chart R, R-3) 2-(carboxymethoxy)-5-((2S)-3-oxo-3-(pentylamino)-2-(((2S)-3-phenyl-2-[[2-(1H-1,2,3-triazol-5-yl)sulfanyl]acetyl] amino]propanoyl)amino]propyl)benzoic acid

[0293] Prepared by a procedure analogous to that described for Example 128. ¹H NMR (DMSO-*d*₆) δ 8.25 (d, *J* = 8 Hz, 1 H), 8.21 (d, *J* = 8 Hz, 1 H), 7.83 (bt, 1 H), 7.73 (bs, 1 H), 7.43 (bs, 1 H), 7.26 (dm, *J* = 7 Hz, 1 H), 7.15 (m, 5 H), 7.04 (d, *J* = 7 Hz, 1 H), 4.56 (bs, 2 H), 4.48 (m, 1 H), 4.38 (m, 1 H), 3.56 (bs, 2 H), 2.6-3.1 (m, 6 H), 1.1-1.4 (m, 6 H), 0.84 (t, *J* = 7 Hz, 3 H);

MS (FAB) *m/z* (rel. intensity) 641 (MH⁺, 18), 685 (17), 665 (11), 664 (20), 663 (47), 641 (18), 195 (19), 120 (99), 114 (17), 88 (25), 30 (17).

HRMS (FAB) calcd for C₃₀H₃₆N₆O₈S + H₁ 641.2393, found 641.2388.

EXAMPLE 131 (COMPARISON): 2-[4-((2S)-2-(((2-((carboxymethyl)amino)-2-oxoethyl)amino) carbonyl]amino)-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0294] Prepared by a procedure analogous to that described for Example 10 (Chart D). ¹H NMR (DMSO-*d*₆) δ 8.08 (bt, *J* = 7 Hz, 1 H), 7.82 (bt, *J* = 7 Hz, 1 H), 7.07 (d, *J* = 8.5 Hz, 2 H), 6.80 (d, *J* = 8.5 Hz, 2 H), 5.27 (s, 1 H), 4.22 (m, 1 H), 3.73 (s, 2 H), 3.65 and 3.55 (ABq, *J* = 15 Hz, 2 H), 2.6-3.1 (m, 4 H), 1.1-1.4 (m, 6 H), 0.82 (t, *J* = 7 Hz, 3 H); MS (FAB) *m/z* (rel. intensity) 511 (MH⁺, 99), 512 (25), 511 (99), 409 (17), 371 (30), 133 (25), 129 (16), 107 (22), 88 (30), 59 (27), 43 (22).

HRMS (FAB) calcd for C₂₂H₃₀N₄O₁₀ + H₁ 511.2040, found 511.2057.

EXAMPLE 132: (Chart T, Formula T-5) 2-[4-((2S)-2-(((2S)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropanoyl)amino)-3-oxo-3-(pentylamino)propyl]-2-(2H-1,2,3,4-tetraazol-5-yl)phenoxy]acetic acid

[0295] PREPARATION OF T-1: To a solution of Q-2 (5.94 g, 12.47 mmol) in DMF (anhydrous, 30 mL) in a Heck vial was added Pd(PPh₃)₄ (0.58 g, 0.50 mmol) and zinc cyanide (1.61 g, 13.72 mmol). The vial was flushed with nitrogen, tightly sealed and stirred at 80 °C for 16h. After cooling to room temperature, the mixture was partitioned between EtOAc (50 mL) and 2 M aqueous ammonium hydroxide (50 mL). The organic layer was dried (Na₂SO₄), and concentrated. The residue was purified by column chromatography (SiO₂, EtOAc/n-hexane 1:1) to afford 1.76 g (38%) of T-1 as a white solid: ¹H NMR 400 MHz (MeOH) δ 0.91 (t, 3H, *J* = 7.3, 14.4), 1.18-1.48 (m, 6H), 1.39 (s, 9H), 2.77 (dd, 1H, *J* = 8.6, 13.7), 2.98 (dd, 1H, *J* = 6.45, 13.7), 3.08-3.21 (m, 2H), 4.20 (m, 1H), 6.88 (d, 1H, *J* = 8.5), 7.34 (dd, 1H, *J* = 2.2, 8.5), 7.38 (d, 1H, 2.2).

[0296] PREPARATION OF T-2: Prepared from T-1 (0.61 g, 1.63 mmol) by the general method described for U-2 and U-3, which afforded 0.69 g (94%) of the title compound as a white solid. ¹H NMR 400 MHz (CDCl₃) δ 0.90 (t, 3H, *J* = 7.1, 14.4), 1.22-1.45 (m, 6H), 1.38 (s, 9H), 2.79 (dd, 1H, *J* = 8.8, 13.7), 3.01 (m, 1H), 3.11-3.18 (m, 2H), 3.77 (s, 3H), 4.87 (s, 2H), 7.00 (d, 1H, *J* = 8.7), 7.46 (dd, 1H, *J* = 2.2, 8.7), 7.51 (d, 1H, *J* = 2.2).

[0297] PREPARATION OF T-3: Prepared from T-2 (0.55 mg, 1.23 mmol) by the general method described for U-4 and U-5, which afforded 0.61 g (84%) of the title compound as a white solid. Mp = 121.4-123.0 °C. ¹H NMR 400 MHz (MeOH) δ 0.90 (t, 3H, *J* = 7.2, 14.5), 1.21 (m, 2H), 1.29 (m, 2H), 1.37 (s, 9H), 1.40 (m, 2H), 2.75 (dd, 1H, *J* = 9.2, 13.5), 2.91 (dd, 1H, *J* = 7.7, 13.5), 2.97-3.06 (m, 3H), 3.12 (m, 1H), 3.75 (s, 3H), 4.24 (dd, 1H, *J* = 5.3, 9.2), 4.51 (app t, 1H, *J* = 7.1, 14.1), 4.87 (s, 2H), 6.98 (d, 1H, *J* = 8.6), 7.14-7.27 (m, 5H), 7.44 (dd, 1H, *J* = 1.9, 8.6), 7.49 (d, 1H, *J* = 1.9).

[0298] PREPARATION OF T-4: To a suspension of T-3 (0.45 g, 0.75 mmol) in toluene in a Heck vial was added trimethylsilyl azide (0.30 mL, 2.25 mmol) and dibutyltin oxide (19 mg, 0.075 mmol). The mixture was flushed with nitrogen; tightly sealed and stirred at 110 °C for 16h. The reaction mixture was cooled to room temperature and the volatiles were evaporated in vacuo. The residue was purified by column chromatography (SiO₂, EtOAc) which afforded 90 mg (19%) of T-4 as a white solid. Mp = 189.5-192.8 °C. ¹H NMR 400 MHz (MeOH) δ 0.85 (t, 3H, *J* = 6.9, 14.1), 1.16-1.41 (m, 6H), 1.33 (s, 9H), 2.69 (dd, 1H, *J* = 9.4, 13.7), 2.95-3.17 (m, 5H), 3.79 (s, 3H), 4.22 (dd, 1H, *J* = 4.9, 9.4), 4.58 (m, 1H), 4.99 (s, 2H), 7.09-7.24 (m, 6H), 7.42 (d, 1H, *J* = 1.7, 6.9), 8.09 (d, 1H, *J* = 1.7).

[0299] PREPARATION OF T-5: Prepared from T-4 (19 mg, 0.029 mmol) by the general method described for U-1 0

and U-11, which afforded 16 mg (90%) of the title compound as a white solid. ¹H NMR 400 MHz (MeOH) δ 0.85 (t, 3H, J = 7.0, 14.2), 1.18 (m, 2H), 1.24 (m, 2H), 1.33 (s, 9H), 1.38 (m, 2H), 2.70 (dd, 1H, J = 9.4, 13.8), 2.96-3.08 (m, 3H), 3.14 (m, 2H), 4.22 (dd, 1H, J = 5.1, 9.4), 4.58 (m, 1H), 4.95 (s, 2H), 7.12-7.29 (m, 6H), 7.44 (dd, 1H, J = 1.9, 8.5), 7.86 (br m, 0.5H), 8.01 (br m, 1H), 8.10 (d, 1H, J = 1.9); ¹³C NMR (MeOH) δ 14.3, 23.4, 28.7, 30.0, 30.2, 38.2, 39.0, 40.6, 55.8, 67.0, 80.9, 114.0, 114.9, 127.8, 129.5, 130.3, 131.5, 132.8, 135.4, 138.5, 155.9, 157.9, 172.5, 173.1, 174.1. MS (ESI) 622 (M-H). HRMS (EI) calcd for C₃₁H₄₁N₇O₇ 623.3068, found 623.3071. Anal. Calcd for C₃₁H₄₁N₇O₇: C, 59.70; H, 6.63; N, 15.72. Found: C, 59.5; H, 6.7; N, 14.0.

EXAMPLES 133-135 (Chart U)

[0300] PREPARATION OF U-1: Triethylamine (1.71 mL, 12.5 mmol) and benzyl alcohol (6.45 mL, 62 mmol) is added to a stirring suspension of Q-2 (2.97 g, 6.23 mmol) palladium(II)acetate (42 mg, 0.19 mmol) and 1,1'-bis(diphenylphosphino)ferrocene (DPPF, 207 mg, 0.37 mmol) in DMF (15 mL). The mixture is saturated with CO (1 atm) and stirred at 70 °C for 16 h. The mixture is allowed to reach room temperature and extracted with EtOAc (40 mL), the organic layer is washed with 10% aqueous HCl (20 mL) and brine (20 mL), dried (Na₂SO₄) and concentrated. The residue is purified by column chromatography (SiO₂, EtOAc/n-hexane 1:2), which furnished 5.7 g of a yellow oil. This crude material still contains some benzyl alcohol. Crystallization in EtOAc/n-hexane gives 0.82 g (27%) of pure U-1 as a white solid. ¹H NMR 400 MHz (CDCl₃) δ 0.84 (t, 3H, J = 7.1, 14.4), 1.13 (m, 2H), 1.23 (m, 2H), 1.35 (m, 2H), 1.39 (s, 9H), 2.95 (d, 2H, J = 6.7), 3.11 (m, 2H), 4.18 (m, 1H), 5.07 (br m, 1H), 5.37 (d, 2H, J = 1.7), 5.77 (br m, 1H), 6.91 (d, 1H, J = 8.5), 7.30 (dd, 1H, J = 2.2, 8.5), 7.35-7.47 (m, 5H), 7.69 (d, 1H, J = 2.2), 10.69.

[0301] PREPARATION OF U-2: Methyl bromoacetate (0.35 mL, 3.75 mmol) and freshly grounded K₂CO₃ (0.52 g, 3.75 mmol) is added to a stirring solution of U-1 (0.61 g, 1.25 mmol) in acetone (15 mL). The mixture is stirred at 50 °C over night. After cooling to ambient temperature, H₂O (10 mL) is added and the mixture is extracted with EtOAc (10 mL). The organic layer is dried (Na₂SO₄), and concentrated. The residue is purified by column chromatography (SiO₂, EtOAc/ n-hexane 1:1) which furnished 0.46 g (66%) of U-2 as a white solid. ¹H NMR 400 MHz (MeOH) δ 0.87 (t, 3H, J = 7.1, 14.3), 1.17-1.40 (m, 6H), 1.35 (s, 9H), 2.79 (dd, 1H, J = 8.4, 13.6), 2.98-3.14 (m, 3H), 3.74 (s, 3H), 4.19 (m, 1H), 4.75 (s, 2H), 5.33 (s, 2H), 6.95 (d, 1H, J = 8.5), 7.32-7.39 (m, 5H), 7.46 (dd, 1H, J = 1.9, 8.5), 7.67 (d, 1H, J = 1.9).

[0302] PREPARATION OF U-3: From Q-3 as described for the preparation of U-2. ¹H NMR 400 MHz (MeOH) δ 0.89 (t, 3H, J = 6.8, 13.9), 1.22 (m, 2H), 1.29 (m, 2H), 1.37 (s, 9H), 1.39 (m, 2H), 2.81 (dd, 1H, J = 8.5, 13.3), 3.00 (dd, 1H, J = 7.4, 13.3), 3.07-3.15 (m, 2H), 3.82 (s, 3H), 4.20 (m, 1H), 4.79 (s, 2H), 5.21 (s, 2H), 6.91 (d, 1H, J = 8.5), 7.31-7.35 (m, 6H), 7.66 (s, 1H).

[0303] PREPARATION OF U-4: Trifluoroacetic acid (0.90 mL) is carefully added to a stirring solution of U-2 (0.44 mg, 0.78 mmol) in CH₂Cl₂ at 0 °C. The mixture is stirred for 4h allowing the solution to warm to ambient temperature. The volatiles are removed by vacuo and the residue is partitioned between EtOAc (15 mL) and saturated aqueous NaHCO₃ (2x10 mL). The organic layer is dried (Na₂SO₄) and concentrated to give 0.35 g (98%) of the crude amine as a yellowish oil. The amine is dissolved in CH₂Cl₂ (7 mL) and cooled with ice. Boc-(L)-phenylalanine (0.20 g, 0.77 mmol), 1-hydroxybenzotriazole (0.10 g, 0.77 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC, 0.15 g, 0.77 mmol) is added to the solution, which is then stirred at room temperature over night. The reaction mixture is diluted with CH₂Cl₂ (5 mL) and washed with saturated aqueous NaHCO₃ (5 mL), brine (5 mL) and 10% aqueous HCl. The organic layer is dried (Na₂SO₄) and concentrated. The residue is purified by column chromatography (SiO₂, EtOAc) which gave 0.46 g (86%) of U-4 as a white solid. ¹H NMR 400 MHz (MeOH) δ 0.86 (t, 3H, J = 7.2, 14.5), 1.18 (m, 2H), 1.24 (m, 2H), 1.34 (s, 9H), 1.38 (m, 2H), 2.72 (dd, 1H, J = 9.4, 13.8), 2.89-3.11 (m, 3H), 3.71 (s, 3H), 4.22 (m, 1H), 4.49 (m, 1H), 4.74 (s, 2H), 5.31 (s, 2H), 6.94 (d, 1H, J = 8.6), 7.14-7.46 (m, 6H), 7.65 (d, 1H, J = 1.7).

[0304] PREPARATION OF U-5: From U-3 as described for U-4. ¹H NMR 400 MHz (MeOH) δ 0.88 (t, 3H, J = 7.1, 14.4), 1.18 (m, 2H), 1.28 (m, 2H), 1.34 (s, 9H), 1.37 (m, 2H), 2.74 (dd, 1H, J = 9.4, 13.8), 2.93 (dd, 1H, J = 7.5, 13.8), 2.97-3.05 (m, 3H), 3.12 (m, 1H), 3.82 (s, 3H), 4.23 (m, 1H), 4.50 (m, 1H), 4.79 (s, 2H), 5.18 (s, 2H), 6.69 (br d, 0.6H), 6.91 (d, 1H, J = 8.6).

EXAMPLE 133: (Chart U, Formula U-6) 5-[(2S)-2-(((2S)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropanoyl)amino)-3-oxo-3-(pentylamino)propyl]-2-(2-methoxy-2-oxoethoxy)benzoic acid

[0305] A mixture of U-4 (0.41 g, 0.58 mmol) and 10% Pd/C (80 mg) in methanol (25 mL) is hydrogenated at atmospheric pressure for 2h. The mixture is filtered through Celite and solvent removed in vacuo to afford 0.34g (96%) of U-6 as a white solid. ¹H NMR 400 MHz (MeOH) δ 0.89 (t, 3H, J = 7.1, 14.4), 1.19-1.42 (m, 6H), 1.36 (s, 9H), 2.73 (dd, 1H, J = 9.5, 13.8), 2.93-3.07 (m, 3H), 3.14 (m, 1H), 3.75 (s, 2H), 4.23 (m, 1H), 4.51 (br m, 1H), 4.81 (s, 2H), 6.96 (d, 1H, J = 8.5), 7.17-7.27 (m, 5H), 7.35 (dd, 1H, J = 1.8, 8.5), 7.72 (d, 1H, J = 1.8), 7.84 (br m, 0.5H), 7.98 (br m, 0.2H); ¹³C NMR (MeOH) δ 14.6, 23.7, 28.9, 30.2, 30.2, 30.4, 38.3, 39.3, 40.8, 53.0, 56.1, 58.0, 67.4, 81.1, 115.6, 122.7, 128.0, 129.7, 130.6, 132.0, 134.2, 135.9, 138.8, 157.9, 158.2, 169.7, 171.2, 172.7, 172.8, 174.3. MS (ESI) 612 (M-H).

[0306] Anal. Calcd for $C_{32}H_{43}N_3O_9$: C, 62.63; H, 7.06; N, 6.85. Found: C, 62.0; H, 7.0; N, 6.8.

[0307] PREPARATION OF U-7: From U-5 as described for U-6. 1H NMR 400 MHz (MeOH) δ 0.89 (t, 3H, $J = 7.1$, 14.4), 1.21 (m, 2H), 1.29 (m, 2H), 1.35 (s, 9H), 1.37 (m, 2H), 2.75 (dd, 1H, $J = 9.3$, 13.8), 2.94 (dd, 1H, $J = 7.4$, 13.8), 2.98-3.05 (m, 3H), 3.13 (m, 1H), 3.86 (s, 3H), 4.24 (m, 1H), 4.50 (m, 1H), 4.69 (s, 2H), 6.95 (d, 1H, $J = 8.6$), 7.17-7.27 (m, 5H), 7.35 (dd, 1H, $J = 1.5$, 8.6), 7.64 (d, 1H, $J = 1.5$), 7.83 (br m, 1H), 7.98 (br d, 0.6H).

[0308] PREPARATION OF U-8: To a solution of U-6 (116 mg, 0.19 mmol) in a mixture of THF (2.5 mL) and DMF (0.2 mL) was added 1,1'-carbonyldiimidazole (CDI, 61 mg, 0.38 mmol), and the mixture was refluxed at 80 °C for 1h. The mixture was cooled to ambient temperature and hydroxylamine hydrochloride (39 mg, 0.57 mmol) was added. The reaction mixture was refluxed at 80 °C for 4h. After cooling to room temperature the mixture was partitioned between EtOAc (3 mL) and 3 M aqueous HCl (3 mL), the organic layer was dried (Na_2SO_4) and concentrated. The residue was purified by column chromatography (SiO_2 , 5% MeOH in CH_2Cl_2) to afford 76 mg of a red/brownish solid. This material could be further purified by crystallization in EtOAc to give 34 mg (28%) of U-8 as a white solid. 1H NMR 400 MHz (MeOH) δ 0.89 (t, 3H, $J = 7.1$, 14.4), 1.20 (m, 2H), 1.29 (m, 2H), 1.35 (s, 9H), 1.38 (m, 2H), 2.70 (dd, 1H, $J = 9.4$, 13.3), 2.91-3.17 (m, 5H), 3.81 (s, 3H), 4.21 (m, 1H), 4.54 (m, 1H), 4.86 (s, 2H, obscured by solvent peak), 6.98 (d, 1H, $J = 8.1$), 7.16-7.27 (m, 5H), 7.34 (br d, 1H, $J = 8.1$), 7.84 (br s, 1H).

[0309] PREPARATION OF U-9: From U-7 as described for U-8. 1H NMR 400 MHz (MeOH) δ 0.88 (t, 3H, $J = 7.1$, 14.5), 1.19 (m, 2H), 1.28 (m, 2H), 1.35 (s, 9H), 1.37 (m, 2H), 2.77 (dd, 1H, $J = 9.2$, 13.7), 2.93 (dd, 1H, $J = 7.5$), 2.98-3.06 (m, 3H), 3.13 (m, 1H), 3.90 (s, 3H), 4.22 (dd, 1H, $J = 5.3$, 9.1), 4.50 (app t, 1H), 4.66 (s, 2H), 7.06 (d, 1H, $J = 8.5$), 7.17-7.27 (m, 5H), 7.43 (dd, 1H, $J = 1.8$, 8.5), 7.75 (d, 1H, 1.8).

EXAMPLE 134 (COMPARISON): (Chart U, Formula U-10) 2-[4-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]-2-[(hydroxyamino)carbonyl]phenoxy}acetic acid

[0310] To a solution of U-8 (10 mg, 0.0165 mmol) in THF (200 μ L) was added a 2.5 M aqueous solution of LiOH (9.9 μ L, 0.0248 mmol). The mixture was stirred at ambient temperature for 2h. The reaction mixture was acidified with 3 M HCl and extracted with EtOAc (2 mL). The organic layer was dried (Na_2SO_4), and concentrated to afford 8.0 mg (79%) of U-10 as a white solid. 1H NMR 400 MHz (MeOH) δ 0.88 (t, 3H, $J = 7.1$, 14.5), 1.19 (m, 2H), 1.29 (m, 2H), 1.35 (s, 9H), 1.38 (m, 2H), 2.71 (dd, 1H, $J = 9.5$, 13.5), 2.94 (m, 1H), 2.98-3.07 (m, 3H), 3.12 (m, 2H), 4.22 (dd, 1H, $J = 5.1$, 9.5), 4.54 (m, 1H), 4.81 (s, 2H), 6.99 (d, 1H, $J = 8.5$), 7.17-7.27 (m, 5H), 7.35 (dd, 1H, $J = 1.8$, 8.5), 7.85 (d, 1H, $J = 1.8$); ^{13}C NMR (MeOH) δ 14.6, 23.7, 28.9, 30.2, 30.4, 38.4, 39.2, 40.7, 56.1, 58.0, 67.3, 81.1, 114.8, 121.9, 128.0, 129.7, 130.6, 132.4, 133.4, 135.4, 138.8, 156.3, 158.1, 165.7, 172.4, 172.8, 174.3. MS (ESI) 613 (M-H). Anal. Calcd for $C_{31}H_{42}N_4O_9 \cdot \frac{1}{2} H_2O$: C, 59.70; H, 6.95; N, 8.98. Found: C, 59.5; H, 7.0; N, 8.8.

EXAMPLE 135 (COMPARISON): (Chart U, Formula U-11) 5-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]-2-[2-(hydroxyamino)-2-oxoethoxy]benzoic acid

[0311] Prepared from U-9 (10 mg, 0.016 mmol) by the above method which afforded 8.1 mg (83%) of the title compound as a white solid. 1H NMR 400 MHz (MeOH) δ 0.88 (t, 3H, $J = 7.1$, 14.5), 1.19 (m, 2H), 1.29 (m, 2H), 1.35 (s, 9H), 1.38 (m, 2H), 2.78 (dd, 1H), 2.95 (dd, 1H), 3.14 (m, 1H), 4.23 (dd, 1H, $J = 5.2$, 9.2), 4.51 (m, 1H), 4.68 (s, 2H), 7.04 (d, 1H, $J = 8.5$), 7.17-7.27 (m, 5H), 7.42 (dd, 1H, $J = 2.1$, 8.5), 7.78 (d, 1H, 2.1); ^{13}C NMR (MeOH) δ 14.3, 23.4, 286, 29.9, 30.1, 38.1, 39.0, 40.5, 55.8, 57.7, 68.9, 80.9, 115.8, 121.2, 127.7, 129.4, 130.3, 131.9, 134.2, 136.4, 138.4, 158.2, 167.6, 169.0, 172.4, 172.5, 174.0. MS (ESI) 613 (M-H). Anal. Calcd for $C_{31}H_{42}N_4O_9 \cdot \frac{1}{2} H_2O$: C, 59.70; H, 6.95; N, 8.98. Found: C, 59.6; H, 7.2; N, 8.8.

EXAMPLE 136 (COMPARISON): (Chart V, Formula V-6) 2-[4-[(2S,3E,Z)-2-[(3-carboxypropanoyl)amino]-3-nonenyl]phenoxy]malonic acid

[0312] PREPARATION OF V-2: Diisobutylaluminum hydride (DIBAL-H, 20 wt.% in toluene, 72.2 mL, 101.6 mmol) was added dropwise, during 10 min, to a stirred solution of N-Boc-L-tyrosine methyl ester (V-1, 6.0 g, 20.3 mmol) in diethylether (dried over molecular sieves, 120 mL) kept at -78 °C, under nitrogen atmosphere. After being stirred at -78 °C for 1 h, the reaction was quenched with MeOH (20 mL) and the mixture was poured into a saturated aqueous solution of potassium sodium tartrate (Rochelle salt, 500 mL). The mixture was allowed to warm to ambient temperature and diethylether (100 mL) was added. The ethereal layer was separated, dried ($MgSO_4$) and concentrated. The crude product was passed through a short pad of silica gel, eluting with EtOAc/n-hexane 1:1, which furnished 4.9 g (91%) of V-2 as a slightly yellow oil. 1H NMR 400 MHz ($CDCl_3$) δ 1.44 (s, 9H), 3.02 (d, 2H, $J = 6.5$), 4.41 (dd, 1H, $J = 6.5$, 13.3), 5.14 (m, 1H), 6.75 (d, 2H, $J = 8.1$), 7.00 (d, 2H, $J = 8.1$), 9.61 (s, 1H).

[0313] PREPARATION OF V-3: Potassium-*tert*-butoxide (12.5 g, 111.2 mmol) was added to solution of N-hexyltriphenylphosphonium bromide (5.9 g, 22.2 mmol) in THF (dried over molecular sieves, 60 mL) at -5 °C (ice/acetone

bath), under nitrogen atmosphere. After 10 min stirring, the aldehyde V-2 (5.9 g, 22.2 mmol) dissolved in THF (30 mL) was added dropwise. The mixture was stirred at 0 °C for 10 min, and then at ambient temperature for an additional 25 min. The reaction was quenched with crushed ice and the pH was adjusted to 3-4 with 10 % aqueous HCl. The mixture was extracted with EtOAc (50 mL), washed with brine (2 x 50 mL), dried (MgSO₄), and concentrated. The residue was purified by flash chromatography (SiO₂, EtOAc/n-hexane 1:1.5), which furnished 4.31 g (58%) of V-3 as a yellow oil. ¹H NMR 400 MHz (CDCl₃) δ 0.85 (t, 3H, J = 6.8, 14.0), 1.19 - 1.28 (m, 6H), 1.43 (s, 9H), 1.85 - 1.97 (m, 2H), 2.64 (m, 1H), 2.82 (br m, 2H), 4.45 (br s, 1H), 5.18 (dd, 1H, J = 8.9, 10.7), 5.42 (ddd, 1H, J = 7.4, 10.7, 14.8), 6.73 (d, 2H, J = 8.5), 7.00 (d, 2H, J = 8.5).

[0314] PREPARATION OF V-4: Dibenzyl bromomalonate (4.6 g, 13.7 mmol) and potassium carbonate (2.71 g, 19.6 mmol) were added to a solution of V-3 (3.27 g, 9.81 mmol) in acetone (40 mL). The mixture was stirred at ambient temperature for 16 h. The reaction was quenched with H₂O (100 mL) and the mixture was extracted with EtOAc (2 x 50 mL). The organic layer was dried (MgSO₄) and concentrated. The residue was purified by flash chromatography (SiO₂, EtOAc/n-hexane 1:5) which furnished 5.86 g (97%) of V-4 as a yellow oil. ¹H NMR 400 MHz (CDCl₃) δ 0.88 (t, 3H), 1.15 - 1.29 (m, 6H), 1.42 (s, 9H), 1.80 - 2.00 (m, 2H), 1.63 (m, 1H), 1.72 (m, 1H), 3.59 (s, 1H), 4.52 (br s, 1H), 5.02 - 5.42 (m, 7H), 6.7 - 7.3 (m, 14H).

[0315] PREPARATION OF V-5: Trifluoroacetic acid (2.4 mL, 31.1 mmol) was carefully added to a stirring solution of V-4 (1.74 g, 2.83 mmol) in CH₂Cl₂ (20 mL) at 0 °C. The mixture was stirred for 2 h allowing the solution to warm to ambient temperature. The volatiles were removed *in vacuo* and the residue was partitioned between diethylether (20 mL) and saturated aqueous NaHCO₃ (2 x 10 mL). The organic layer was dried (MgSO₄), and concentrated to dryness to afford 1.49 g (> 100%) of the crude amine as a brown/yellow oil. The amine was dissolved in CH₂Cl₂ (20 mL) and cooled with ice to 0 °C. Benzylhydrogen succinate (0.59 g, 2.83 mmol) and N'-((3-dimethylaminopropyl)-N-ethylcarbodiimide hydrochloride (EDC, 0.54 g, 2.83 mmol) was added and the mixture was stirred for 16 h allowing the solution to warm to ambient temperature. The mixture was diluted with CH₂Cl₂ (10 mL) and the organic phase was washed with 10 % aqueous HCl (2 x 20 mL), dried (MgSO₄), and concentrated. TLC indicated that the product was a mixture of 4 compounds. By repeated flash chromatography, (SiO₂, EtOAc/n-hexane 1:3), 0.57 g (28%) of V-5 as a colorless oil could be isolated. ¹H NMR 400 MHz (CDCl₃) δ 0.83 (t, 3H, J = 6.8, 14.0), 1.14 - 1.26 (m, 8H), 1.77 (m, 1H), 1.94 (m, 1H), 2.41 (t, 2H), 2.69 (m, 2H), 2.83 (dd, 1H, J = 5.1, 13.4), 4.86 (br m, 1H), 5.11 (s, 2H), 5.21 (s, 4H), 5.24 (s, 1H), 5.43 (m, 1H), 5.58 (br d, 1H), 6.82 (d, 2H, J = 8.5), 7.05 (d, 2H, J = 8.5), 7.25 - 7.34 (m, 15H).

[0316] PREPARATION OF V-6: A solution of V-5 (172 mg, 0.25 mmol) and 2.5 M aqueous LiOH (609 µL, 1.52 mmol) in THF (4 mL) was stirred at ambient temperature for 16 h. The reaction mixture was partitioned between EtOAc (5 mL) and saturated aqueous NaHCO₃ (5 mL). The water layer was acidified with 10% aqueous HCl and extracted with EtOAc (4 x 2 mL). The organic layer was dried (MgSO₄) and concentrated to dryness which furnished 96 mg (87%) of V-6 as a white solid. ¹H NMR 400 MHz (CDCl₃) δ 0.85 (t, 3H, J = 6.8, 14.0), 1.19 - 1.28 (m, 6H), 1.43 (s, 9H), 1.85 - 1.97 (m, 2H), 2.64 (m, 1H), 2.82 (br m, 2H), 4.54 (br s, 1H), 5.18 (dd, 1H, J = 8.9, 10.7), 5.42 (ddd, 1H, J = 7.4, 10.7, 14.8), 6.73 (d, 2H, J = 8.5), 7.00 (d, 2H, J = 8.5); ¹³C NMR (CDCl₃) δ 14.0, 22.5, 27.7, 28.4, 29.0, 31.4, 41.4, 60.5, 80.3, 115.2, 128.9, 129.1, 130.6, 133.0, 154.7, 173.2. MS (ESI) 434 (M - H). Anal. Calcd. for C₂₂H₂₉O₈N • ¼ H₂O: C, 60.06; H, 6.76; N, 3.18; O, 30.0. Found: C, 60.1; H, 6.7; N, 3.0; O, 29.8.

EXAMPLE 137 (COMPARISON): (Chart W, W-6) 2-[4-((2S)-2-(((2S)-2-((tert-butoxycarbonyl)amino)-3-phenylpropanoyl)amino)-3-oxo-3-(pentylamino)propyl)] (carboxymethyl)anilino]acetic acid

[0317] PREPARATION OF W-2: To a stirring solution of Boc-Phe (4-NO₂)-OH (1.0 g, 3.22 mmol) and amyl amine (0.0.280 g, 3.22 mmol) in CH₂Cl₂ at 0 °C was added EDC (0.617 g, 3.22 mmol) and the resulting solution stirred overnight allowing the solution to warm to ambient temperature. CH₂Cl₂ (100 mL) was added and the solution washed with 10% HCl/H₂O (3 x 100 mL). The organic layer was further washed with saturated aqueous NaHCO₃ (3 x 100 mL) dried over MgSO₄ and solvent removed under reduced pressure to afford 0.856 g title compound as a white solid. m.p. 151-152 °C. ¹H NMR δ 0.86 (t, 3H), 1.22 (m, 6H), 1.40 (s, 9H), 3.16 (m, 4H), 4.31 (q, 1H), 5.02 (brs, 1H), 5.87 (brs, 1H), 7.38 (d, 2H), 8.14 (d, 2H).

[0318] PREPARATION OF W-3: A mixture of W-2 (3.43 g, 9.04 mmol) and 10% Pd/C (0.9 g) in abs. EtOH (125 mL) was hydrogenated at atmospheric pressure and room temperature for 4 h. The reaction mixture was filtered over diatomaceous earth and evaporated *in vacuo* leaving a off-white solid, that was purified on a silica gel flash-column eluting with ethyl acetate/n-hexane (2:1 v/v). A pinkish solid (3.01 g, 95%) was isolated after pooling and evaporating pure fractions. ¹H NMR δ 0.86 (t, 3H), 1.17-1.41 (m, 15H), 2.84-2.97 (m, 2H), 3.06-3.19 (m, 2H), 3.69 (s, 2H), 4.22 (m, 1H), 5.28 (br s, NH), 6.08 (t, NH), 6.59 (d, 2H), 6.96 (d, 2H).

[0319] PREPARATION OF W-4: A magnetically stirred mixture of W-3 (410 mg, 1.17 mmol), methyl bromoacetate (300 µL, 2.2 eq.), K₂CO₃ (500 mg) and KI (100 mg) in acetonitrile (4 mL) was heated at 60 °C for 15 h and then for 6 h at ambient temperature under N₂-atmosphere. The reaction mixture was treated with H₂O (20 mL), extracted with ethyl acetate (3 x 15 mL) and washed with brine (30 mL). The combined organic layers were dried over Na₂SO₄,

filtered and reduced to dryness giving 0.83 g of a yellow oil. Purification with flash-chromatography on silica gel eluting with ethyl acetate/n-hexane (1:1 v/v) gave 429 mg (74 %) of a yellow waxy solid. ^1H NMR δ 0.87 (t, 3H), 1.16-1.42 (m, 15H), 2.85-3.04 (m, 2H), 3.08-3.22 (m, 2H), 3.75 (s, 6H), 4.13 (s, 4H), 4.25 (m, 1H), 5.05 (br s, NH), 5.76 (br s, NH), 7.04 (d, 2H), 7.53 (d, 2H).

[0320] PREPARATION OF W-5: At 0 °C, TFA (0.5 mL) was added dropwise to a solution of W-4 (200 mg, 0.41 mmol) in DCM (4 mL). The resulting solution was stirred for 4 h at room temperature. Evaporation in vacuo gave a red oil that was taken up in methanol and heated shortly with activated carbon. Filtration over diatomaceous earth and removal of the solvent gave 246 mg (97%) the title compound di-TFA salt as a yellow oil. This oil was dissolved in DCM (3 mL) in stirred at 0 °C. Then, HOBT (54 mg, 1 eq), Boc-L-Phe (106 mg, 1 eq), EDC (77 mg, 1 eq) and TEA (166 μL , 3 eq) were added, and this mixture was stirred for 16 h at room temperature. Ethyl acetate (30 mL) was added and washed with 2% aqueous HCl (20 mL) and saturated aqueous NaHCO_3 (20 mL). The organic layer was dried (Na_2SO_4) and reduced to dryness giving 210 mg of a yellow oil. Purification with flash-chromatography on silica gel eluting with ethyl acetate/n-hexane (1:1 v/v) gave 132 mg (50 %) after two steps of a colorless oil. A small fraction was recrystallized from toluene giving a white solid. ^1H NMR δ 0.87 (t, 3H), 1.16-1.42 (m, 15H), 2.74-2.79 (m, 1H), 2.97-3.14 (m, 5H), 3.74 (s, 6H), 4.11 (s, 4H), 4.28 (m, 1H), 4.51 (m, 1H), 4.91 (br s, NH), 5.99 (br s, NH), 6.41 (m, NH), 6.48 (d, 2H), 6.94 (d, 2H), 7.16 (d, 2H), 7.23-7.32 (m, 3H).

[0321] PREPARATION OF W-6: To a stirred solution of W-5 (88 mg, 137 μmol) in THF (2 mL), aqueous LiOH (2.5 M, 400 μL , 7.3 eq) was added. After 3 h the reaction mixture was acidified with 10% aqueous HCl until pH 5. The milky suspension was extracted with warm ethyl acetate (3×15 mL). The combined organic layers were dried (Na_2SO_4) and evaporated in vacuo leaving a white solid (64 mg, 76%). A small portion was recrystallized from acetonitrile giving white flakes. ^1H NMR (CD_3OD) δ 0.91 (t, 3H), 1.20-1.42 (m, 15H), 2.01 (m, 1H), 2.79 (dd, 2H), 2.90-3.14 (m, 3H), 4.17 (s, 4H), 4.26 (dd, 1H), 4.46 (m, 1H), 6.53 (d, 2H), 7.05 (d, 2H), 7.20-7.30 (m, 5H); MS (ionspray, $[\text{M}-\text{H}]^+$) m/z 611.0; Anal. Calcd. (found) for $\text{C}_{32}\text{H}_{46}\text{N}_4\text{O}_9 \cdot \text{H}_2\text{O}$: C 60.9 (60.8) % H 7.4 (7.1) % N 8.9 (9.0) %.

EXAMPLE 138 (COMPARISON): (Chart X, Formula X-7) 5-[(2S)-2-(((2S)-2-[(tert-butoxycarbonyl) amino]-3-phenylpropanoyl)amino)-3-hydroxypropyl]-2-(carboxymethoxy)benzoic acid

[0322] PREPARATION OF X-1: 3-Iodo-L-tyrosine (5.0 g, 16.3 mmol) was suspended in benzyl alcohol (100 mL) and at 0 °C, thionyl chloride (20 mL) was added dropwise over a 20-min period. The temperature was raised to 80 °C and HCl (g) started to evolve. The reaction mixture became yellow turbid and turned to clear colorless after 30 min. After 8 h of heating, the mixture was stirred overnight at ambient temperature. Dry diethyl ether (150 mL) was added and the flask was stored overnight at -10 °C. The white product was collected on a glass-sintered funnel and dried (1.91 g). An additional amount of 2.65 g (product/ start. mat. 3:1) was obtained after the addition of i-hexane and storage at -10 °C. The combined material was taken up in 5% NaHCO_3 (200 mL) and extracted with ethyl acetate (3×150 mL). The combined organic layers were dried (Na_2SO_4) and evaporated in vacuo leaving a crude yellow oil (4.00 g; 64 %). ^1H NMR (HCl salt, CD_3OD) δ 3.02 (d, J = 6.83, 2H), 4.22 (t, J = 6.83, 1H), 5.15 (? , J = 3.93, 2H), 6.68 (d, J = 8.28, 1H), 6.91 (dd, J_1 = 8.28, J_2 = 2.18, 1H), 7.23-7.31 (m, 5H), 7.51 (d, J = 2.18, 1H).

[0323] PREPARATION OF X-2: The free base of X-1 (3.97 g, 10.0 mmol) was dissolved in dichloromethane (75 mL) and stirred at 0 °C under N_2 -atmosphere. Then, EDC (1.92 g, 10.0 mmol), HOBT (1.35 g, 10.0 mmol) and BOC-L-Phe (2.65 g, 10.0 mmol) were added simultaneously and triethylamine 1.39 mL, 10.0 mmol) was added dropwise. This reaction mixture was stirred for 15 h allowing to warm to ambient temperature. Ethyl acetate (200 mL) was added and the organic layer was washed with 5 % HCl (2×200 mL). The combined aqueous phases were extracted with ethyl acetate (100 mL) after which the combined organic layers were washed with 10% NaHCO_3 (100 mL). Drying (Na_2SO_4), filtration and evaporation in vacuo gave an off-white foam (6.01 g, 93%). The product was purified by flash column chromatography on silica gel eluting with chloroform: white foam C.Y. 78%. ^{13}C NMR (CDCl_3) δ 28.24, 36.52, 38.26, 53.34, 55.85, 67.36, 85.47, 115.02, 127.06, 128.59, 128.66, 128.71, 128.75, 129.30, 129.58, 131.04, 134.86, 136.39, 138.79, 151.04, 170.58, 170.91. (Rf 0.30 / i-hexane/ ethyl acetate 3:1).

[0324] PREPARATION OF X-3: A mixture of X-2 (4.43 g, 6.87 mmol), $\text{Pd}(\text{OAc})_2$ (50 mg, 3.3 mol%) and DPPF (230 mg, 6.2 mol%) in acetonitrile (20 mL) was treated with triethylamine (1.9 mL, 13.74 mmol) and methanol (4.4 mL). A carbon monoxide atmosphere was established and the reaction mixture was heated at 70 °C (Essential! Solvent vapour displaces CO if temperature is too high) for 16 h. The darkbrown reaction mixture was directly coated on silica gel and subjected to column chromatography (3×20 cm) eluting with chloroform. Pure fractions were pooled giving 2.45 g (62%) off-white solid after evaporation of the eluent. Pure material can be obtained by recrystallization from abs. ethanol. (Rf 0.15 / chloroform).

[0325] PREPARATION OF X-4: A mixture of X-3 (1.68 g, 2.91 mmol), methyl bromoacetate (0.83 μL , 3 eq.) and K_2CO_3 (activated, 1.20 g, 3 eq.) in acetone (20 mL) was heated at 50 °C overnight. TLC showed complete conversion and water (20 mL) was added. Extraction with dichloromethane (3×25 mL), drying (Na_2SO_4) and removal of the solvent at the rotavapor afforded 2.27 g of a yellow oil. Flash column chromatography on silica gel (2×20 cm) eluting

with chloroform gave 1.17 g (62%) of a pure colorless oil, that solidified on standing. An additional amount (0.45 g) impure colorless oil was isolated. (Rf 0.12 / chloroform).

[0326] PREPARATION OF X-5: X-4 (0.97 g, 1.50 mmol) was hydrogenated (atmospheric pressure) in abs ethanol (30 mL) using 10% Pd/C (100 mg) for 3 h. Filtration over diatomaceous earth and evaporation in vacuo of the filtrate

yielded 0.76 g (91 %) of a light-grey foam.
[0327] PREPARATION OF X-6: To a solution of X-5 (1.24 g, 2.24 mmol) in dry THF (10 mL) is added 1,1'-carbonyl-diimidazole (CDI, 0.54 g, 3.35 mmol). The solution is stirred at room temperature over night under nitrogen atmosphere. The reaction mixture is cooled with ice and a solution of NaBH₄ (0.21 g, 5.59 mmol) in H₂O (5 mL) is slowly added. After addition is complete, the mixture is stirred at room temperature for 10 min. The mixture is quenched with 10% aqueous HCl, and extracted with EtOAc. The organic layer is dried (Na₂SO₄) and concentrated. The residue is purified by flash chromatography (SiO₂, EtOAc) with furnished 160 mg (13%) of X-6 as a sticky foam. ¹H NMR 400 MHz (CDCl₃) δ 1.38, 2.49, 2.68, 2.73, 3.00, 3.42, 3.57, 3.78, 3.87, 4.05, 4.27, 4.68, 6.80, 7.13-7.30, 7.61.

[0328] PREPARATION OF X-7: To a solution of X-6 (36 mg, 0.066 mmol) in THF (1.5 mL) was added a 2.5 M aqueous solution of LiOH (106 µL, 0.26 mmol). The mixture was stirred at room temperature for 4 h, and then acidified with 10% aqueous HCl and extracted with EtOAc. The organic layer was dried (Na₂SO₄) and concentrated to afford 33 mg (96%) of X-7 as a white solid. Mp = 168.8-172.3 °C. ¹H NMR 400 MHz (MeOH) δ 1.35, 2.31, 2.64-2.97, 3.52, 4.06, 4.22, 4.78, 7.00, 7.11-7.27, 7.43, 7.77; ¹³C NMR (MeOH) δ 20.7, 28.6, 30.9, 36.8, 39.5, 57.3, 67.5, 80.6, 111.0115.6, 121.3, 127.6, 129.3, 130.3, 133.6, 134.0, 136.1, 138.6, 157.5, 169.1, 172.2, 174.1. MS (ESI) 516 (M-H). Anal. Calcd for C₂₆H₃₂N₂O₉ · H₂O: C, 59.42; H, 6.33; N, 5.33. Found; C, 59.4; H, 6.3; N, 5.35.4.

EXAMPLE 139 (COMPARISON) : (Chart Y, Formula Y-6) 2-[4-[2-[(3-carboxypropanoyl)amino]-2-methyl-3-oxo-3-(pentylamino)propyl]phenoxy]malonic acid

[0329] PREPARATION OF Y-2: To a solution of DL-α-methyl tyrosine (2.72 g; 13.9 mmol) and tetramethylammonium hydroxide pentahydrate (5.62 g; 31.0 mmol) in acetonitrile (270 mL) was added di-*tert*-butyldicarbonate (3.79 g; 17.4 mmol) and the resulting solution was allowed to stir 18 h at rt and concentrated [Khalil, E.M.; Subasinghe, N.L.; Johnson, R.L.: *Tet. Lett.* **1996**, 37, 3441]. The residue was partitioned between Et₂O/H₂O; the phases were separated and the aqueous phase extracted twice more with Et₂O. The aqueous phase was brought to pH 4 with solid citric acid and extracted with CHCl₃ (3 x 100 mL). The organic extracts were combined, dried (Na₂SO₄) and concentrated to afford 2.58 g (63%) 9 as a white foam. ¹H NMR (CDCl₃) δ 6.95 (d, J = 8 Hz, 2 H), 6.70 (d, J = 8 Hz, 2 H), 5.17 (br s, 1 H), 3.29 (br m, 1 H), 3.11 (br m, 1 H), 1.56 (s, 3 H), 1.47 (s, 9 H).

[0330] PREPARATION OF Y-3: To a solution of Y-2 (3.23 g; 10.9 mmol), diisopropylethylamine (2.09 mL; 12.0 mmol), and amylamine (1.39 mL; 12.0 mmol) in CH₂Cl₂ (250 mL) was added *N*[(dimethylamino)-1*H*-1,2,3-triazolo[4,5-*b*]pyridin-1-ylmethylene]-*N*-methylmethan ammonium hexafluorophosphonate *N*-oxide (4.63 g; 12.2 mmol), and the reaction was allowed to stir 18 h at rt. The reaction was washed with 1 M HCl (3 x 100 mL), saturated aq. NaHCO₃ (3 x 100 mL), and dried (Na₂SO₄). Upon concentration *in vacuo*, the residue was dissolved in a minimum of hot EtOAc and cooled to 4°C. The resulting precipitate was collected to afford 1.92 g (5.26 mmol; 48%) Y-3 as a white solid. Mp 170-2°C; UV λ_{max} 225 (9820, 95% EtOH); ¹H NMR (300 MHz, DMSO) δ 0.84 (t, J = 7 Hz, 3 H), 1.17-1.24 (m, 7 H), 1.38 (s, 9 H), 2.90-3.10 (m, 4 H), 3.35 (s, 2 H), 6.32 (br s, 1 H), 6.60 (d, J = 8 Hz, 2 H), 6.82 (d, J = 8 Hz, 2 H), 7.67 (br t, 1 H), 9.10 (s, 1 H).

[0331] PREPARATION OF Y-4: To a solution of Y-3 (1.94 g; 5.33 mmol) in acetone (100 mL) was added finely ground K₂CO₃ (2.28 g; 16.5 mmol) and dibenzyl bromomalonate (3.07 g; 8.46 mmol). After 18 h at rt, the reaction was diluted with H₂O (250 mL), extracted with EtOAc (3 x 100 mL), and dried (Na₂SO₄). Chromatography (mpic, E. Merck silica 60, 230-400) with 30% EtOAc/hexane afforded 1.11 g (32%) y-4 as a colorless oil. UV λ_{max} 222 (13000, 95% EtOH); ¹H NMR (300 MHz, CDCl₃) δ 7.34-7.25 (m, 10 H), 7.01 (d, J = 9 Hz, 2 H), 6.83 (d, J = 9 Hz, 2 H), 6.33 (br s, 1 H), 5.26 (s, 1 H), 5.22 (s, 4 H), 3.31 (d, J = 14 Hz, 1 H), 3.24-3.20 (m, 2 H), 3.00 (d, J = 14 Hz, 1 H), 1.45 (s, 9 H), 1.36 (s, 3 H), 1.38-1.29 (m, 6 H), 0.88 (t, J = 7 Hz, 3 H).

[0332] PREPARATION OF Y-5: HCl gas was bubbled into a solution of Y-4 (1.044 g; 1.61 mmol) for 15 min. After 4 h, the reaction was concentrated *in vacuo* and the residue dissolved in CH₂Cl₂ and triethylamine (0.49 mL; 3.54 mmol). Succinic anhydride (0.186 g; 1.86 mmol) was added and the reaction was allowed to stir 20 h at rt. An additional 0.186 g of succinic anhydride and 0.49 mL of triethylamine were added and the reaction allowed to stir an additional 20 h. The reaction was washed with 1 M HCl (2 x 20 mL), dried (MgSO₄), and chromatographed (mpic, E. Merck silica gel 60, 230-400 mesh) with 2% MeOH/CH₂Cl₂/0.5% AcOH to afford 0.567 g (54%) Y-5 as a colorless oil. ¹H NMR (300 MHz, CDCl₃) δ 7.33-7.25 (m, 10 H), 6.98 (d, J = 9 Hz, 2 H), 6.81 (d, J = 9 Hz, 2 H), 6.32 (br s, 1 H), 6.26 (br t, 1 H), 5.26 (s, 1 H), 5.22 (s, 4 H), 3.31-3.14 (m, 4 H), 2.70-2.67 (m, 2 H), 2.42-2.39 (m, 2 H), 1.50 (s, 3 H), 1.50-1.43 (m, 2 H), 1.30-1.24 (m, 4 H), 0.88 (t, J = 7 Hz, 3 H).

[0333] PREPARATION OF Y-6: A solution of Y-5 (0.225 g; 0.348 mmol) and 10% Pd/C (0.031 g) in MeOH (10 mL) was hydrogenated at atmospheric pressure for 18 h. The reaction was filtered through Celite, concentrated, and purified

by preparative RP HPLC using the conditions outlined below to afford 0.088g (54%) Y-6 as a hygroscopic white solid after lyophilization. UV λ_{\max} 223 (11400, 95% EtOH); ^1H NMR (300 MHz, DMSO- d_6) δ 7.59 (br s, 1 H), 7.45 (br t, 1 H), 6.96 (d, J = 8 Hz, 2 H), 6.79 (d, J = 8 Hz, 2 H), 5.24 (s, 1 H), 3.26 (d, J = 13 Hz, 1 H), 3.05-2.93 (m, 3 H), 2.45-2.42 (m, 2 H), 2.35-2.29 (m, 2 H), 1.38-1.29 (m, 2 H), 1.24-1.14 (m, 4 H), 1.13 (s, 3 H), 0.84 (t, J = 7 Hz, 3 H); ^{13}C NMR (75 MHz, DMSO- d_6) δ 174.8, 173.8, 171.3, 167.9, 155.9, 131.9, 130.8, 114.6, 76.1, 59.7, 30.9, 29.4, 29.1, 29.0, 24.1, 22.3, 14.1; IR (mull) 3362, 2729, 2669, 2592, 1732, 1631, 1612, 1544, 1511, 1299, 1226, 1184, 1120, 855, 839 cm^{-1} ; MS (FAB) m/z 467 (MH^+), 468, 467, 252, 208, 177, 157, 88, 43, 42, 23; HRMS (FAB) calcd for $\text{C}_{22}\text{H}_{30}\text{N}_2\text{O}_9 + \text{H}_1$ 467.2029, found 467.2040.

EXAMPLE 140 (COMPARISON) : (Chart Z, Formula Z-6) 2-[4-[2-[(3-carboxypropanoyl)amino]-3-oxo-3-(pentylamino)propyl]-2-fluorophenoxy]malonic acid

[0334] PREPARATION OF Z-2: To a suspension of commercially available 3-fluoro-DL-tyrosine (1.58 g; 7.93 mmol) in 0.5 M NaOH (19 mL) was added di-*tert*-butyl dicarbonate (2.12 g; 9.72 mmol) in THF (19 mL). After 26 h at rt, and additional 1.5 g of di-*tert*-butyl dicarbonate was added and the reaction stirred for an additional 2 h. The reaction was concentrated *in vacuo* to remove THF, and the residue was washed with Et₂O. The aqueous phase was acidified with 1 M citric acid, extracted with CHCl₃, and dried (Na₂SO₄) to afford 2.05 g (86%) Z-2 as a white foam. UV λ_{\max} 223 (8000, 95% EtOH); ^1H NMR (300 MHz, DMSO- d_6) δ 7.04 (d, J = 8 Hz, 1 H), 6.98 (d, J = 13 Hz, 1 H), 6.82-6.78 (m, 2 H), 3.99 (ddd, J = 5, 10, 14 Hz, 1 H), 2.88 (dd, J = 5, 14 Hz, 1 H), 2.68 (dd, J = 10, 14 Hz, 1 H), 1.30 (s, 9 H).

[0335] PREPARATION OF Z-3: N-Ethyl-N'-3-dimethylaminopropylcarbodiimide hydrochloride (1.38 g; 7.20 mmol) was added to a solution of Z-2 (1.78 g; 5.96 mmol) in CH₂Cl₂ (50 mL). After 10 min, amyl amine (0.83 mL; 7.2 mmol) was added and the reaction was allowed to stir 18h at rt. The reaction was washed with 1 M citric acid (3 x 25 mL), dried (Na₂SO₄), and concentrated *in vacuo* to afford 1.50 g (68%) Z-3 as a white solid. Mp. 63-4°C; UV λ_{\max} 272 (1600, 95% EtOH); ^1H NMR (300 MHz, CDCl₃) δ 6.94-6.79 (m, 3 H), 6.07 (t, J = 6 Hz, 1 H), 5.19 (br s, 1 H), 4.23 (q, J = 8 Hz, 1 H), 3.35-3.04 (m, 2 H), 2.93 (d, J = 7 Hz, 2 H), 1.40 (s, 9 H), 1.32-1.18 (m, 6 H), 0.84 (t, J = 7 Hz, 3 H).

[0336] PREPARATION OF Z-4: To a solution of Z-3 (1.49 g; 4.04 mmol) in acetone (60 mL) was added finely ground K₂CO₃ (1.72 g; 12.5 mmol) and dibenzyl bromomalonate (1.75 g; 4.83 mmol). After 18 h at rt, the reaction was diluted with H₂O (250 mL), extracted with EtOAc (3 x 100 mL), and dried (Na₂SO₄). Chromatography (mpc, E. Merck silica 60, 230-400) with 30% EtOAc/hexane afforded 0.799 g (30%) Z-4 as a colorless oil which solidified upon standing. Mp. 77-78.5°C; UV λ_{\max} 223 (12200, 95% EtOH); ^1H NMR (300 MHz, CDCl₃) δ 7.34-7.29 (m, 10 H), 6.97-6.81 (m, 3 H), 5.74 (br t, 1 H), 5.24 (s, 1 H), 5.23 (s, 4 H), 4.96 (br s, 1 H), 4.22-4.15 (m, 1 H), 3.15 (q, J = 7 Hz, 2 H), 2.97 (d, J = 7 Hz, 2 H), 1.41 (s, 9 H), 1.43-1.34 (m, 2 H), 1.29-1.16 (m, 4 H), 0.86 (t, J = 7 Hz, 3 H).

[0337] PREPARATION OF Z-5: HCl gas was bubbled into a solution of Z-4 (0.490 g; 0.753 mmol) for 10 min. After 18 h, the reaction was concentrated *in vacuo* and the residue dissolved in CH₂Cl₂ (15 mL) and triethylamine (0.25 mL; 1.8 mmol). Succinic anhydride (0.088 g; 0.879 mmol) was added and the reaction was allowed to stir 18 h at rt. The reaction was diluted with CH₂Cl₂ (50 mL), washed with 1 M HCl (2 x 30 mL), dried (MgSO₄), and chromatographed (mpc, E. Merck silica gel 60, 230-400 mesh) with 2% MeOH/CH₂Cl₂/1% AcOH to afford 0.0167 g (34%) Z-5 as a colorless oil. UV λ_{\max} 223 (11500, 95% EtOH); ^1H NMR (300 MHz, CDCl₃) δ 7.56 (d, J = 8 Hz, 1 H), 7.32-7.23 (m, 10 H), 6.95-6.78 (m, 3 H), 5.25 (s, 1 H), 5.19 (dd, J = 12, 14 Hz, 4 H), 4.69-4.61 (m, 1 H), 3.21-3.09 (m, 1 H), 3.05-2.86 (m, 3 H), 2.59-2.54 (m, 2 H), 2.47-2.40 (m, 2 H), 1.35-1.07 (m, 6 H), 0.82 (t, J = 7 Hz, 3 H).

[0338] PREPARATION OF Z-6: A solution of Z-5 (0.145 g; 0.223 mmol) and 10% Pd/C (0.026 g) in MeOH (10 mL) was hydrogenate at atmospheric pressure for 18 h. The reaction was filtered through Celite, concentrated, and purified by preparative RP HPLC using the conditions outlined below to afford 0.068g (65%) Z-6 as a hygroscopic white solid after lyophilization. UV λ_{\max} 222 (10200, 95% EtOH); ^1H NMR (300 MHz, DMSO- d_6) δ 8.08 (d, J = 8 Hz, 1 H), 7.82 (br t, 1 H), 7.07 (d, J = 12 Hz, 1 H), 6.96-6.89 (m, 2 H), 5.33 (s, 1 H), 4.39-4.32 (m, 1 H), 3.06-2.94 (m, 2 H), 2.90-2.84 (m, 1 H), 2.99-2.63 (m, 1 H), 2.35-2.21 (m, 4 H), 1.46-1.11 (m, 6 H), 0.82 (t, J = 7 Hz, 3 H); IR (mull) 3333, 2727, 2670, 2600, 1733, 1626, 1549, 1516, 1341, 1278, 1216, 1174, 1134, 1091, 722 cm^{-1} ; MS (FAB) m/z 471 (MH^+), 471, 193, 171, 167, 153, 135, 133, 121, 103, 89; HRMS (FAB) calcd for $\text{C}_{21}\text{H}_{27}\text{FN}_2\text{O}_9 + \text{H}_1$ 471.1779, found 471.1797.

EXAMPLE 141 (COMPARISON) : (Chart AA, Formula AA-#) 2-[4-[3-[(2-carboxyethyl)amino]-3-oxo-2-[(pentylamino)carbonyl]propyl]phenoxy]malonic acid

[0339] PREPARATION OF AA-2: To a solution of 4-hydroxybenzaldehyde (5.04 g, 41.3 mmol) was added finely ground K₂CO₃ (17.3 g; 125 mmol) and diethyl chloromalonate (7.34 mL; 45.4 mmol). After 18 h at rt, the reaction was concentrated *in vacuo*, and the residue taken up in water (200 mL). This was extracted with EtOAc (3 x 100 mL), dried (Na₂SO₄) and chromatographed (mpc, E. Merck silica gel 60, 230-400 mesh) to afford 7.97 g (69%) AA-2 as a pale yellow oil UV λ_{\max} 267 (15300, 95% EtOH); ^1H NMR (300 MHz, CDCl₃) δ 9.76 (s, 1 H), 7.73 (d, J = 9 Hz, 2 H), 6.96 (d, J = 9 Hz, 2 H), 5.24 (s, 1 H), 4.28-4.15 (m, 4 H), 1.18 (t, J = 7 Hz, 6 H).

[0340] PREPARATION OF AA-4: To a solution of commercially available ethyl hydrogen malonate (12.1 g; 91.7 mmol) and amyl amine (10.6 mL; 91.7 mmol) in CH_2Cl_2 (100 mL) was added diethyl cyanophosphonate (13.9 mL; 91.7 mmol) over 15 min. After 1.5 h at rt, the reaction was washed with 1 M citric acid (2 x 50 mL), H_2O (50 mL), saturated aqueous NaHCO_3 (2 x 50 mL), and dried (Na_2SO_4). Chromatography (mpc, E. Merck silica gel 60, 230-400 mesh) with 30% EtOAc/heptane afforded 8.9 g (48%) AA-4 as a pale yellow oil. ^1H NMR (300 MHz, CDCl_3) δ 7.28 (br s, 1 H), 4.00 (q, $J = 7$ Hz, 2 H), 3.12 (s, 2 H), 3.06 (q, $J = 7$ Hz, 2 H), 1.35 (quintet, $J = 7$ Hz, 2 H), 1.16-1.11 (m, 4 H), 1.10 (t, $J = 7$ Hz, 3 H), 0.71 (t, $J = 6$ Hz, 3 H).

[0341] PREPARATION OF AA-5: A solution of AA-4 (7.2 g; 36 mmol), 1.0 M NaOH (35.8 mL) and THF (100 mL) was allowed to stir at rt for 1 h. The reaction was concentrated *in vacuo* to remove THF, diluted with H_2O (200 mL), and washed with Et_2O (3 x 100 mL). The aqueous phase was brought to pH 1 with 3 M HCl and extracted with CH_2Cl_2 (3 x 100 mL). Upon drying (Na_2SO_4) and concentrating, 5.02 g (81 %) AA-5 was obtained as a white solid. Mp. 66-7°C; ^1H NMR (300 MHz, CDCl_3) δ 12.23 (br s, 1 H), 7.53 (br t, 1 H), 3.34 (s, 2 H), 3.25 (q, $J = 7$ Hz, 2 H), 1.51 (quintet, $J = 7$ Hz, 2 H), 1.33-1.27 (m, 4 H), 0.86 (t, $J = 7$ Hz, 3 H).

[0342] PREPARATION OF AA-6: To a solution of β -alanine ethyl ester hydrochloride (1.76 g; 11.5 mmol) and triethylamine (1.60 mL; 11.5 mmol) was added AA-5 (1.99 g; 11.5 mmol) and diethyl cyanophosphonate (1.74 mL; 11.5 mmol). After 18 h at rt, the reaction was washed with 1 M citric acid (3 x 50 mL) and dried (MgSO_4). Chromatography (mpc, E. Merck silica gel 60, 230-400 mesh) with 5% MeOH/ CH_2Cl_2 afforded 1.65 g (53%) AA-6 as a white solid. Mp. 102-3°C; UV λ_{max} 331 (8.99, 95% EtOH); ^1H NMR (300 MHz, CDCl_3) δ 7.52 (br s, 1 H), 7.26 (br s, 1 H), 4.12 (q, $J = 7$ Hz, 2 H), 3.50 (q, $J = 6$ Hz, 2 H), 3.20 (q, $J = 7$ Hz, 2 H), 3.13 (s, 2 H), 2.51 (s, 2 H), 1.48 (quintet, $J = 7$ Hz, 2 H), 1.29-1.26 (m, 4 H), 1.23 (t, $J = 7$ Hz, 3 H), 0.86 (t, $J = 7$ Hz, 3 H).

[0343] PREPARATION OF AA-7: A solution of AA-67 (0.981 g; 3.60 mmol), AA-2 (1.02 g; 3.62 mmol) and piperidine (1 mL) in toluene (100 mL) was brought to reflux under a Dean-Stark trap. After 21 h, the reaction was cooled, concentrated, and chromatographed (mpc, E. Merck silica gel 60, 230-400 mesh) with 1:1 EtOAc/heptane to afford 0.593 g (31 %) AA-7 as an orange oil which solidified upon standing. A second chromatography using 20%-50% EtOAc/heptane separated the two isomers. Earlier eluting isomer: ^1H NMR (300 MHz, CDCl_3) δ 7.72 (s, 1 H), 7.40 (br t, 1 H), 7.33 (d, $J = 9$ Hz, 2 H), 6.91 (d, $J = 9$ Hz, 2 H), 6.33 (br t, 1 H), 5.18 (s, 1 H), 4.34-4.27 (m, 4 H), 4.00 (q, $J = 7$ Hz, 2 H), 3.49 (q, $J = 6$ Hz, 2 H), 3.29 (q, $J = 7$ Hz, 2 H), 2.43 (t, $J = 6$ Hz, 2 H), 1.54 (quintet, $J = 7$ Hz, 2 H), 1.33-1.27 (m, 4 H), 1.30 (t, $J = 7$ Hz, 6 H), 1.17 (t, $J = 7$ Hz, 3 H), 0.89 (t, $J = 7$ Hz, 3 H). Later eluting isomer: ^1H NMR (300 MHz, CDCl_3) δ 7.79 (br t, 1 H), 7.69 (s, 1 H), 7.37 (d, $J = 9$ Hz, 2 H), 6.91 (d, $J = 9$ Hz, 2 H), 5.78 (br t, 1 H), 5.19 (s, 1 H), 4.35-4.27 (m, 4 H), 4.16 (q, $J = 7$ Hz, 2 H), 3.59 (q, $J = 6$ Hz, 2 H), 3.21 (q, $J = 7$ Hz, 2 H), 2.57 (t, $J = 6$ Hz, 2 H), 1.38 (quintet, $J = 7$ Hz, 2 H), 1.35-1.24 (m, 11 H), 1.24-1.14 (m, 2 H), 0.84 (t, $J = 7$ Hz, 3 H).

[0344] PREPARATION OF AA-8: A mixture of E and Z AA-7 (0.580 g; 1.08 mmol) in MeOH (25 mL) was hydrogenated at atmospheric pressure with 10% Pd/C (0.059 g) for 18 h. The reaction was filtered through a pad of Celite, concentrated, and chromatographed (flash, silica gel) with 60% EtOAc/heptane to afford 0.408 g (70%) AA-8 as a white solid. Mp. 88-90°C; UV λ_{max} 222 (11900, 95% EtOH); ^1H NMR (300 MHz, CDCl_3) δ 7.07 (d, $J = 9$ Hz, 2 H), 7.00 (br t, 1 H), 6.82 (d, $J = 9$ Hz, 2 H), 6.71 (br t, 1 H), 5.12 (s, 1 H), 4.32-4.25 (m, 4 H), 4.10 (q, $J = 7$ Hz, 2 H), 3.39 (q, $J = 6$ Hz, 2 H), 3.14-3.02 (m, 5 H), 2.45-2.30 (m, 2 H), 1.38 (quintet, $J = 7$ Hz, 2 H), 1.34-1.18 (m, 13 H), 0.85 (t, $J = 7$ Hz, 3 H).

[0345] PREPARATION OF AA-9: A solution of AA-8 (0.313 g; 0.583 mmol), 1.0 M NaOH (2.92 mL), H_2O (7 mL) and THF (10 mL) was stirred at rt. After 3 hr, the reaction was concentrated *in vacuo* to remove THF, diluted with H_2O (10 mL) and washed with CH_2Cl_2 (5 mL). The aqueous phase was brought to pH 2 with 1 M HCl, and allowed to stir until a white precipitate appeared (approx. 1 h). This was collected, washed with H_2O and dried to afford 0.216 g (82%) AA-9. UV λ_{max} 225 (10700, 95% EtOH); ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 7.86 (br t, 1 H), 7.75 (br t, 1 H), 6.99 (d, $J = 9$ Hz, 2 H), 6.78 (d, $J = 9$ Hz, 2 H), 5.12 (s, 1 H), 3.23-3.17 (m, 3 H), 2.99-2.95 (m, 2 H), 2.87-2.85 (m, 2 H), 2.27 (t, $J = 7$ Hz, 2 H), 1.31-1.11 (m, 6 H), 0.81 (t, $J = 7$ Hz, 3 H); ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) δ 173.3, 169.4, 169.0, 168.7, 156.2, 131.9, 129.8, 115.0, 75.6, 55.4, 32.2, 34.8, 34.1, 29.0, 28.9, 22.2, 14.4; IR (drift) 3283, 3053, 2953, 2928, 2871, 2859, 1768, 1725, 1663, 1639, 1551, 1512, 1275, 1245, 1205 cm^{-1} ; MS (FAB) m/z 453 (MH^+), 475, 454, 453, 431, 139, 107, 105, 103, 91, 23; HRMS (FAB) calcd for $\text{C}_{21}\text{H}_{28}\text{N}_2\text{O}_9$, 453.1873, found 453.1859.

EXAMPLE 142 (COMPARISON): (Chart CC, Formula CC-7) 2-[4-[2-(((2R)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropanoyl)amino)-3-oxo-3-(pentylamino)propyl]-2-(2-hydroxy-2-oxoethoxy)phenoxy]acetic acid

[0346] PREPARATION OF CC-2: K_2CO_3 (0.776 g; 5.61 mmol) and ethyl bromoacetate (0.63 mL; 5.6 mmol) were added to a solution of benzyl 2-[(tert-butoxycarbonyl)amino]-3-(3,4-dihydroxyphenyl)propanoate (CC-1) [Kaiser, Ado; Koch, Wolfgang; Scheer, Marcel; Woelcke, Uwe. N-Acyl-3,4-dihydroxy-L-phenylalanines. Ger. Offen., 32 pp. DE 2153811 720504] (0.997 g; 2.57 mmol) in acetone (100 mL). After 18 h at rt, the reaction was concentrated *in vacuo*, and the residue partitioned between H_2O (100 mL)/EtOAc (50 mL). The phases were separated, the aqueous phase was extracted with EtOAc (50 mL) and the organic phases were combined and dried (Na_2SO_4). Chromatography (mpc, E. Merck silica gel 60, 230-400 mesh) with 30% EtOAc/hexane afforded 0.691 g (48%) benzyl 3-[3,4-bis(2-ethoxy-

2-oxoethoxy)phenyl]-2-[(*tert*-butoxycarbonyl)amino]propanoate (CC-2) as a colorless oil. UV λ_{\max} 224 (8550, 95% EtOH); ^1H NMR (300 MHz, CDCl_3) δ 7.36-7.28 (m, 5 H), 6.73 (d, J = 8 Hz, 1 H), 6.62 (m, 2 H), 5.16-5.10 (m, 2 H), 4.94 (br d, 1 H), 4.67-4.51 (m, 5 H), 4.28-4.21 (m, 4 H), 2.92-2.88 (m, 2 H), 1.41 (s, 9 H), 1.32-1.26 (m, 6 H).

[0347] PREPARATION OF CC-3: 10% Pd/C (554 mg) was added to CC-2 (2.94 g, 5.25 mmol) followed by the addition of 100 mL of MeOH. The reaction mixture was stirred at rt under a H_2 atmosphere for 14 h, filtered through celite, concentrated and purified by column chromatography (1 % AcOH, 5% MeOH/ CH_2Cl_2) to afford CC-3 as a clear oil (1.86 g, 3.75 mmol, 71% yield). UV λ_{\max} 221 (7890, 95% Ethanol); (400 MHz, CDCl_3) δ 6.74-6.85 (m, 3 H), 4.97 (s, 1 H), 4.70 (s, 2 H), 4.69 (s, 2 H), 4.53 (s, 1 H), 4.27 (m, 4 H), 3.06 (s, 2 H), 1.44 (s, 9 H), 1.29 (m, 6 H).

[0348] PREPARATION OF CC-4: CC-3 (1.64 g, 3.50 mmol) was dissolved in 75 mL of CH_2Cl_2 , to the solution was added amylamine (450 μL , 3.88 mmol), triethylamine (610 μL , 4.02 mmol), and diethyl cyanophosphonate (610 μL , 4.38 mmol). The reaction mixture was stirred for 18 h, quenched with 10% aq citric acid, extracted with CH_2Cl_2 and washed with water. The organic layer was dried over NaSO_4 , filtered and condensed. The residue was purified by flash chromatography (30% EtOAc-50% EtOAc/ heptane) to afford CC-4 as a white solid (964 mg, 1.79 mmol, 51 % yield). UV λ_{\max} 221 (8920, 95% ETHANOL); ^1H NMR (400 MHz, CDCl_3) δ 7.28 (s, 1 H), 6.76-6.84 (m, 3 H), 5.75 (s, 1 H), 5.05 (s, 1 H), 4.70 (s, 2 H), 4.68 (s, 2 H), 4.21-4.30 (m, 4 H), 4.19 (m, 1 H), 3.14 (m, 2 H), 3.02 (dd, J = 14, 6 Hz, 1 H), 2.90 (dd, J = 14, 8 Hz, 1 H), 1.43 (s, 9 H), 1.15-1.41 (m, 12 H), 0.87 (t, J = 7 Hz, 3 H).

[0349] PREPARATION OF CC-5: CC-4 (765 mg, 1.42 mmol) was dissolved in 25 mL of 20% TFA/ CH_2Cl_2 solution, the reaction mixture was stirred for 1 h at rt then condensed to afford a yellow oil. The oil was redissolved in CH_2Cl_2 , washed with NaHCO_3 and water. The organic layer was dried over NaSO_4 , filtered and condensed. The residue was purified by flash chromatography (5% MeOH sat. NH_3 / CH_2Cl_2) to afford CC-5 as a clear oil. ^1H NMR (400 MHz, CDCl_3) δ 6.76-6.84 (m, 3 H), 4.69 (s, 4 H), 4.25 (q, J = 10 Hz, 4 H), 3.54 (m, 1 H), 3.13-3.25 (m, 3 H), 2.60 (dd, J = 18, 13 Hz, 1 H), 1.50 (m, 2 H), 1.26-1.32 (m, 10 H), 0.89 (t, J = 9 Hz, 3 H).

[0350] PREPARATION OF CC-6: CC-5 (362 mg, 0.826 mmol) was dissolved in 50 mL of CH_2Cl_2 to the solution was added triethylamine (160 μL , 1.18 mmol), diethylcyanophosphonate (160 μL , 1.06) and Boc-L-Phe (262 mg, 0.99 mmol). The reaction mixture was stirred at rt for 16 h, quenched with 10% aq citric acid, extracted with CH_2Cl_2 and washed with water. The organic layer was dried over NaSO_4 , filtered, and condensed. The residue was purified by column chromatography (5% MeOH/ CH_2Cl_2) to afford CC-6 (1:1 mixture of diastereomers) as a white solid (418 mg, 0.610 mmol, 74% yield). UV λ_{\max} 224 (10300, MeOH); ^1H NMR (400 MHz, CDCl_3) δ 7.18-7.32 (m, 5 H), 6.63-6.46 (m, 3 H), 6.34 (m, 1 H), 5.98-6.07 (m, 1 H), 5.00-5.07 (m, 1 H), 4.67-4.71 (m, 4 H), 4.55 (m, 1 H), 4.09-4.28 (m, 5 H), 2.90-3.18 (m, 5 H), 2.52-2.82 (m, 1H), 1.15-1.43 (m, 21 H), 0.86 (m, 3 H).

[0351] PREPARATION OF CC-7: CC-6 (160 mg, 0.243 mmol) was dissolved in 15 mL of THF, to the solution was added 1.0 N NaOH (800 μL , 0.800 mmol), the reaction mixture was stirred for 16 h at rt. The reaction was quenched with 10% citric acid, extracted with CH_2Cl_2 , the organic layer was dried over NaSO_4 , filtered and condensed to afford CC-7 as a white solid (122 mg, 0.201 mmol, 83% yeild). UV λ_{\max} 226 (9460, MeOH); ^1H NMR (400 MHz, DMSO- d_6) δ 12.90 (s, 2 H), 7.76-8.24 (m, 2 H), 7.10-7.24 (m, 5 H), 6.65-6.90 (m, 4 H), 4.61 (m, 4 H), 4.40 (m, 1 H), 4.11 (m, 1 H), 2.63-2.87 (m, 6 H), 1.15-1.35 (m, 15 H), 0.84 (m, 3 H); ^{13}C NMR (100 MHz, DMSO- d_6) δ 180.9, 180.8, 180.7, 179.9, 179.8, 179.7, 179.7, 179.6, 164.9, 164.6, 156.5, 155.5, 155.4, 147.5, 147.5, 147.4, 140.7, 140.3, 138.6, 138.5, 137.5, 137.4, 135.6, 131.6, 131.4, 124.8, 223.6, 123.6, 87.7, 87.6, 81.9, 74.9, 74.7, 74.6, 65.4, 65.2, 63.6, 63.3, 52.2, 48.0, 47.9, 46.9, 46.7, 39.9, 38.1, 38.0, 38.0, 37.9, 37.6, 37.2, 31.3, 23.3; IR (drift) 3316, 2958, 2931, 1756, 1710, 1688, 1646, 1614, 1551, 1517, 1271, 1250, 1192, 1168, 1145 cm^{-1} . HRMS (FAB) calcd for $\text{C}_{32}\text{H}_{43}\text{N}_3\text{O}_{10}+\text{H}_1$ 630.3026, found 630.3038.

EXAMPLE 143: (Chart DD, Formula DD-3) 2-(carboxymethoxy)-5-[(2S)-3-oxo-2-[[[(2R)-2-(2-oxo-1-pyrrolidinyl)-3-phenylpropanoyl]amino]-3-(pentylamino)propyl]benzoic acid

[0352] PREPARATION OF DD-1: Q-4 (2.30 g, 4.53 mmol) was dissolved in 20 mL of 20% TFA/ CH_2Cl_2 solution, the reaction mixture was stirred at rt for 5 h. The solution was condensed, dissolved in CH_2Cl_2 , washed with saturated aq NaHCO_3 , organics dried over NaSO_4 . Purified by column chromatography (5%MeOH (sat. NH_3)/ CH_2Cl_2) to afford DD-1 as a clear oil (1.48 g, 3.62 mmol, 80% yield). $[\alpha]_D^{25} = 18^\circ$ (c 0.64, methanol); UV λ_{\max} 226 (9750, MeOH); ^1H NMR (300 MHz, CDCl_3) δ 7.65 (d, J = 2 Hz, 1 H), 7.32 (d, J = 2 Hz, 1 H), 7.29 (d, J = 2 Hz, 1 H), 6.85 (d, J = 9 Hz, 1 H), 4.68 (s, 2 H), 4.35 (q, J = 7 Hz, 2 H), 4.25 (q, J = 7 Hz, 2 H), 3.56 (dd, J = 9, 4 Hz, 1 H), 3.21 (m, 3 H), 2.71 (dd, J = 14, 9 Hz, 1 H), 1.25-1.50 (m, 12 H), 0.89 (t, J = 7 Hz, 3 H).

[0353] PREPARATION OF DD-3: (2S)-2-(2-Oxo-1-pyrrolidinyl)-3-phenylpropanoic acid [Ocaïn, T.D.; Deininger, D. D., US Patent 5,023,338] (124 mg, 0.532 mmol) was added to 15 ml of CH_2Cl_2 , to the reaction mixture was added triethyl amine (75 μL , 0.54 mmol), DD-1 (225 mg, 0.551 mmol) and diethyl cyanophosphonate (90 μL , 0.59 mmol). The reaction mixture was stirred at rt for 12 h, diluted with CH_2Cl_2 , quenched with 10% aq citric acid, washed with water, organics dried over NaSO_4 , filtered and condensed. The resulting product DD-2 was dissolved in 10 mL THF/ water (1:1), to the reaction mixture was added 1 N NaOH solution (600 μL , 0.600 mmol), the reaction mixture was stirred at

rt for 2 h then quenched with 10% aq citric acid, extracted with EtOAc, washed with water, dried organics over Na₂SO₄, filtered, and condensed. Purified by column chromatography (10% MeOH/ CH₂Cl₂, 1% AcOH, to give a white solid as a diastereomer mixture 7:1, (175 mg, 0.31 mmol, 56% yield). Separated major diastereomers by HPLC to give DD-3 as a white solid. Mp. 96-97°C; ¹H NMR (400 MHz, DMSO) δ 7.98 (d, J = 8 Hz, 1 H), 7.87 (t, J = 4 Hz, 1 H), 7.56 (d, J = 2 Hz, 1 H), 7.32 (dd, J = 10, 2 Hz, 1 H), 7.20 (m, 2 H), 7.12 (m, 3 H), 6.91 (d, J = 9 Hz, 1 H), 4.80 (dd, J = 10.5 Hz, 1 H), 4.71 (s, 3 H), 4.45 (m, 1 H), 2.90-3.12 (m, 5 H), 2.72-2.83 (m, 3 H), 2.07 (m, 1 H), 1.89 (m, 1 H), 1.75 (m, 1 H), 1.62 (m, 1 H), 1.36 (m, 2 H), 1.14-1.23 (m, 5 H), 0.85 (t, J = 7 Hz, 3 H); ¹³C NMR (100 MHz, DMSO) δ 174.2, 170.5, 170.1, 169.2, 167.0, 155.3, 137.7, 133.6, 131.6, 130.4, 128.7, 128.0, 126.2, 121.0, 113.4, 65.5, 54.6, 54.2, 43.0, 38.5, 36.3, 33.8, 30.2, 28.6, 28.5, 21.8, 17.6, 13.9; MS (ESI+) for C₃₀H₃₇N₃O₈ m/z 568.2 (M+H)⁺.

EXAMPLE 144 (COMPARISON): (Chart EE, Formula EE-4) 5-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropanoyl]amino)-3-oxo-3-(pentylamino)propyl]-2-[carboxy(fluoro)methoxy]benzoic acid

[0354] PREPARATION OF BE-1: Triethylamine (3.37 mL; 24.2 mmol) was added to a solution of Q-2 (5.76 g; 12.1 mmol), 1,1'-bis(diphenylphosphino)ferrocene (0.400 g; 0.721 mmol) and palladium (II) acetate (0.0814 g; 0.362 mmol) in 25 mL DMF/10 mL benzyl alcohol. The reaction was placed under a CO atmosphere and heated to 60°C for 12 h. The reaction was diluted with EtOAc (75 mL), washed with 1 M HCl (3 x 25 mL), and dried (Na₂SO₄). The organic phase was filtered, absorbed onto silica gel, and chromatographed (flash, silica gel) with 20% EtOAc/heptane. The white precipitate which formed in the resulting fractions was collected and washed with 20% EtOAc/heptane to afford 1.47 g EE-1. The filtrate was combined with the remaining product-containing fractions and rechromatographed with 10%-20% EtOAc/heptane to afford an additional 0.902 g EE-1 as a solid. The mixed cuts (containing residual benzyl alcohol) were evaporated to dryness at 70°C/1 Torr to afford an additional 0.527 g EE-1 as a solid. Total yield: 2.90 g (49%). ¹H NMR (300 MHz, CDCl₃) δ 10.70 (s, 1 H), 7.68 (d, J = 2 Hz, 1 H), 7.46-7.35 (m, 5 H), 7.30 (dd, J = 2.9 Hz, 1 H), 6.89 (d, J = 9 Hz, 1 H), 6.12 (br t, 1 H), 5.36 (s; 2 H), 5.25 (br d, 1 H), 4.25-4.22 (m, 1 H), 3.15-3.07 (m, 2 H), 2.97-2.92 (m, 2 H), 1.37 (s, 9 H), 1.41-1.14 (m, 6 H), 0.85 (t, J = 7 Hz, 3 H).

[0355] PREPARATION OF EE-2: Ethyl bromofluoroacetate (0.26 mL; 2.2 mmol) was added to EE-1 (0.902 g; 1.86 mmol) and finely ground K₂CO₃ (1.28 g; 9.24 mmol) in acetone (100 mL) and the reaction was stirred 18 h at rt. The reaction was filtered through Whatman 42 paper, concentrated, and chromatographed (flash, silica) with 30%-50% EtOAc/heptane to afford 0.809 g (74%) EE-2 as a white solid. Mp. 88-90°C; [α]_D²⁵ = 1° (c 0.91, chloroform); UV λ_{max} 229 (10300, 95% EtOH); ¹H NMR (300 MHz, CDCl₃) δ 7.73 (t, J = 3 Hz, 1 H), 7.42-7.28 (m, 6 H), 7.16 (d, J = 8 Hz, 1 H), 6.27 (br s, 1 H), 5.88 (dd, J = 1, 60 Hz, 1 H), 5.35-5.30 (m, 3 H), 4.28-4.21 (m, 3 H), 3.12-3.08 (m, 3 H), 2.99-2.94 (m, 1 H), 1.39-1.17 (m, 6 H), 1.36 (s, 9 H), 1.29 (t, J = 7 Hz, 3 H), 0.85 (t, J = 7 Hz, 3 H).


[0356] PREPARATION OF EE-3: Trifluoroacetic acid (12 mL) was added to a solution of EE-2 (0.659 g; 1.12 mmol) in CH₂Cl₂ (36 mL) and allowed to stir at rt for 20 min. The reaction was concentrated, and the residue was dissolved in CH₂Cl₂ (70 mL), washed with 5% NaHCO₃ (12 mL), water (2 x 12 mL), and dried (MgSO₄). Removal of the solvent *in vacuo* afforded (0.543 g) of a yellow oil, which was used without further purification. A 0.438 g portion of this oil in CH₂Cl₂ (40 mL) was treated with triethylamine (0.187 g; 1.34 mmol), N-tert-butyl-L-phenylalanine (0.286 g; 1.08 mmol), and diethylcyanophosphonate (0.203 g; 1.34 mmol). After 18 h at rt, the reaction was diluted with CH₂Cl₂ (40 mL), washed with 1 M citric acid (2 x 20 mL), and dried (MgSO₄). Chromatography (flash, silica) with 50% EtOAc/heptane afforded 0.504 g (76%) EE-3 as a white solid. [α]_D²⁵ = -24° (c 1.02, chloroform); UV λ_{max} 230 (10500, 95% EtOH); ¹H NMR (300 MHz, CDCl₃) δ 7.42-7.14 (m, 13 H), 6.74 (br s, 1 H), 6.36 (br s, 1 H), 5.88 (dd, J = 5, 60 Hz, 1 H), 5.31 (s, 2 H), 5.14 (br t, 1 H), 4.65-4.61 (m, 1 H), 4.35-4.32 (m, 1 H), 4.26-4.22 (m, 2 H), 3.20-2.89 (m, 6 H), 1.33-1.16 (m, 18 H), 0.85 (t, J = 7 Hz, 3 H).

[0357] PREPARATION OF EE-4: A solution of EE-3 (0.389g; 0.529mmol) in MeOH (25 mL) was hydrogenated at atmospheric pressure with 10% Pd/C (80.3 mg). After 23 h the catalyst was removed by filtration through Whatman 42 paper, and the filtrate concentrated to a white solid. Mass spectroscopy showed a mixture of methyl and ethyl esters: MS (ESI-) for C₃₉H₄₈FN₃O₉ m/z 630.1 (M-H)⁻. MS (ESI-) for C₄₀H₅₀FN₃O₉ m/z 644.1 (M-H)⁻. A 99.3 mg portion of this mixture was treated with 0.0509 M LiOH (6.35 mL) in THF (6 mL). After 15 min. rxn concentrated *in vacuo* to remove THF, acidified with 1 M citric acid. The resulting solid was purified by preparative RP HPLC to afford 21 mg EE-4 as a white solid after lyophilization. UV λ_{max} 229 (8470, 95% EtOH); ¹H NMR (300 MHz, DMSO-d₆) δ 7.96-7.94 (m, 2 H), 7.62 (s, 1 H), 7.6-7.4 (m, 1 H), 7.24-7.17 (m, 6 H), 9.91 (br d, 1 H), 6.17 (d, J = 59 Hz, 1 H), 4.50-4.40 (m, 1 H), 4.14-4.04 (m, 1 H), 3.08-2.91 (m, 3 H), 2.86-2.79 (m, 2 H), 2.71-2.61 (m, 1 H), 1.37-1.12 (m, 15 H), 0.84 (t, J = 7 Hz, 3 H); IR (drift) 3311, 2959, 2931, 2871, 2860, 1689, 1646, 1525, 1498, 1450, 1441, 1368, 1291, 1242, 1166 cm⁻¹; MS (FAB) m/z 618 (MH⁺), 518, 120, 88, 86, 57, 43, 41, 39, 29, 23; HRMS (FAB) calcd for C₃₁H₄₀FN₃O₉+H₁ 618.2827, found 618.2851.

EXAMPLE 145: (Chart FF, Formula FF-2) 5-((2S)-2-(((2S)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropanoyl)amino)-3-oxo-3-[(4-phenylbutyl)amino]propyl)-2-(carboxymethoxy)benzoic acid

[0358] To a solution of X-5 (100 mg, 0.18 mmol) in dichloromethane (500 μ L) was added 1-hydroxybenzotriazole (31 mg, 0.23 mmol) in N,N-dimethylformamide (100 μ L) and 4-phenylbutylamine (46 μ L, 0.29 mmol). The mixture was cooled in an ice bath and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (44 mg, 0.23 mmol) in dichloromethane (2 mL) was added. The mixture was stirred in room temperature for 3h. The reaction mixture was applied on a small silica gel column (5 mL) packed in dichloromethane and the amide FF-1 was eluted using a stepwise gradient of dichloromethane-acetonitrile. The amide was then dissolved in tetrahydrofuran-methanol (3 mL, 2:1) and aqueous sodium hydroxide (1.5 mL, 2%) was added. The mixture was stirred at room temperature for 6 h. Acetic acid (40 μ L) was added and the mixture was concentrated. The material was purified by reversed phase HPLC (Vydac C-18 column) using a water-acetonitrile gradient and lyophilized to give FF-2 (34 mg) as a white solid. $^1\text{H-NMR}$ (400 MHz, CD_3OD) δ 7.76 (s, 1H), 7.38 (d, $J=8.5$ Hz, 1H), 7.26-7.11 (m, 10H), 6.98 (d, $J=8.5$ Hz, 1H), 4.76 (s, 2H), 4.50 (t, $J=6.9$ Hz, 1H), 4.23 (dd, $J=5.2$ Hz, $J=9.3$ Hz, 1H), 3.21-2.72 (m, 6H), 2.58 (t, $J=7.5$ Hz, 2H), 1.54 (m, 2H), 1.42 (m 2H), 1.34 (s, 9H); IR (KBr) 3296, 2925, 1738, 1687, 1643 cm^{-1} ; HRMS m/z 661.2987 (calc. of monoisotopic mass for $\text{C}_{36}\text{H}_{43}\text{N}_3\text{O}_9$ gives 661.2999).

CHART A






CHART D









CHART E




E-1



E-2




E-3




E-4: R = CH₂Ph

E-5: R = Me




E-6: R = CH₂Ph

E-7: R = Me



E-8



E-9

CHART F




CHART G




CHART H




CHART I




CHART J




CHART K




CHART L




CHART M






CHART N




CHART O




CHART P




CHART Q




CHART R




CHART S




CHART T





CHART U



The reaction scheme illustrates the synthesis of compound V-6 from V-1 through a series of chemical transformations:

- V-1** (4-hydroxy-2-(tert-butoxycarbonylamino)benzoic acid methyl ester) is converted to **V-2** (4-hydroxy-2-(tert-butoxycarbonylamino)benzaldehyde).
- V-2** is converted to **V-3** (4-hydroxy-2-(tert-butoxycarbonylamino)-3-(hept-5-en-1-yl)benzoic acid methyl ester).
- V-3** is converted to **V-4** (4-hydroxy-2-(tert-butoxycarbonylamino)-3-(hept-5-en-1-yl)benzoic acid).
- V-4** is converted to **V-5** (4-(2-(2-((tert-butoxycarbonylamino)(hept-5-en-1-yl)vinyl)oxy)acetic acid)benzoic acid).
- V-5** is converted to **V-6** (4-(2-(2-((tert-butoxycarbonylamino)(hept-5-en-1-yl)vinyl)oxy)acetic acid)benzoic acid).

CHART W









CHART X




CHART Y




CHART Z




CHART AA




CHART BB







CHART CC






CHART DD




CHART EE




CHART FF




TABLE 1

Example Number, Structure	PTP1 Inhib	
	Conc (uM)	% Inhib
Example 1 	100 10 1	80 8 3
Example 3 	100 100 10 1	81 76 25 1
Example 2 	10 1 100	9 4 39
Example 12 	10 1 100	17 2 67

TABLE 1 (CONT'D.)




Example Number, Structure	PTP1 Inhib	
	Conc (uM)	% Inhib
Example 13 	10 1 100	13 3 62
Example 14 	10 1 100	12 3 53
Example 15 	10 1 100	20 2 72

TABLE 1 (CONTINUED)

Example Number, Structure	PTP1 Inhib	
	Conc (uM)	% Inhib
Example 16 	10 100 1	20 74 4
Example 56 	100 10 1	34 1 0
Example 17 	100 10 1	67 18 0

TABLE 1 (CONT'D.)

Example Number, Structure	PTP1 Inhib	
	Conc (uM)	% Inhib
Example 18 	100 10 1	63 14 2
Example 19 	100 10 1	41 5 4
Example 29 	100 10 1	74 17 3

TABLE 1 (CONT'D.)



Example Number, Structure	PTP1 Inhib	
	Conc (uM)	% Inhib
Example 30 	100 10 1	73 21 0
Example 21 	100 10 1	83 33 1

TABLE 1 (CONT'D.)




Example Number, Structure	PTP1 Inhib	
	Conc (uM)	% Inhib
Example 22 	100 10 1	57 11 0
Example 31 	100 10 1	46 7 0
Example 32 	100 10 1	48 6 0

TABLE 1 (CONT'D.)

Example Number, Structure	PTP1 Inhib	
	Conc (uM)	% Inhib
Example 33 	100 10 1	23 2 0
Example 55 	10 100 1	9 50 0
Example 34 	100 10 1	35 6 0
Example 57 	100 10 1	39 4 3

TABLE 1 (CONT'D.)




Example Number, Structure	PTP1 Inhib	
	Conc (uM)	% Inhib
Example 41 	100 10 1	45 3 0
Example 35 	100 10 1	26 3 2
Example 23 	100 10 1	37 4 0

TABLE 1 (CONT'D.)





Example Number, Structure	PTP1 Inhib	
	Conc (uM)	% Inhib
Example 24 	100 10 1	72 20 1
Example 4 	10 100 1	23 75 4
Example 5 	10 100 1	18 63 6
Example 26 	100 10 1	96 75 27

TABLE 1 (CONT'D.)





Example Number, Structure	PTP1 Inhib	
	Conc (uM)	% Inhib
Example 25 	100 10 1	90 53 13
Example 42 	100 10 1	63 24 12
Example 43 	100 10 1	72 23 8
Example 44 	100 10 1	61 18 7

TABLE 1 (CONT'D.)





Example Number, Structure	PTP1 Inhib	
	Conc (uM)	% Inhib
Example 45 	100 10 1	51 3 4
Example 46 	100 10 1	63 19 4
Example 6 	100 10 1	81 34 8
Example 39 	10 100 1	3 34 0

TABLE 1 (CONT'D.)





Example Number, Structure	PTP1 Inhib	
	Conc (uM)	% Inhib
Example 7 	100 10 1 100 10 1	72 21 2 74 21 3
Example 8 	10 100 1	64 94 16
Example 47 	10 100 1	15 66 4
Example 27 	100 10 1	83 34 5

TABLE 1 (CONT'D.)





Example Number, Structure	PTP1 Inhib	
	Conc (uM)	% Inhib
Example 28 	100 10 1	96 76 25
Example 9 	100 10 1	94 64 17
Example 38 	100 10 1	79 29 5
Example 10 	10 100 1	25 75 3

TABLE 1 (CONT'D.)

Example Number, Structure	PTP1 Inhib	
	Conc (uM)	% Inhib
Example 36 	100 10 1	93 63 16
Example 48 	100 10 1	57 13 1
Example 52 	100 10 1	90 51 10
Example 37 	10 100 1	57 91 14

TABLE 1 (CONT'D.)





Example Number, Structure	PTP1 Inhib	
	Conc (uM)	% Inhib
Example 40 	100 10 1	33 6 3
Example 49 	100 10 1	43 7 1
Example 51 	100 10 1	51 11 0
Example 53 	100 10 1	94 62 16

TABLE 1 (CONT'D.)





Example Number, Structure	PTP1 Inhib	
	Conc (uM)	% Inhib
Example 54 	100 10 1	98 87 42
Example 50 	100 10 1	56 14 3
Example 11 	100 10 1	90 35 5
Example 58 	100 10 1	93 63 17

TABLE 2





PNU-number, Structure	PTP1B Inhib		PTP1 Inhib	
	PTP1B. Conc (uM)	PTP1B. % Inhib	PTP1N. Conc (uM)	PTP1N. % Inhib
EXAMPLE 59 			100 10 1	98 80 30
EXAMPLE 65 	100 10 1	90 42 9	100 10 1	91 44 10
EXAMPLE 60 			100 10 1	54 13 4
EXAMPLE 98 			100 10 1	60 13 1

TABLE 2 CONTINUED





PNU-number, Structure	PTP1B Inhib		PTP1 Inhib	
	PTP1B. Conc (uM)	PTP1B. % Inhib	PTP1N. Conc (uM)	PTP1N. % Inhib
EXAMPLE 72 			100 10 1	89 47 9
EXAMPLE 61 			100 10 1	98 70 11
EXAMPLE 66 			100 10 1	85 12 3
EXAMPLE 62 			10 1 100	83 37 97

TABLE 2 CONTINUED

PNU-number, Structure	PTP1B Inhib		PTP1 Inhib	
	PTP1B. Conc (μ M)	PTP1B. % Inhib	PTP1N. Conc (μ M)	PTP1N. % Inhib
EXAMPLE 67 			100 10 1	89 36 5
EXAMPLE 63 	100 10 1	99 87 45	100 10 1	97 84 37
EXAMPLE 68 			100 10 1	90 36 8
EXAMPLE 69 	100 10 1	91 48 7	100 10 1	88 42 9

TABLE 2 CONTINUED

PNU-number, Structure	PTP1B Inhib		PTP1 Inhib	
	PTP1B. Conc (uM)	PTP1B. % Inhib	PTP1N. Conc (uM)	PTP1N. % Inhib
EXAMPLE 64 			100 10 1	98 83 32
EXAMPLE 70 	100 10 1	83 36 7	100 10 1	76 22 0
EXAMPLE 116 see comment			100 10 1	93 57 8
EXAMPLE 71 			100 10 1	32 0 0

TABLE 2 CONTINUED






PNU-number, Structure	PTP1B Inhib		PTP1 Inhib	
	PTP1B. Conc (μ M)	PTP1B. % Inhib	PTP1N. Conc (μ M)	PTP1N. % Inhib
EXAMPLE 117 	100 10 1	97 74 21	100 10 1	93 55 8
EXAMPLE 118 			100 10 1	92 51 4
EXAMPLE 131 			100 10 1	93 56 12
EXAMPLE 119 	100 10 1	97 63 14	100 10 1	96 55 8
EXAMPLE 139 			100 10 1	34 6 3

TABLE 2 CONTINUED





PNU-number, Structure	PTP1B Inhib		PTP1 Inhib	
	PTP1B. Conc (uM)	PTP1B. % Inhib	PTP1N. Conc (uM)	PTP1N. % Inhib
EXAMPLE 140 			100 10 1	78 29 4
EXAMPLE 99 			100 10 1	57 15 3
EXAMPLE 73 			100 10 1	87 42 9
EXAMPLE 120 	100 10 1	90 45 6		

TABLE 2 CONTINUED





PNU-number, Structure	PTP1B Inhib		PTP1 Inhib	
	PTP1B. Conc (uM)	PTP1B. % Inhib	PTP1N. Conc (uM)	PTP1N. % Inhib
EXAMPLE 81 	100 10 1 100 10 1 100 10 1	97 78 30 89 54 15 87 43 8	100 10 1	97 84 40
EXAMPLE 121 	100 10 1	58 9 3		
EXAMPLE 122 	100 10 1	42 8 5		
EXAMPLE 123 	100 10 1 100 10 1	95 65 18 93 59 14		

TABLE 2 CONTINUED





PNU-number, Structure	PTP1B Inhib		PTP1N Inhib	
	PTP1B. Conc (uM)	PTP1B. % Inhib	PTP1N. Conc (uM)	PTP1N. % Inhib
EXAMPLE 124 	100 10 1 100 10 1	95 68 19 92 63 19		
EXAMPLE 125 	100 10 1	94 63 17		
EXAMPLE 100 	100 10 1	44 8 2		
EXAMPLE 101 	100 10 1 10 100 1	75 14 2 16 83 3		

TABLE 2 CONTINUED





PNU-number, Structure	PTP1B Inhib		PTP1 Inhib	
	PTP1B. Conc (uM)	PTP1B. % Inhib	PTP1N. Conc (uM)	PTP1N. % Inhib
EXAMPLE 74 	100 10 1	86 40 9		
EXAMPLE 102 	100 10 1 100 10 1	93 29 1 90 29 3		
EXAMPLE 103 	100 10 1	86 33 3		
EXAMPLE 104 	100	79		

TABLE 2 CONTINUED





PNU-number, Structure	PTP1B Inhib		PTP1 Inhib	
	PTP1B. Conc (uM)	PTP1B. % Inhib	PTP1N. Conc (uM)	PTP1N. % Inhib
EXAMPLE 104	10 1	28 6		
EXAMPLE 126 	100 10 1	87 45 11		
EXAMPLE 105 	100 10 1 100 10 1	74 25 5 82 30 5		
EXAMPLE 106 	100 10 1	67 21 7		
EXAMPLE 107 	100	80		

TABLE 2 CONTINUED





PNU-number, Structure	PTP1B Inhib		PTP1 Inhib	
	PTP1B. Conc (uM)	PTP1B. % Inhib	PTP1N. Conc (uM)	PTP1N. % Inhib
EXAMPLE 107	10 1	25 5		
EXAMPLE 97 	100 10 1	60 15 4		
EXAMPLE 108 	100 10 1	76 24 4		
EXAMPLE 109 	100 10 1	71 19 3		
EXAMPLE 110 	100 10 1	85 40 9		

TABLE 2 CONTINUED





PNU-number, Structure	PTP1B Inhib		PTP1N Inhib	
	PTP1B. Conc (uM)	PTP1B. % Inhib	PTP1N. Conc (uM)	PTP1N. % Inhib
EXAMPLE 111 	100 10 1	66 19 6		
EXAMPLE 112 	100 10 1	90 50 10		
EXAMPLE 113 	100 10 1 100 10 1	58 7 2 77 17 2		
EXAMPLE 82 	100 10 1 100 10 1	98 93 64 98 92 66		

TABLE 2 CONTINUED





PNU-number, Structure	PTP1B Inhib		PTP1 Inhib	
	PTP1B. Conc (uM)	PTP1B. % Inhib	PTP1N. Conc (uM)	PTP1N. % Inhib
EXAMPLE 75 	100 10 1	86 40 9		
EXAMPLE 76 	100 10 1	92 54 12		
EXAMPLE 77 	100 10 1	93 62 14		
EXAMPLE 114 	100 10 1	69 19 2		

TABLE 2 CONTINUED









PNU-number, Structure	PTP1B Inhib		PTP1 Inhib	
	PTP1B. Conc (uM)	PTP1B. % Inhib	PTP1N. Conc (uM)	PTP1N. % Inhib
EXAMPLE 115 	100 10 1 100 10 1	97 81 32 96 78 28		
EXAMPLE 128 	100 10 1	96 82 36		
EXAMPLE 129 	100 10 1	98 91 57		
EXAMPLE 130 	100 10 1	98 88 49		

TABLE 2 CONTINUED

PNU-number, Structure	PTP1B Inhib		PTP1 Inhib	
	PTP1B. Conc (μ M)	PTP1B. % Inhib	PTP1N. Conc (μ M)	PTP1N. % Inhib
EXAMPLE 132 	100 10 1 100 10 1 100 10 1	87 39 8 87 33 0 97 77 27		
EXAMPLE 133 	100 10 1	25 5 3		
EXAMPLE 141 	100 10 1	48 15 9		
EXAMPLE 78 	100 10 1	62 16 4		

EP 1 019 364 B1





PNU-number, Structure	PTP1B Inhib		PTP1 Inhib	
	PTP1B. Conc (uM)	PTP1B. % Inhib	PTP1N. Conc (uM)	PTP1N. % Inhib
EXAMPLE 83 	100 10 1	98 88 47		
EXAMPLE 84 	100 10 1	98 87 45		
EXAMPLE 85 	100 10 1	96 70 19		
EXAMPLE 86 	100 10 1	96 77 28		

TABLE 2 CONTINUED





PNU-number, Structure	PTP1B Inhib		PTP1 Inhib	
	PTP1B. Conc (uM)	PTP1B. % Inhib	PTP1N. Conc (uM)	PTP1N. % Inhib
EXAMPLE 87 	100 10 1	97 83 37		
EXAMPLE 137 	100 10 1	29 6 3		
EXAMPLE 79 	100 10 1	94 65 19		
EXAMPLE 134 	100 10 1	54 13 5		

TABLE 2 CONTINUED





PNU-number, Structure	PTP1B Inhib		PTP1N Inhib	
	PTP1B. Conc (uM)	PTP1B. % Inhib	PTP1N. Conc (uM)	PTP1N. % Inhib
EXAMPLE 88 	100 10 1	96 78 23		
EXAMPLE 89 	100 10 1	96 75 19		
EXAMPLE 90 	100 10 1	95 73 15		
EXAMPLE 127 	100 10 1	89 44 10		

TABLE 2 CONTINUED





PNU-number, Structure	PTP1B Inhib		PTP1 Inhib	
	PTP1B. Conc (uM)	PTP1B. % Inhib	PTP1N. Conc (uM)	PTP1N. % Inhib
EXAMPLE 135 	100 10 1	30 8 6		
EXAMPLE 91 	100 10 1	98 86 43		
EXAMPLE 92 	100 10 1	98 88 46		
EXAMPLE 93 	100 10 1	98 85 39		

TABLE 2 CONTINUED





PNU-number, Structure	PTP1B Inhib		PTP1N Inhib	
	PTP1B. Conc (uM)	PTP1B. % Inhib	PTP1N. Conc (uM)	PTP1N. % Inhib
EXAMPLE 94 	100 10 1	98 85 39		
EXAMPLE 95 	100 10 1	97 87 46		
EXAMPLE 142 	100 10 1	25 5 1		
EXAMPLE 80 	100 10 1	86 43 8		

TABLE 2 CONTINUED








PNU-number, Structure	PTP1B Inhib		PTP1 Inhib	
	PTP1B. Conc (μ M)	PTP1B. % Inhib	PTP1N. Conc (μ M)	PTP1N. % Inhib
EXAMPLE 96 	100 10 1	95 74 22		
EXAMPLE 143 	100 10 1	94 71 24		
EXAMPLE 144 	100 10 1	95 76 34		
EXAMPLE 145 	100 10 1	98 93 64		

TABLE 2 CONTINUED

PNU-number, Structure	PTP1B Inhib		PTP1 Inhib	
	PTP1B. Conc (μ M)	PTP1B. % Inhib	PTP1N. Conc (μ M)	PTP1N. % Inhib
EXAMPLE 138 	100 10 1	80 31 5		
EXAMPLE 136 			100 10 1	70 14 1

Claims

1. A compound of formula 1



I


wherein G^1 is

- a) $-R^2$, or
- b) $-NR^8 R^4$;

R^2 is

- a) $-C_1-C_{10}$ alkyl optionally substituted with one or two $-CO_2R^5$ bonded to the same or different carbon atoms or with one $-CO-NH_2$,

- b) -C₃-C₈ cycloalkyl optionally substituted with one -CO₂R⁵,
 c) -C₀-C₆ alkyl-phenyl optionally substituted with one or two -CO₂R⁵ bonded to the same or different carbon atoms or with -CH₂CH(CO₂R⁵)₂,
 d) -CH(R⁷)NHXR⁶, or
 e)



R³ is

- a) -C₁-C₁₂ alkyl, optionally substituted with one to three -O-, -S-, -N-, -O-C₁-C₄ alkyl, -S-C₁-C₄ alkyl, -O-G³, -S-G³, or -OH,
 b) -C₁-C₄ alkyl -C₃-C₆ cycloalkyl,
 c) -C₂-C₁₂ alkenyl,
 d) -C₃-C₁₂ alkynyl,
 e) -C₀-C₁₀ alkyl(G³)_n wherein alkyl is optionally substituted with one to three -O-, -S- or -N-, or
 f) -CH(CONH₂)C₁-C₁₂ alkyl;

R⁴ is

- a) -H,
 b) up to C₁₈ alkyl or alkenyl, or
 c) -C₀-C₆-alkyl-G³;

R⁵ is

- a) -H,
 b) -C₁-C₁₀ alkyl, or
 c) -C₁-C₅ alkyl-phenyl;

R⁶ is

- a) C₁-C₁₀ alkyl,
 b) C₀-C₆ alkyl-G³,
 c) C₁-C₆ alkyl CONH₂,
 d) C₁-C₆ alkyl NHCO₂R⁵,
 e) C₁-C₆ alkyl-OR⁵,
 f) C₁-C₆ alkyl-NHSO₂Me,
 g) C₁-C₆ alkyl-O- G³,
 h) C₁-C₆ alkyl-S- G³, or
 i) -C₁-C₆ alkyl-CO₂R⁵;

R⁷ is

- a) -H,
 b) -C₁-C₆ alkyl- G³,
 c) -C₁-C₆ alkyl-CO₂R⁵,
 d) C₁-C₆ alkyl CONH₂,
 e) C₁-C₆ alkyl NHCO₂R⁵,
 f) C₁-C₁₀ alkyl,
 g) up to C₁₀ cycloalkyl,
 h) -C₁-C₆ alkyl- SR⁵, or
 i) -C₁-C₆ alkyl- S(=O)R⁵;

R⁸ is

- a) C₀-C₆ alkyl-G³,
- b) CH(R⁷)CO₂R⁵,
- c) CH(R⁷)CH₂CO₂R⁵, or
- d) CH(R⁷)CONHCH₂CO₂R⁵;

G³ is

- a) phenyl substituted by zero (0) to three (3) R⁹, or
- b) naphthyl substituted by zero (0) to three (3) R⁹;

R⁹ is

- a) C₁-C₈ alkyl substituted by zero (0) to three (3) halo,
- b) C₂-C₈ alkenyl,
- c) OH,
- d) O-C₁-C₅ alkyl,
- e) O-C₀-C₅ alkyl-phenyl
- f) -(CH₂)_n-O-C₁-C₅ alkyl substituted by zero (0) to three (3) hydroxy,
- g) halo,
- h) NH₂,
- i) amino-C₁-C₅ alkyl,
- j) mono-or di-C₁-C₅ alkylamino,
- k) -C(O)-C₁-C₅ alkyl,
- l) -CHO,
- m) -C(O)-C₀-C₅ alkyl-phenyl
- n) -COOR⁵
- o) -CON(R⁵)₂,
- p) -C₃-C₇ cycloalkyl,
- q) -NO₂,
- r) -CN,
- s) -SO₃H,
- t) -SO₂N(R⁵)₂,
- u) -O[(CH₂)₂-O]_n-CH₃,
- v) -[CH₂-O]_n-C₁-C₃ alkyl,
- w) -NR⁵(CO)-NR⁵,
- x) -CF₃,
- y) -NR⁵(CO)C₁-C₅ alkyl
- z) -N(R⁵)-SO₂-R⁵,
- a1) -O-C(O)-R⁵,
- b1) -S(O)-R⁵,
- c1) -SR⁵, or
- d1) -SO₂-R⁵;

R¹⁰ is

- a) CO₂R⁵, or
- b) 5-tetrazolyl;


R¹¹ is

- a) H, or
- b) methyl;


X is -CO- or -SO₂- or -CO₂-; and
n is zero, one, two or three;

or a pharmaceutically acceptable salt thereof.

2. The compound according to claim 1 of formula III or IV



III




IV

wherein G¹, R³ and R⁵ are as defined in claim 1, and the chiral center depicted in III is in the (S) configuration.

3. The compound according to claim 2, wherein G¹ is

- a) -CH(CH₂phenyl)NHCO₂t-Bu,
- b) -CH(CH₂phenyl)NHCOC₁₋₃ alkyl-G³,
- c) -CH(CH₂phenyl)NHCOC₁₋₃ alkyl-CO₂R⁵, or
- d)



R³ is

- a) -C₅-C₆ alkyl, or
- b) -C₃-C₆ alkyl-phenyl;

R⁵ is -H; and

the configuration of the chiral center(s) is (S).

4. The compound according to any one of claims 1 to 3, selected from the group consisting of:


(S)-2-(Carboxymethoxy)-5-[2-[(3-carboxy-1-oxopropyl)amino]-3-oxo-3-(pentylamino)propyl]benzoic acid;
 5-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]-
 2-(carboxymethoxy)benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-2-[(dibenzylamino)carbonyl]amino]-3-oxo-3-(pentylamino)propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(3-phenylpropanoyl)amino]propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-2-[(2S)-2-[(3-carboxypropanoyl)amino]-3-phenylpropanoyl]amino]-3-oxo-
 3-(pentylamino)propyl]benzoic acid;
 5-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]-
 2-(carboxymethoxy)benzoic acid;
 5-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-(4-hydroxyphenyl)propanoyl]amino]-3-oxo-3-(pentylamino)
 propyl]-2-(carboxymethoxy)benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phenyl-2-[(2H-1,2,4-triazol-3-ylsulfanyl)
 acetyl]amino]propanoyl]amino]propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phenyl-2-[(2-(5-sulfanyl-1H-1,2,3,4-tetraazol-
 1-yl)acetyl]amino]propanoyl]amino]propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phenyl-2-[(2H-1,2,3-triazol-5-ylsulfanyl)

acetyl]amino]propanoyl)amino]propyl]benzoic acid;
 2-[4-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]-
 2-(2H-1,2,3,4-tetraazol-5-yl)phenoxy]acetic acid;
 5-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]-
 2-(2-methoxy-2-oxoethoxy)benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-2-[(2S)-2-[(2-(1H-indol-3-yl)acetyl]amino)-3-phenylpropanoyl]amino]-3-oxo-
 3-(pentylamino)propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phenyl-2-[(2-phenylacetyl)amino]propanoyl]
 amino]propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phenyl-2-[(4-phenylbutanoyl)amino]pro-
 panoyl]amino]propyl]benzoic acid;
 5-[(2S)-2-[(2S)-2-(acetylamino)-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]-2-(carboxymeth-
 oxy)benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-2-[(2S)-2-[(3-methoxypropanoyl)amino]-3-phenylpropanoyl]amino]-3-oxo-
 3-(pentylamino)propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-2-[(2S)-2-[(4-hydroxybutanoyl)amino]-3-phenylpropanoyl]amino]-3-oxo-
 3-(pentylamino)propyl]benzoic acid;
 5-[(2S,5S)-5-benzyl-13,13-dimethyl-4,7,11-trioxo-2-[(pentylamino)carbonyl]-12-oxa-3,6,10-triazatetradec-
 1-yl]-2-(carboxymethoxy)benzoic acid;
 5-[(2S,5S)-5-benzyl-4,7,11,11-tetraoxo-2-[(pentylamino)carbonyl]-11 λ 6-thia-3,6,10-triazadodec-1-yl]-
 2-(carboxymethoxy)benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-2-[(2S)-2-[(2-(3-hydroxyphenyl)acetyl]amino)-3-phenylpropanoyl]amino]-
 3-oxo-3-(pentylamino)propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-2-[(2S)-2-[(2-(4-hydroxyphenyl)acetyl]amino)-3-phenylpropanoyl]amino]-
 3-oxo-3-(pentylamino)propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-2-[(2S)-2-[(2-(4-methylphenyl)acetyl]amino)-3-phenylpropanoyl]amino]-3-oxo-
 3-(pentylamino)propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phenyl-2-[(2-[4-(trifluoromethyl)phenyl]acetyl]
 amino]propanoyl]amino]propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-2-[(2S)-2-[(2-(4-methoxyphenyl)acetyl]amino)-3-phenylpropanoyl]amino]-
 3-oxo-3-(pentylamino)propyl]benzoic acid;
 5-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]propanoyl]amino]-3-oxo-3-(pentylamino)propyl]-2-(car-
 boxymethoxy)benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phenyl-2-[(3-phenylpropanoyl)amino]pro-
 panoyl]amino]propyl]benzoic acid;
 2-(carboxymethoxy)-5-[(2S)-3-oxo-2-[(2R)-2-(2-oxo-1-pyrrolidinyl)-3-phenylpropanoyl]amino]-3-(pentylami-
 no)propyl]benzoic acid; and
 5-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-[(4-phenylbutyl)amino]
 propyl]-2-(carboxymethoxy)benzoic acid.

5. Use of a compound according to any one of claims 1 to 4, in the manufacture of a medicament useful for the treatment and/or prevention of Non-Insulin Dependent Diabetes Mellitus.

Patentansprüche

1. Eine Verbindung der Formel I




I

worin G^1 darstellt

- a) $-R^2$, oder
- b) $-NR^8R^4$;

R^2 darstellt

- a) $-C_1-C_{10}$ -Alkyl, gegebenenfalls mit einem oder zwei $-CO_2R^5$, die am gleichen oder an unterschiedlichen Kohlenstoffatomen gebunden sind, oder mit einem $-CO-NH_2$ substituiert,
- b) $-C_3-C_8$ -Cycloalkyl, gegebenenfalls mit einem $-CO_2R^5$ substituiert,
- c) $-C_0-C_6$ -Alkylphenyl, gegebenenfalls mit einem oder zwei $-CO_2R^5$, die am gleichen oder an verschiedenen Kohlenstoffatomen gebunden sind, oder mit $-CH_2CH(CO_2R^5)_2$ substituiert,
- d) $-CH(R^7)NHR^6$, oder
- e)



R^3 darstellt

- a) $-C_1-C_{12}$ -Alkyl, gegebenenfalls mit einem bis drei $-O-$, $-S-$, $-N-$, $-O-C_1-C_4$ -Alkyl, $-S-C_1-C_4$ -Alkyl, $-O-G^3$, $-S-G^3$ oder $-OH$,
- b) $-C_1-C_4$ -Alkyl- $-C_3-C_6$ -cycloalkyl,
- c) $-C_2-C_{12}$ -Alkenyl,
- d) $-C_3-C_{12}$ -Alkynyl,
- e) $-C_0-C_{10}$ -Alkyl(G^3) $_n$, worin Alkyl gegebenenfalls mit einem bis 3 $-O-$, $-S$ oder $-N-$ substituiert ist, oder
- f) $-CH(CONH_2)C_1-C_{12}$ -Alkyl;

R^4 darstellt

- a) $-H$,
- b) bis zu C_{18} -Alkyl oder Alkenyl, oder
- c) $-C_0-C_6$ -Alkyl- G^3 ;

R^5 darstellt

- a) $-H$,
- b) $-C-C_{10}$ -Alkyl, oder
- c) $-C_1-C_5$ -Alkylphenyl;

R^6 darstellt

- a) C_1-C_{10} -Alkyl,
- b) C_0-C_6 -Alkyl- G^3 ,
- c) C_1-C_6 -Alkyl- $CONH_2$,
- d) C_1-C_6 -Alkyl- $NHCO_2R^5$,
- e) C_1-C_6 -Alkyl- OR^5 ,
- f) C_1-C_6 -Alkyl- $NHSO_2Me$,
- g) C_1-C_6 -Alkyl- $O-G^3$,
- h) C_1-C_6 -Alkyl- $S-G^3$, oder
- i) $-C_1-C_6$ -Alkyl- CO_2R^5 ;

R^7 darstellt

- a) $-H$,
- b) $-C_1-C_6$ -Alkyl- G^3 ,

- c) $-C_1-C_6\text{-Alkyl-CO}_2R^5$,
- d) $C_1-C_6\text{-Alkyl-CONH}_2$,
- e) $C_1-C_6\text{-Alkyl-NHCO}_2R^5$,
- f) $C_1-C_{10}\text{-Alkyl}$
- g) bis zu $C_{10}\text{-Cycloalkyl}$,
- h) $-C_1-C_6\text{-Alkyl-SR}^5$, oder
- i) $-C_1-C_6\text{-Alkyl-S(=O)R}^5$;

R^8 darstellt

- a) $C_0-C_6\text{-Alkyl-G}^3$,
- b) $CH(R^7)CO_2R^5$,
- c) $CH(R^7)CH_2CO_2R^5$, oder
- d) $CH(R^7)CONHCH_2CO_2R^5$;

G^3 darstellt

- a) Phenyl, das mit Null (0) bis drei (3) R^9 substituiert ist, oder
- b) Naphthyl, das mit Null (0) bis drei (3) R^9 substituiert ist;

R^9 darstellt

- a) $C_1-C_8\text{-Alkyl}$, substituiert mit Null (0) bis drei (3) Halogen,
- b) $C_2-C_8\text{-Alkenyl}$,
- c) OH,
- d) $O-C_1-C_5\text{-Alkyl}$
- e) $O-C_0-C_5\text{-Alkylphenyl}$
- f) $-(CH_2)_n-O-C_1-C_5\text{-Alkyl}$, substituiert mit Null (0) bis drei (3) Hydroxy,
- g) $-(CH_2)_n-O-C_2-C_7\text{-Alkenyl}$, substituiert mit Null (0) bis drei (3) Hydroxy,
- h) Halogen,
- i) NH_2 ,
- j) Amino- $C_1-C_5\text{-Alkyl}$,
- k) Mono- oder Di- $C_1-C_5\text{-Alkylamino}$
- l) $-C(O)-C_1-C_5\text{-Alkyl}$
- m) $-CHO$,
- n) $-C(O)-C_0-C_5\text{-Alkylphenyl}$
- o) $-COOR^5$,
- p) $-CON(R^5)_2$,
- q) $-C_3-C_7\text{-Cycloalkyl}$,
- r) $-NO_2$,
- s) $-CN$,
- t) $-SO_3H$,
- u) $-SO_2N(R^5)_2$,
- v) $-O[(CH_2)_2-O]_n-CH_3$,
- w) $-[CH_2-O]_n-C_1-C_3\text{-Alkyl}$,
- x) $-NR^5(CO)-NR^5$,
- y) $-CF_3$,
- z) $-NR^5(CO)C_1-C_5\text{-Alkyl}$
- a1) $-N(R^5)-SO_2-R^5$,
- b1) $-O-C(O)-R^5$,
- c1) $-S(O)-R^5$,
- d1) $-SR^5$, oder
- d1) $-SO_2-R^5$;

R^{10} darstellt


- a) CO_2R^5 , oder
- b) 5-Tetrazolyl;

R¹¹ darstellt


- a) H oder
b) Methyl;

X ist -CO- oder -SO₂- oder -CO₂-; und
n ist null, eins, zwei oder drei;
oder ein pharmazeutisch verträgliches Salz davon.

2. Die Verbindung gemäß Anspruch 1 der Formel III oder IV



III




IV

worin G¹, R³ und R⁵ wie in Anspruch 1 definiert sind und worin das in Formel III dargestellte Chiralitätszentrum in der (S)-Konfiguration vorliegt.

3. Die Verbindung gemäß Anspruch 2, worin
G¹ darstellt

- a) -CH(CH₂Phenyl)NHCO₂t-Bu,
b) -CH(CH₂Phenyl)NHCOC₁-C₃-alkyl-G³,
c) -CH(CH₂Phenyl)NHCOC₁-C₃-alkyl-CO₂R⁵, oder
d)



R³ darstellt

- a) -C₅-C₆-Alkyl oder
b) -C₃-C₆-Alkylphenyl;

R⁵ ist -H, und
die Konfiguration des Chiralitätszentrums bzw. der Chiralitätszentren (S) ist.

4. Die Verbindung gemäß einem der Ansprüche 1 bis 3, die aus der Gruppe ausgewählt ist, bestehend aus:


(S)-2-(Carboxymethoxy)-5-[2-[(3-carboxy-1-oxopropyl)amino]-3-oxo-3-(pentylamino)propyl]benzoesäure;
5-[(2S)-2-(((2S)-2-[Tert-butoxycarbonyl]amino)-3-phenylpropanoyl)amino)-3-oxo-3-(pentylamino)propyl]-
2-(carboxymethoxy)benzoesäure;
2-(Carboxymethoxy)-5-[(2S)-2-[(dibenzylamino)carbonyl]amino]-3-oxo-3-(pentylamino)propyl]benzoesäure;
2-(Carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(3-phenylpropanoyl)amino]propyl]benzoesäure;
2-(Carboxymethoxy)-5-[(2S)-2-(((2S)-2-[3-carboxypropanoyl]amino)-3-phenylpropanoyl)amino)-3-oxo-
3-(pentylamino)propyl]benzoesäure;
5-[(2S)-2-(((2S)-2-[Tert-butoxycarbonyl]amino)-3-phenylpropanoyl)amino)-3-oxo-3-(pentylamino)propyl]-

2-(carboxymethoxy)benzoesäure;
 5-[(2S)-2-[(2S)-2-[(Tert-butoxycarbonyl)amino]-3-(4-hydroxyphenyl)propanoyl]amino]-3-oxo-3-(pentylamino)propyl]-2-(carboxymethoxy)benzoesäure;
 2-(Carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phenyl-2-[[2-(4H-1,2,4-triazol-3-ylsulfanyl)acetyl]amino]propanoyl]amino]propyl]benzoesäure;
 2-(Carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phenyl-2-[[2-(5-sulfanyl-1H-1,2,3,4-tetraazol-1-yl)acetyl]amino]propanoyl]amino]propyl]benzoesäure;
 2-(Carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phenyl-2-[[2-(1H-1,2,3-triazol-5-ylsulfanyl)acetyl]amino]propanoyl]amino]propyl]benzoesäure;
 2-[4-[(2S)-2-[(2S)-2-[(Tert-butoxycarbonyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]-2-(2H-1,2,3,4-tetraazol-5-yl)phenoxy]essigsäure;
 5-[(2S)-2-[(2S)-2-[(Tert-butoxycarbonyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]-2-(2-methoxy-2-oxoethoxy)benzoesäure;
 2-(Carboxymethoxy)-5-[(2S)-2-[(2S)-2-[[2-(1H-indol-3-yl)acetyl]amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoesäure;
 2-(Carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phenyl-2-[(2-phenylacetyl)amino]propanoyl]amino]propyl]benzoesäure;
 2-(Carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phenyl-2-[(4-phenylbutanoyl)amino]propanoyl]amino]propyl]benzoesäure;
 5-[(2S)-2-[(2S)-2-(Acetyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]-2-(carboxymethoxy)benzoesäure;
 2-(Carboxymethoxy)-5-[(2S)-2-[(2S)-2-[(3-methoxypropanoyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoesäure;
 2-(Carboxymethoxy)-5-[(2S)-2-[(2S)-2-[(4-hydroxybutanoyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoesäure;
 5-[(2S,5S)-5-Benzyl-13,13-dimethyl-4,7,11-trioxo-2-[(pentylamino)carbonyl]-12-oxa-3,6,10-triazatetradec-1-yl]-2-(carboxymethoxy)benzoesäure;
 5-[(2S,5S)-5-Benzyl-4,7,11,11-tetraoxo-2-[(pentylamino)carbonyl]-11 λ 6-thia-3,6,10-triazadodec-1-yl]-2-(carboxymethoxy)benzoesäure;
 2-(Carboxymethoxy)-5-[(2S)-2-[(2S)-2-[[2-(3-hydroxyphenyl)acetyl]amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoesäure;
 2-(Carboxymethoxy)-5-[(2S)-2-[(2S)-2-[[2-(4-hydroxyphenyl)acetyl]amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoesäure;
 2-(Carboxymethoxy)-5-[(2S)-2-[(2S)-2-[[2-(4-methylphenyl)acetyl]amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoesäure;
 2-(Carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phenyl-2-[[2-(4-(trifluormethyl)phenyl)acetyl]amino]propanoyl]amino]propyl]benzoesäure;
 2-(Carboxymethoxy)-5-[(2S)-2-[(2S)-2-[[2-(4-methoxyphenyl)acetyl]amino]-3-phenylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoesäure;
 5-[(2S)-2-[(2S)-2-[(Tert-butoxycarbonyl)amino]propanoyl]amino]-3-oxo-3-(pentylamino)propyl]-2-(carboxymethoxy)benzoesäure;
 2-(Carboxymethoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phenyl-2-[(3-phenylpropanoyl)amino]propanoyl]amino]propyl]benzoesäure;
 2-(Carboxymethoxy)-5-[(2S)-3-oxo-2-[(2R)-2-(2-oxo-1-pyrrolidinyl)-3-phenylpropanoyl]amino]-3-(pentylamino)propyl]benzoesäure; und
 5-[(2S)-2-[(2S)-2-[(Tert-butoxycarbonyl)amino]-3-phenylpropanoyl]amino]-3-oxo-3-[(4-phenylbutyl)amino]propyl]-2-(carboxymethoxy)benzoesäure.

5. Verwendung einer Verbindung gemäß einem der Ansprüche 1 bis 4 bei der Herstellung eines Arzneimittels, das zur Behandlung und/oder Prävention von nicht-Insulin-abhängiger Diabetes Mellitus geeignet ist.

Revendications

1. Composé de formule I




dans laquelle G¹ représente :

- a) -R² ou
- b) -NR⁸R⁴ ;

R² représente :

- a) -alkyle en C₁-C₁₀ éventuellement substitué par un ou deux -CO₂R⁵ liés au même atome de carbone ou à des atomes de carbone différents ou par un -CO-NH₂,
- b) -cycloalkyle en C₃-C₈ éventuellement substitué par un -CO₂R⁵,
- c) -alkyle en C₀-C₆-phényle éventuellement substitué par un ou deux -CO₂R⁵ liés au même atome de carbone ou à des atomes de carbone différents ou par -CH₂CH(CO₂R⁵)₂,
- d) -CH(R⁷)NHR⁶ ou
- e)



R³ représente :

- a) -alkyle en C₁-C₁₂, éventuellement substitué par un à trois -O-, -S-, -N-, -O-alkyle en C₁-C₄, -S-alkyle en C₁-C₄, -O-G³, -S-G³ ou -OH,
- b) -alkyle en C₁-C₄-cycloalkyle en C₃-C₆,
- c) -alcényle en C₂-C₁₂,
- d) -alcynyle en C₃-C₁₂,
- e) -alkyle en C₀-C₁₀ (G³)_n dans lequel l'alkyle est éventuellement substitué par un à trois -O-, -S- ou -N-, ou
- f) -CH(CONH₂)alkyle en C₁-C₁₂ ;

R⁴ représente :

- a) -H,
- b) alkyle ou alcényle ayant jusqu'à C₁₈ ou
- c) -alkyle en C₀-C₆-G³ ;

R⁵ représente :

- a) -H,
- b) -alkyle en C₁-C₁₀ ou
- c) -alkyle en C₁-C₅-phényle ;

R⁶ représente :

- a) alkyle en C₁-C₁₀,
- b) alkyle en C₀-C₆-G³,
- c) alkyle en C₁-C₆ CONH₂,
- d) alkyle en C₁-C₆ NHCO₂R⁵,
- e) alkyle en C₁-C₆-OR⁵,
- f) alkyle en C₁-C₆-NHSO₂Me,
- g) alkyle en C₁-C₆-O-G³,
- h) alkyle en C₁-C₆-S-G³ ou
- i) -alkyle en C₁-C₆-CO₂R⁵ ;

R⁷ représente :

- a) -H,
- b) -alkyle en C₁-C₆-G³,
- c) -alkyle en C₁-C₆-CO₂R⁵,
- d) alkyle en C₁-C₆ CONH₂,
- e) alkyle en C₁-C₆ NHCO₂R⁵,
- f) alkyle en C₁-C₁₀,
- g) cycloalkyle jusqu'à C₁₀,
- h) -alkyle en C₁-C₆-SR⁵ ou
- i) -alkyle en C₁-C₆-S(=O)R⁵ ;

R⁸ représente :

- a) alkyle en C₀-C₆-G³,
- b) CH (R⁷) CO₂R⁵,
- c) CH(R⁷)CH₂CO₂R⁵ ou
- d) CH(R⁷)CONHCH₂CO₂R⁵ ;

G³ représente :

- a) phényle substitué par de zéro (0) à trois (3) R⁹, ou
- b) naphthyle substitué par de zéro (0) à trois (3) R⁹ ;

R⁹ représente :

- a) alkyle en C₁-C₈ substitué par de zéro (0) à trois (3) halo,
- b) alcényle en C₂-C₈
- c) OH,
- d) O-alkyle en C₁-C₅,
- e) O-alkyle en C₀-C₅-phényle,
- f) -(CH₂)_n-O-alkyle en C₁-C₅ substitué par de zéro (0) à trois (3) hydroxy,
- g) halo,
- h) NH₂,
- i) amino-alkyle en C₁-C₅,
- j) mono- ou di-alkylamino en C₁-C₅,
- k) -C(O)-alkyle en C₁-C₅,
- l) -CHO,
- m) -C(O)-alkyle en C₀-C₅-phényle,
- n) -COOR⁵,
- o) -CON(R⁵)₂,
- p) -cycloalkyle en C₃-C₇,
- q) -NO₂,
- r) -CN,
- s) -SO₃H,
- t) -SO₂N(R⁵)₂,
- u) -O[(CH₂)₂-O]_n-CH₃,

- v) $-\text{[CH}_2\text{-O]}_n\text{-alkyle en C}_1\text{-C}_3$,
 w) $-\text{NR}^5(\text{CO})\text{-NR}^5$,
 x) $-\text{CF}_3$,
 y) $-\text{NR}^5(\text{CO})\text{ alkyle en C}_1\text{-C}_5$,
 z) $-\text{N(R}^5\text{)-SO}_2\text{-R}^5$,
 a1) $-\text{O-C(O)-R}^5$,
 b1) $-\text{S(O)-R}^5$,
 c1) $-\text{SR}^5$ ou
 d1) $-\text{SO}_2\text{-R}^5$;


R^{10} représente :

- a) CO_2R^5 ou
 b) 5-tétrazolyne ;
 R^{11} représente :


- a) H ou
 b) méthyle ;

X représente $-\text{CO}-$ ou $-\text{SO}_2-$ ou $-\text{CO}_2-$; et
 n vaut zéro, un, deux ou trois ;
 ou sel pharmaceutiquement acceptable de celui-ci.

2. Composé selon la revendication 1, de formule III ou IV



III




IV

dans laquelle G^1 , R^3 et R^5 sont tels que définis dans la revendication 1, et le centre chiral représenté en III est en configuration (S).

3. Composé selon la revendication 2, dans lequel G^1 représente :

- a) $-\text{CH}(\text{CH}_2\text{phényl})\text{NHCO}_2\text{t-Bu}$,
 b) $-\text{CH}(\text{CH}_2\text{phényl})\text{NHCO alkyle en C}_1\text{-C}_3\text{-G}^3$,
 c) $-\text{CH}(\text{CH}_2\text{phényl})\text{NHCO alkyle en C}_1\text{-C}_3\text{-CO}_2\text{R}^5$ ou
 d)



R³ représente :

- a) -alkyle en C₅-C₆ ou
- b) -alkyle en C₃-C₆-phényle ;

R⁵ représente -H ; et

la configuration du (des) centre(s) chiral(aux) est (S).

4. Composé selon l'une quelconque des revendications 1 à 3, choisi dans le groupe constitué par :

l'acide (S)-2-(carboxyméthoxy)-5-[2-[(3-carboxy-1-oxopropyl)amino]-3-oxo-3-(pentylamino)propyl]benzoïque ;
 l'acide 5-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)-amino]-3-phénylpropanoyl]amino]-3-oxo-3-(pentylamino)-propyl]-2-(carboxyméthoxy)benzoïque ;
 l'acide 2-(carboxyméthoxy)-5-[(2S)-2-[(dibenzylamino)carbonyl]amino]-3-oxo-3-(pentylamino)propyl]benzoïque ;
 l'acide 2-(carboxyméthoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(3-phénylpropanoyl)amino]propyl]benzoïque ;
 l'acide 2-(carboxyméthoxy)-5-[(2S)-2-[(2S)-2-[(3-carboxypropanoyl)amino]-3-phénylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoïque ;
 l'acide 5-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)-amino]-3-phénylpropanoyl]amino]-3-oxo-3-(pentylamino)-propyl]-2-(carboxyméthoxy)benzoïque ;
 l'acide 5-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)-amino]-3-(4-hydroxyphényl)propanoyl]amino]-3-oxo-3-(pentylamino)propyl]-2-(carboxyméthoxy)benzoïque ;
 l'acide 2-(carboxyméthoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phényl-2-[[2-(4H-1,2,4-triazol-3-yl)sulfanyl]acétyl]amino]propanoyl]amino]propyl]benzoïque ;
 l'acide 2-(carboxyméthoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phényl-2-[[2-(5-sulfanyl-1H-1,2,3,4-tétrazol-1-yl)acétyl]amino]propanoyl]amino]-propyl]benzoïque ;
 l'acide 2-(carboxyméthoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phényl-2-[[2-(1H-1,2,3-triazol-5-yl)sulfanyl]acétyl]amino]propanoyl]amino]propyl]benzoïque ;
 l'acide 2-[4-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)amino]-3-phénylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]-2-(2H-1,2,3,4-tétrazol-5-yl)-phénoxy]acétique ;
 l'acide 5-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)-amino]-3-phénylpropanoyl]amino]-3-oxo-3-(pentylamino)-propyl]-2-(2-méthoxy-2-oxoéthoxy)benzoïque ;
 l'acide 2-(carboxyméthoxy)-5-[(2S)-2-[(2S)-2-[[2-(1H-indol-3-yl)acétyl]amino]-3-phénylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoïque ;
 l'acide 2-(carboxyméthoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phényl-2-[(2-phénylacétyl)-amino]propanoyl]amino]propyl]benzoïque ;
 l'acide 2-(carboxyméthoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phényl-2-[(4-phénylbutanoyl)-amino]propanoyl]amino]propyl]benzoïque ;
 l'acide 5-[(2S)-2-[(2S)-2-(acétylamino)-3-phénylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]-2-(carboxyméthoxy)benzoïque ;
 l'acide 2-(carboxyméthoxy)-5-[(2S)-2-[(2S)-2-[(3-méthoxypropanoyl)amino]-3-phénylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoïque ;
 l'acide 2-(carboxyméthoxy)-5-[(2S)-2-[(2S)-2-[(4-hydroxybutanoyl)amino]-3-phénylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoïque ;
 l'acide 5-[(2S,5S)-5-benzyl-13,13-diméthyl-4,7,11-trioxo-2-[(pentylamino)carbonyl]-12-oxa-3,6,10-triazatétradéc-1-yl]-2-(carboxyméthoxy)benzoïque ;
 l'acide 5-[(2S,5S)-5-benzyl-4,7,11,11-tétraoxo-2-[(pentylamino)carbonyl]-11λ⁶-thia-3,6,10-triazadodéc-1-yl]-2-(carboxyméthoxy)benzoïque ;
 l'acide 2-(carboxyméthoxy)-5-[(2S)-2-[(2S)-2-[[2-(3-hydroxyphényl)acétyl]amino]-3-phénylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoïque ;
 l'acide 2-(carboxyméthoxy)-5-[(2S)-2-[(2S)-2-[[2-(4-hydroxyphényl)acétyl]amino]-3-phénylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoïque ;
 l'acide 2-(carboxyméthoxy)-5-[(2S)-2-[(2S)-2-[[2-(4-méthylphényl)acétyl]amino]-3-phénylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoïque ;
 l'acide 2-(carboxyméthoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-[(2S)-3-phényl-2-[[2-(4-(trifluorométhyl)phényl)acétyl]amino]propanoyl]amino]propyl]benzoïque ;
 l'acide 2-(carboxyméthoxy)-5-[(2S)-2-[(2S)-2-[[2-(4-méthoxyphényl)acétyl]amino]-3-phénylpropanoyl]amino]-3-oxo-3-(pentylamino)propyl]benzoïque ;
 l'acide 5-[(2S)-2-[(2S)-2-[(tert-butoxycarbonyl)-amino]propanoyl]amino]-3-oxo-3-(pentylamino)propyl]-2-

(carboxyméthoxy)benzoïque ;
 l'acide 2-(carboxyméthoxy)-5-[(2S)-3-oxo-3-(pentylamino)-2-(((2S)-3-phényl-2-[(3-phénylpropanoyl)-amino]propanoyl)amino)propyl]benzoïque ;
 l'acide 2-(carboxyméthoxy)-5-[(2S)-3-oxo-2-(((2R)-2-(2-oxo-1-pyrrolidinyl)-3-phénylpropanoyl)amino)-3-(pentylamino)propyl]benzoïque ; et
 l'acide 5-[(2S)-2-(((2S)-2-[(tert-butoxycarbonyl)-amino]-3-phénylpropanoyl)amino)-3-oxo-3-[(4-phénylbutyl)-amino]propyl]-2-(carboxyméthoxy)benzoïque.

5. Utilisation d'un composé selon l'une quelconque des revendications 1 à 4, dans la fabrication d'un médicament utile pour le traitement et/ou la prévention du diabète sucré non insulino-dépendant.